



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

09TT 2900 54 2



LANE MEDICAL LIBRARY STAMFORD

V 141.2
P56
1921

Gelsemium *Meningeal*
Neuralgia

LANE

MEDICAL



LIBRARY


Gift
Dr. Howard W. Fleming

1925

L. B. Burgin

200 3/4 Lynch St

San Francisco.



Epitome of the Pharmacopeia of the United States and the National Formulary. With Comments

**Prepared for the Use of Physicians Under
Authorization of the Council on Pharmacy and
Chemistry of the American Medical Association
by a Committee Consisting of the Following:**

- A. W. HEWLETT, M.D.**, Professor of Medicine, Leland Stanford Junior University.
- TORALD SOLLMANN, M.D.**, Professor of Pharmacology and Materia Medica, Medical Department, Western Reserve University.
- M. I. WILBERT, PH.M., PHAR.D.**, Assistant in the Division of Pharmacology, Hygienic Laboratory, U. S. Public Health Service.
- W. A. PUCKNER, PHAR.D.**, Secretary of the Council on Pharmacy and Chemistry.

...A.M.A. LIBRARY

PERMISSION TO USE FOR COMMENT PARTS OF THE NATIONAL FORMULARY HAS BEEN GRANTED BY THE COUNCIL OF THE AMERICAN PHARMACEUTICAL ASSOCIATION.]

W. A. R. 1990

COPYRIGHT, 1921
AMERICAN MEDICAL ASSOCIATION

P56
1921

PREFACE TO FIRST EDITION

The Pharmacopeia of the United States and the National Formulary are now recognized by federal and state laws as standards for drugs and their preparations. Physicians who prescribe Pharmacopeial or National Formulary drugs or preparations, therefore, are more likely to obtain a good and constant quality than if they prescribe unofficial articles, i. e., articles that have no legalized standard. Hence medical men are interested in knowing what drugs and preparations are included in these two books of standards. Both the Pharmacopeia and the National Formulary, however, contain a vast amount of technical information, of value to pharmacists but of little interest to physicians. THE JOURNAL of the American Medical Association in 1907 issued the "Physicians' Manual of the U. S. Pharmacopeia and National Formulary," a book designed to meet the needs of physicians in this respect. The fact that this book did meet a real need is evidenced by the continued demand for it.

The present "Epitome of the U. S. Pharmacopeia and National Formulary" takes the place of that book, but is based on the new (the ninth) revision of the Pharmacopeia and the new (the fourth) edition of the National Formulary. It has been prepared under the direction of a committee appointed by the Council on Pharmacy and Chemistry of the American Medical Association. The aim has been to include all the matter in the Pharmacopeia and National Formulary which is likely to be of interest to physicians. This, it is believed, consists of the official titles and their abbreviations; synonyms; brief definitions; when necessary, concise descriptions of the physical properties; and dosage. Since both the Pharmacopeia and the National Formulary include many drugs and preparations which are irrational, superfluous

97538

PREFACE

worthless, in many cases the medical members of the committee have added brief comments to aid a discriminating selection of therapeutic agents.

In this epitome drugs are arranged alphabetically by Pharmacopeial or National Formulary titles; preparations of drugs are placed under the drugs themselves. Thus *Tinctura Digitalis* is placed not among the Ts, but among the Ds under *Digitalis*. Names of drugs are in black-face type; names of preparations are in light-face italics. An index is added to assist in ready reference.

Statements of composition refer in some cases to percentage by volume, in others to percentage by weight, and in still others to percentage by weight-volume—that is, the weight of a given chemical contained (or the weight of a given vegetable drug represented by extractives) in 100 Cc. of the preparation. It has not been thought worth while to indicate whether the statement in each case is by volume, weight or weight-volume, since the differences are too small to have any therapeutic importance. Statements of alcoholic content refer to percentage by volume and are approximate only. In a few instances in which the quantity of alcohol in a dose is small, or in which the preparation is intended for external use, the alcohol content is not stated.

Dosage statements, when followed by the letters "U. S. P." or "N. F.," are taken from these books, respectively, and are understood to be the average adult doses. In some cases in which this statement was felt to be inadequate or unsatisfactory, further suggestions as to dosage have been added.

PREFACE TO SECOND EDITION

The text of this edition is the same as that of the first, except that a few typographical errors which were in the first edition have been corrected.

The new edition differs from the first in the following particulars :

In the first edition, a star indicates that the drug is included in the second (1916) edition of Useful Drugs. To bring the present edition up to date, the star (*) here indicates that the drug is included in the fifth (1921) edition of Useful Drugs.

In compliance with the opinion expressed by some that a more complete index would be helpful, the index to the second edition has been materially extended. It is hoped that this will help to familiarize those who purchase the book with its contents.

TABLES OF METRIC AND APOTHECARIES' WEIGHTS AND MEASURES

METRIC MEASURES OF WEIGHT

- 1 Milligram (1 mg.)=0.001 gram.
- 1 Centigram=0.01 gram.
- 1 Decigram=0.1 gram.
- 1 Gram (1 Gm.)=1.*
- 1 Decagram=10 grams.
- 1 Hektogram=100 grams.
- 1 Kilogram (1 kg.)=1000 grams.†
- * The weight of 1 cubic centimeter of water at 4 C.
- † The weight of 1 cubic decimeter of water at 4 C.

METRIC FLUID MEASURE

- 1 Milliliter (1 cubic centimeter, 1 Cc.)=0.001 cubic decimeter.
- 1 Centiliter (10 Cc.)=0.01 cubic decimeter.
- 1 Deciliter (100 Cc.)=0.1 cubic decimeter.
- 1 Liter (1000 Cc.)=1 cubic decimeter.

APOTHECARIES' WEIGHT

- 20 Grains=1 scruple (℥).
- 3 Scruples=1 drachm (℥).
- 8 Drachms=1 ounce (℥).
- 12 Ounces=1 pound (lb).

APOTHECARIES' FLUID MEASURE

- 60 Minims=1 fluidrachm (℥).
- 8 Fluidrachms=1 fluidounce (℥).
- 16 Fluidounces=1 pint (O).

EQUIVALENTS OF APOTHECARIES' WEIGHT IN METRIC

- 1 Grain=0.06479895 Gm.
- 1 Drachm=3.8879369 Gm.
- 1 Ounce=31.1034956 Gm.
- 1 Pound=373.241948 Gm.

EQUIVALENTS OF APOTHECARIES' FLUID MEASURE IN METRIC

- 1 Minim=0.06161 Cc.
- 1 Fluidrachm=3.6966 Cc.
- 1 Fluidounce=29.573 Cc.
- 1 Pint=473.17 Cc.

EQUIVALENTS OF METRIC WEIGHT IN APOTHECARIES'

- 0.001 Gm. (1 milligram, 1 mg.)=0.01543 grain.
- 0.01 Gm. (1 centigram)=0.15432 grain.
- 0.1 Gm. (1 decigram)=1.54324 grains.
- 1 Gm. (1 gram)=15.4324 grains.
- 10 Gm. (1 decagram)=2 drachms, 34.324 grains.
- 100 Gm. (1 hectogram)=3 ounces, 1 drachm, 43.24 grains.
- 1000 Gm. (1 kilogram, 1 kg.)=2 pounds, 8 ounces, 52.4 grains.

EQUIVALENTS OF METRIC FLUID MEASURE IN APOTHECARIES'

1 Cc.=16.231 minims.
10 Cc.=2 fluidrachms, 42.311 minims.
100 Cc.=3 fluidounces, 3 fluidrachms, 3.11 minims.
1000 Cc. (1 Liter)=2 pints, 1 fluidounce, 6 fluidrachms, 31.1 minims.

METRIC AND APOTHECARIES' EQUIVALENTS (WEIGHTS)

Gm.	Grains	Grains	Gm.
1 =	15.4324	1 =	0.06480
2 =	30.8647	2 =	0.12960
3 =	46.2971	3 =	0.19440
4 =	61.7294	4 =	0.25920
5 =	77.1618	5 =	0.32399
6 =	92.5941	6 =	0.38879
7 =	108.0265	7 =	0.45359
8 =	123.4589	8 =	0.51839
9 =	138.8912	9 =	0.58319

METRIC AND APOTHECARIES' EQUIVALENTS (VOLUME)

Cc.	Minims	Minims	Cc.
1 =	16.231	1 =	0.061610
2 =	32.462	2 =	0.123220
3 =	48.693	3 =	0.184831
4 =	64.924	4 =	0.246441
5 =	81.156	5 =	0.308051
6 =	97.387	6 =	0.369661
7 =	113.618	7 =	0.431271
8 =	129.849	8 =	0.492882
9 =	146.080	9 =	0.554492

CENTIGRADE AND FAHRENHEIT THERMOMETRIC EQUIVALENTS

C.°	F.°	C.°	F.°
-40	-40	65	149
-30	-22	70	158
-20	-4	75	167
-10	14	80	176
0	32	85	185
5	41	90	194
10	50	95	203
15	59	100	212
20	68	110	230
25	77	120	248
30	86	130	266
35	95	140	284
40	104	150	302
45	113	200	392
50	122	250	482
55	131	300	572
60	140	350	662

CLINICAL TABLE OF CENTIGRADE AND FAHRENHEIT
THERMOMETRIC EQUIVALENTS

C.°	F.°	C.°	F.°
36.0	96.80	39.0	102.20
36.2	97.16	39.1	102.38
36.3	97.34	39.2	102.56
36.4	97.52	39.3	102.74
36.5	97.70	39.4	102.92
36.6	97.88	39.5	103.10
36.7	98.06	39.6	103.28
36.8	98.24	39.7	103.46
36.9	98.42	39.8	103.64
37.0	98.60	39.9	103.82
37.1	98.78	40.0	104.00
37.2	98.96	40.1	104.18
37.3	99.14	40.2	104.36
37.4	99.32	40.3	104.54
37.5	99.50	40.4	104.72
37.6	99.68	40.5	104.90
37.7	99.86	40.6	105.08
37.8	100.04	40.8	105.44
37.9	100.22	41.0	105.80
38.0	100.40	41.2	106.16
38.1	100.58	41.4	106.52
38.2	100.76	41.6	106.88
38.3	100.94	41.8	107.24
38.4	101.12	42.0	107.60
38.5	101.30	42.2	107.96
38.6	101.48	42.4	108.32
38.7	101.66	42.6	108.68
38.8	101.84	42.8	109.04
38.9	102.02	43.0	109.40

Epitome of the U. S. Pharmacopeia and National Formulary

[STARS (★) DESIGNATE PREPARATIONS DESCRIBED IN
"USEFUL DRUGS," A BOOK PREPARED TO FURTHER A
MORE RATIONAL MATERIA MEDICA AND PUBLISHED BY
THE AMERICAN MEDICAL ASSOCIATION, PRICE 50 CENTS]

Absinthium (Absinth.), Absinthium, N. F. (Wormwood,
Madderwort, Vermuth).

Leaves and flowering tops.

ACTION AND USES: Obsolete aromatic bitter; without
advantage over gentian or other bitter drugs.

***Acacia (Acac.), Acacia, U. S. P.** (Gum Arabic, Gum Sene-
gal).

A gum occurring in tears, fragments or powder. Slowly and
almost completely soluble in water and insoluble in alcohol. Incom-
patible with sodium borate, lead acetate and ferric salts.

ACTION AND USES: Demulcent; chiefly as a vehicle to
suspend insoluble substances in aqueous mixtures.

***Mucilago Acaciae (Mucil. Acac.), Mucilage of Acacia,**
U. S. P.—Acacia (35%) in water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Syrupus Acaciae (Syr. Acac.), Syrup of Acacia, U. S. P.
Acacia (10%) in sugar and water.

ACETA, VINEGARS.—A vinegar is a solution of vegetable
principles either in diluted acetic acid or in a mixture
of diluted acetic acid and alcohol. For *Acetum Aromaticum*,
see under *Oleum Lavandulae*; for *Acetum Opii* and
Acetum Scillae, see under *Opium* and *Scilla*, respectively.

***Acetanilidum (Acetanil.), Acetanilid, U. S. P.** (Acetanilide,
Antifebrin).— $C_6H_5NH.CH_3CO$.

A white, odorless, crystalline powder, with a slightly burning
taste. Slightly soluble in water, (1:190) very soluble in alcohol
(1:3.4) and in chloroform (1:3.7). Incompatible with spirit of
nitrous ether and forms pasty masses when triturated with chloral
or antipyrin.

ACTION AND USES: Analgesic, antipyretic and, in
sive doses, a cardiac depressant. Used particu-

relieve headache and neuralgic pains. A habit-forming drug. If used over a long period of time, it may produce a severe anemia.

DOSAGE: 0.2 Gm. or 3 grains (U. S. P.) in powders which may be placed in capsules, or tablets which should be crushed before swallowing. It is well to begin with 0.1 Gm. or about $1\frac{1}{2}$ grains, and to repeat cautiously.

Pulvis Acetanilidi Compositus (*Pulv. Acetanil. Co.*), *Compound Acetanilid Powder, N. F.*—Acetanilid (70%), caffeine (10%) and sodium bicarbonate (20%).

USES: Irrational acetanilid preparation. The caffeine does not diminish the toxicity, nor does the bicarbonate increase the solubility of acetanilid, as was at one time supposed.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Acetonum (Aceton.), Acetone, U. S. P. (Dimethyl-Ketone).

A colorless liquid, with an ethereal odor and a pungent sweetish taste. Miscible with water, alcohol, ether and chloroform.

ACTION AND USES: Pharmaceutic solvent.

***Acetphenetidinum (Acetphen.), Acetphenetidin, U. S. P. (Phenacetin).**— $C_6H_5(OC_2H_5).NH.CH_2CO$.

White, odorless, slightly bitter, crystalline scales or powder. Very slightly soluble in water (1:1310), soluble in alcohol (1:15) and in chloroform (1:14).

ACTION AND USES: Analgesic, antipyretic and, in excessive doses, a cardiac depressant. Used particularly for the relief of headache and neuralgic pains and in the treatment of mild fevers. A habit-forming drug. If used for a long period of time, it may produce a severe anemia.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.), in powders or capsules. If small doses fail to relieve headache, larger doses are also usually ineffective.

***Acidum Aceticum (Acid. Acet.), Acetic Acid, U. S. P.**—Acetic acid CH_3COOH (about 36.5%).

ACTION AND USES: Caustic and rubefacient. In the form of vinegar, is antidote to alkalis.

Acidum Aceticum Dilutum (*Acid. Acet. Dil.*), *Diluted Acetic Acid, U. S. P.*— CH_3COOH (about 6%).

DOSAGE: 2 Cc. or 30 minims.

Acidum Aceticum Glaciale (*Acid. Acet. Glac.*), *Glacial Acetic Acid, U. S. P.*— CH_3COOH .

A colorless liquid, with a vinegar-like odor and a pungent, sour taste. Miscible with water or alcohol.

- ***Acidum Benzoicum** (Acid. Benz.), Benzoic Acid, U. S. P.— $C_6H_5.COOH$. Obtained from benzoin or prepared synthetically.

Lustrous, almost colorless scales or needles, with a pungent, acid taste. Slightly soluble in water (1:275) and freely soluble in alcohol (1:2.3); also readily soluble in solutions of alkali hydroxides or carbonates with the production of alkali benzoates.

ACTION AND USES: Mild antiseptic and diuretic.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), best given in the form of soluble benzoates (see under **Sodii Benzoas**).

- ***Acidum Boricum** (Acid. Bor.) Boric Acid, U. S. P. (Boracic Acid).— H_3BO_3 .

Colorless, odorless scales or powder with a faintly bitter taste. Soluble in water and in alcohol (1:18); freely soluble in glycerin (1:4).

ACTION AND USES: Mild antiseptic and astringent.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in solution. A watery solution, ranging from 2 per cent. to saturated, is used in the treatment of conjunctivitis, cystitis, etc. Externally used as dusting powder.

- ***Glyceritum Boroglycerini** (Glycer. Boroglyc.), Glycerite of Boroglycerin, U. S. P.—Boroglycerin in glycerin representing 31% boric acid.

Liquor Antisepticus (Liq. Antisept.), Antiseptic Solution, N. F.—Boric acid (2.5%), thymol, eucalyptol, methyl salicylate, oil of thyme, menthol, sodium salicylate and sodium benzoate in alcohol and water.

USES: Needlessly complex aromatic mouthwash.

- ***Unguentum Acidi Borici** (Ung. Acid. Bor.), Ointment of Boric Acid, U. S. P.—Boric acid (10%) in paraffin and white petrolatum.

- Acidum Bromauricum** (Acid. Bromaur.), Bromauric Acid, N. F.—Corresponds to not less than 32 per cent. of metallic gold.

A dark-brown, odorless solid, with a metallic acid taste. Very soluble in water and in alcohol.

ACTION AND USES: "Alterative" apparently of no value.

DOSAGE: 0.006 Gm. or $\frac{1}{10}$ grain (N. F.).

- Liquor Auri et Arseni Bromidi** (Liq. Aur. et Arsen. Brom.), Solution of Gold and Arsenic Bromide, N. F.—Contains hydrogen bromaurate ($HAuBr$) and arsenic acid (H_3AsO_4) obtained by oxidizing a solution of arsenous oxide with bromine and adding bromauric acid.

DOSAGE: 0.2 Cc. or 3 minims (N. F.).

***Acidum Citricum (Acid. Cit.), Citric Acid, U. S. P.**

Colorless, odorless crystals. Very soluble in water (1:0.5) and in alcohol (1:1.8).

ACTION AND USES: Substitute for lemon juice; also substitute for dilute hydrochloric acid in the treatment of gastric hypoacidity.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in solution.

Syrupus Acidi Citrici (*Syr. Acid. Cit.*), *Syrup of Citric Acid, U. S. P.*—Citric acid (1%), flavored with tincture of lemon peel, in syrup.

USES: Vehicle for salty substances such as bromides.

Acidum Formicum (Acid. Formic.), Formic Acid, N. F.—HCOOH (about 25%).

A clear, colorless liquid, with a characteristic pungent odor, and an acid reaction.

ACTION AND USES: Externally as caustic irritant, but without advantage over mustard. Internally as slightly irritant diuretic, but without advantage over nitrates. The elixirs containing various formates are intended as tonics, but there is no good evidence that they have such action. They are needlessly complex and irrational.

DOSAGE: 0.3 Cc. or 5 minims (N. F.).

Elixir Formatum (*Elix. Format.*), *Elixir of Formates, N. F.*—Potassium formate (5%), sodium formate (5%) and aromatic elixir. Absolute alcohol content about 15 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Formatum Compositum (*Elix. Format. Co.*), *Compound Elixir of Formates, N. F.*—Sodium formate (3%), magnesium formate (2.5%), strontium formate (2.5%), lithium formate (1%), quinine formate (0.8%), compound spirit of cardamom, acetic ether, alcohol, glycerin and distilled water. Absolute alcohol content about 9.9 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Spiritus Acidi Formici (*Sp. Ac. Formic.*), *Spirit of Formic Acid, N. F.* (*Spiritus Formicarum, Spirit of Ants.*)—Formic acid (4%) in distilled water and alcohol. Absolute alcohol content about 70 per cent.

USES: Rubefacient.

DOSAGE: 4 Cr or 1 fluidrachm (N. F.).

Acidum Gallicum (Acid. Gallic.), Gallic Acid, U. S. P.

An odorless, almost white powder, with an astringent taste. Soluble in water (1:87) and freely soluble in alcohol (1:4.6) and in glycerin (1:10).

ACTION AND USES: Mild astringent; not very effective.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Acidum Hydriodicum Dilutum (Acid. Hydriod. Dil.), Diluted Hydriodic Acid, U. S. P.—HI (about 10%).

A colorless, odorless liquid, with a strong acid taste. Incompatible with alkalis and their carbonates, also with metallic oxides and salts of silver and of lead.

ACTION AND USES: Has the general properties of iodides.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.), diluted. Administered chiefly in the form of syrup as a substitute for the alkali iodides, over which it has no material advantage.

Syrupus Acidi Hydriodici (Syr. Acid. Hydriod.) Syrup of Hydriodic Acid, U. S. P.—Hydriodic acid (about 1.25%).

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Acidum Hydrobromicum Dilutum (Acid. Hydrobrom. Dil.), Diluted Hydrobromic Acid, U. S. P.—HBr (about 10%).

A colorless, odorless liquid, with a strong acid taste. Incompatible with alkalis and their carbonates, metallic oxides and the salts of silver and of lead.

ACTION AND USES: In general, those of bromides; but the preparation is objectionable because of its acidity.

DOSAGE: 1 Cc. or 15 minims (U. S. P.), diluted.

***Acidum Hydrochloricum (Acid. Hydrochl.), Hydrochloric Acid, U. S. P. (Muriatic Acid).—HCl (about 32%).**

A fuming, corrosive liquid. Incompatible with alkalis and their carbonates, metallic oxides and salts of silver and of lead.

ACTION AND USES: Used in gastric hypoacidity; also as caustic; antiseptic.

**Acidum Hydrochloricum Dilutum (Acid. Hydrochl. Dil.), Diluted Hydrochloric Acid, U. S. P.—HCl (about 10%).*

A colorless, odorless, strongly acid solution.

DOSAGE: 1 Cc. or 15 minims (U. S. P.), after meals, diluted in one-half to one glass water and taken through a glass tube to protect the teeth.

**Acidum Hydrocyanicum Dilutum (A.
Diluted Hydrocyanic Acid, U.**

Acid, Acidum Hydrocyanicum Dilutum, P. I.).—HCN (about 2%).

A colorless liquid, with an odor like that of bitter almonds.

ACTION AND USES: Said to be sedative. Was formerly used as an addition to cough mixtures and as an antiemetic but, because of its poisonous properties and instability, is no longer used to the same extent.

DOSAGE: 0.1 Cc. or 1½ minims (U. S. P.)—not drops.

Acidum Hypophosphorosum (Acid. Hypophos.), Hypophosphorous Acid, U. S. P.— H_3PO_3 (about 31%).

A colorless, odorless liquid.

ACTION AND USES: Ingredient of compound hypophosphite preparations.

Acidum Hypophosphorosum Dilutum (Acid. Hypophos. Dil.), *Diluted Hypophosphorous Acid*, U. S. P.— H_3PO_3 (about 10%).

A colorless, odorless liquid with a strong acid taste.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.), diluted.

Acidum Lacticum (Acid. Lact.), Lactic Acid, U. S. P.—Equivalent to about 87 per cent. lactic acid.

A colorless or nearly colorless, nearly odorless, syrupy liquid, with an acid taste. Miscible with water.

ACTION AND USES: Caustic; seldom used internally.

DOSAGE: 2 Cc. or 30 minims (U. S. P.), diluted.

*Acidum Nitricum (Acid. Nitric.), Nitric Acid, U. S. P.— HNO_3 (about 68%).

A colorless, fuming, very corrosive liquid with a suffocating odor. Miscible with water.

ACTION AND USES: Caustic.

Acidum Nitrohydrochloricum (Acid. Nitrohydrochl.), Nitrohydrochloric Acid, U. S. P. (Aqua Regia, Nitromuriatic Acid).—A mixture of hydrochloric acid, nitric acid, nitrosyl chloride and chlorine.

A golden-yellow, fuming, very corrosive liquid. Miscible with water.

ACTION AND USES: Caustic. Formerly thought to be a "hepatic stimulant," but acts much like other mineral acids.

Acidum Nitrohydrochloricum Dilutum (Acid. Nitrohydrochl. Dil.), *Diluted Nitrohydrochloric Acid*, U. S. P. (Diluted Nitromuriatic Acid).—An aqueous solution of about one-

fifth the strength of nitrohydrochloric acid; should be freshly prepared.

DOSAGE: 1 Cc. or 15 minims (U. S. P.), well diluted.

Acidum Oleicum (Acid. Oleic.), Oleic Acid, U. S. P.—
Obtained from fats.

A yellowish or brownish-yellow, oily liquid, having a lard-like odor and taste. Practically insoluble in water, partially soluble in 60 per cent. alcohol.

ACTION AND USES: Solvent for making oleates.

***Acidum Phenylcinchoninicum (Acid. Phenylcinch.), Phenylcinchoninic Acid, U. S. P. (Phenyl-Quinoline-Carboxylic Acid, Atophan).**

A white or nearly white, odorless or nearly odorless powder with a bitter taste. Insoluble in cold water, slightly soluble in cold alcohol.

ACTION AND USES: Increases the excretion of uric acid and diminishes its concentration in the blood. Analgesic. Has been used in acute and chronic gout.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in tablets or powder.

Acidum Phosphoricum (Acid. Phos.), Phosphoric Acid, U. S. P.— H_3PO_4 (about 86.5%).

A colorless, odorless, syrupy liquid having a strongly acid taste. Miscible with water. Incompatible with alkalis, alkali carbonates, ferric chloride, lead acetate and solutions of lime.

ACTION AND USES: Similar to those of hydrochloric acid. Has none of the therapeutic properties of free phosphorus.

Acidum Phosphoricum Dilutum (Acid. Phos. Dil.), Diluted Phosphoric Acid, U. S. P.— H_3PO_4 (about 10%),

DOSAGE: 2 Cc. or 30 minims (U. S. P.), diluted.

Liquor Phosphatum Acidus (Liq. Phos. Acid.), Acid Solution of Phosphates, N. F. (Solution of Acid Phosphates).

—Made by dissolving precipitated calcium carbonate (5%) in phosphoric acid and water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.)

Liquor Phosphatum Compositus (Liq. Phos. Co.), Compound Solution of Phosphates, N. F.—Phosphates of calcium, iron, ammonium, potassium and sodium in citric acid, glycerin, orange flower water and water.

Syrupus Phosphatum Compositum (Syr. Phos. Co.), Compound Syrup of the Phosphates, N. F. (Chemical Food).—Compound solution of phosphates (50%), colored with tincture of cudbear, in glycerin and syrup.

USES: Antiquated and irrational "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Phosphatum cum Quinina et Strychnina.—See under Quininae Hydrochloridum.

Acidum Picricum, Picric Acid.—See Trinitrophenol.

**Acidum Salicylicum (Acid. Salicyl.), Salicylic Acid, U. S. P.*
— $C_6H_4(OH)COOH$.

A white, odorless powder, tasting first sweetish, subsequently acid. Slightly soluble in water (1:460), freely soluble in alcohol (1:2.7) and in ether. Incompatible with solutions of iron and with spirit of nitrous ether.

ACTION AND USES: Antirheumatic, antiseptic, germicide and keratolytic.

DOSAGE: Internally 0.75 Gm. or 12 grains (U. S. P.) best given in the form of soluble salicylates (see *Sodii Salicylas*). Externally in 10 per cent. ointment as a keratolytic.

Collodium Salicylici Compositum (Collod. Salicyl. Co.), Compound Salicylic Collodion, N. F.—Salicylic acid (1.1%) and flexible collodion colored with fluidextract of cannabis.

USES: A popular application for softening corns. The cannabis can play no rôle beyond that of coloring agent.

Glycerogelatinum Acidi Salicylici (Glycerogel. Acid. Salicyl.), Salicylic Acid Glycerogelatin, N. F.—Salicylic acid (10%), in water, glycerin and glycerinated gelatin.

Mulla Acidi Salicylici (Mull. Ac. Salicyl.), Salicylic Acid Mull, N. F. (*Unguentum Salicylatum Extensum, N. F. III*).—Salicylic acid (10%), in benzoinated suet and benzoinated lard.

Mulla Creosoti Salicylata (Mull. Creosot. Salicyl.), Salicylated Creosote Mull, N. F. (*Unguentum Creosoti Salicylatum Extensum, N. F. III*).—Salicylic acid (10%), and creosote (20%) in yellow wax and benzoinated suet.

Pulvis Talcı Compositus (Pulv. Talc. Co.), Compound Powder of Talc, N. F. (*Boro-Salicylated Powder of Talc.*).—Salicylic acid (3%), boric acid (10%) and talc.

USES: Toilet powder, used for local sweating.

Stili Acidi Salicylici Dilubiles (Stil. Acid. Salicyl. Dilub.), Salicylic Acid Pencil, N. F.—Salicylic acid (10%) with tragacanth, starch, white dextrin and sugar.

USES: For the application of salicylic acid to limited areas.

Acidum Stearicum (Acid. Stear.), Stearic Acid, U. S. P.—
Obtained from fats.

A white, hard, unctuous, odorless and tasteless substance. Soluble in alcohol (1:21) and in chloroform (1:2), but insoluble in water.

USES: Used in making glycerin suppositories.

Acidum Sulphuricum (Acid. Sulph.), Sulphuric Acid, U. S. P.
— H_2SO_4 (about 94%).

A colorless, odorless, corrosive liquid. Miscible with water or alcohol with evolution of much heat.

ACTION AND USES: Similar to those of hydrochloric acid. Externally, the dilute acid is used as astringent.

Acidum Sulphuricum Aromaticum (Acid. Sulph. Arom.), Aromatic Sulphuric Acid, U. S. P.—A mixture, containing free sulphuric acid and ethylsulphuric acid equivalent to about 20 per cent. of H_2SO_4 in alcohol, flavored with ginger and cinnamon. Absolute alcohol content about 85 per cent.

A clear, reddish-brown liquid with an aromatic odor.

DOSAGE: 1 Cc. or 15 minims, well diluted.

Acidum Sulphuricum Dilutum (Acid. Sulph. Dil.), Diluted Sulphuric Acid, U. S. P.— H_2SO_4 (about 10%).

DOSAGE: 1 Cc. or 15 minims (U. S. P.), well diluted.

***Acidum Tannicum (Acid. Tann.), Tannic Acid, U. S. P.**
(Gallotannic Acid, Tannin).—Usually obtained from nutgalls.

A yellowish-white to light brown powder, odorless or having a faint, characteristic odor and a strongly astringent taste. Freely soluble in glycerin (1:1) and very soluble in water and in alcohol. Incompatible with soluble preparations of iron.

ACTION AND USES: Astringent; local hemostatic; antidote for certain alkaloidal and metallic poisons.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Collodium Stypticum (Collod. Stypt.), Styptic Collodion, N. F.—Tannic acid (20%) and flexible collodion. There is no advantage in applying tannin as collodion.

***Glyceritum Acidi Tannici (Glycer. Acid. Tann.), Glycerite of Tannic Acid, U. S. P.** (Glycerite of Tannin).—Tannic acid (20%) in glycerin.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Trochisci Acidi Tannici (Troch. Acid. Tann.), Troches of Tannic Acid, U. S. P.—Each troche contains tannic acid (0.06 Gm. or 1 grain), with sugar and tragacanth, flavored with orange flower.

Unguentum Acidi Tannici (*Ung. Acid. Tann.*), *Ointment of Tannic Acid*, U. S. P.—Tannic acid (20%), in glycerin and ointment.

Acidum Tartaricum (*Acid. Tart.*), *Tartaric Acid*, U. S. P.—
Obtained from wine lees or argol.

A white, odorless powder, with an acid taste. Very soluble in water (1:0.75) and in alcohol (1:3.3).

ACTION AND USES: Mild acid; used in the manufacture of effervescent salts.

DOSAGE: 0.5 Gm. or 8 grains, diluted.

Acidum Trichloraceticum (*Acid. Trichloracet.*), *Trichloroacetic Acid*, U. S. P.— CCl_3COOH .

Colorless, deliquescent crystals with a slight characteristic odor. Miscible with water and with alcohol.

ACTION AND USES: Caustic, for the removal of warts and other skin blemishes. Used also as local hemostatic.

Aconitina (*Aconitin.*), *Aconitine*, U. S. P.—An alkaloid obtained from aconite.

A white, odorless powder. Very slightly soluble in water, soluble in alcohol (1:28) and in ether (1:65).

ACTION AND USES: Similar to those described under aconite. As several alkaloidal preparations of aconite, of varying activity, are sold under the name of aconitine, its use is not advisable.

DOSAGE: 0.15 mg. or $\frac{1}{400}$ grain (U. S. P.).

Oleatum Aconitinae (*Oleat. Aconitin.*), *Oleate of Aconitine*, N. F.—Aconitine (2%) in oleic acid and olive oil.

USES: Without advantage over aconite preparations for external application.

***Aconitum** (*Aconit.*), *Aconite*, U. S. P. (*Monkshood*, *Aconite Root*, *Aconiti Tuber*, P. I.).—A tuberous root, yielding about 0.5 per cent. of ether-soluble alkaloids; also assayed biologically. Physicians should specify preparations that have been assayed biologically, since the alkaloidal assay is not a reliable index of activity.

ACTION AND USES: Slows the pulse and lowers the blood pressure, but this action in therapeutic doses is uncertain. Used in the treatment of febrile colds; also for counter-irritation, in the form of the tincture. Used also as a local anesthetic, and by some still believed to be useful in the treatment of the symptoms arising from excessive cardiac

hypertrophy. Toxic symptoms may occur from local absorption.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.); not used as such.

Extractum Aconiti (Ext. Aconit.), Extract of Aconite, U. S. P. (Powdered Extract of Aconite).—Yields about 2 per cent. of ether-soluble alkaloids. Biologically assayed by producing death in the guinea-pig.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.). Physicians who prescribe extract of aconite should specify the biologically assayed product.

Fluidextractum Aconiti (Fldext. Aconit.), Fluidextract of Aconite, U. S. P.—Aconite (100%), yielding about 0.5 per cent. of alkaloids. Absolute alcohol content about 70 per cent. Biologically assayed by producing death in the guinea-pig.

DOSAGE: 0.03 Cc. or $\frac{1}{2}$ minim (U. S. P.). Physicians who prescribe fluidextract of aconite should specify the biologically assayed product. Best to use the tincture.

Linimentum Aconiti et Chloroformi (Lin. Aconit. et Chlorof.), Liniment of Aconite and Chloroform, N. F.—Fluidextract of aconite (4.5%), in alcohol, chloroform and soap liniment.

**Tinctura Aconiti (Tr. Aconit.), Tincture of Aconite, U. S. P.* (Aconiti Tinctura, P. I.).—Aconite (10%) yielding about 0.05 per cent. of alkaloids, in alcohol and water. Alkaloidal assay unreliable. Biologically assayed by producing death in the guinea-pig. Absolute alcohol content about 65 per cent.

DOSAGE: 0.3 Cc. or 5 minims (U. S. P.). Physicians should specify the biologically assayed product.

***Adeps (Adeps), Lard, U. S. P.**

Insoluble in water, and only slightly soluble in alcohol.

ACTION AND USES: Basis for ointments. Used particularly when absorption is desired.

**Adeps Benzoïnatus (Adeps Bens.), Benzoïnated Lard, U. S. P.*—Somewhat antiseptic and less liable to rancidity than ordinary lard.

Unguentum (Ung.), Ointment, U. S. P. (Simple Ointment).—White wax (20%) and benzoïnated lard.

Adeps Lanae (Adeps Lan.), Wool Fat, U. S. P. (Anhydrous Lanolin).—Purified wool fat, freed from water.

Insoluble in but miscible with about twice its weight of water; sparingly soluble in cold alcohol.

ACTION AND USES: Basis for ointments; because of its tenacious consistency, should be mixed with some other base.

**Adeps Lanae Hydrosus (Adeps Lan. Hyd.), Hydrous Wool Fat, U. S. P. (Lanolin).*—Wool fat with about 27 per cent. of water.

USES: Ointment used particularly when absorption through the skin is desired; mixed with white petrolatum or lard.

Adonis (Adonis), Adonis, N. F. (Pheasant's Eye).—The plant deprived of its root.

ACTION AND USES: Cardiac tonic of digitalis group, but of unreliable activity.

DOSAGE: 0.125 Gm. or 2 grains (N. F.).

Fluidextractum Adonidis (Fldext. Adonid.), Fluidextract of Adonis, N. F.—Adonis (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 0.125 Cc. or 2 minims (N. F.).

**Aether (Aether), Ether, U. S. P.*—Contains about 96.5 per cent. of ethyl oxide.

A clear, colorless, very inflammable liquid with a characteristic odor and a burning, sweetish taste. Soluble in water (1:12), and miscible with alcohol, chloroform, petroleum benzin, benzene, and fixed or volatile oils.

ACTION AND USES: General anesthetic, administered by inhalation; when administered by mouth, carminative.

DOSAGE: 1 Cc. or 15 minims.

**Spiritus Aetheris (Sp. Aeth.), Spirit of Ether, U. S. P. (Hoffmann's Drops).*—Ether (32.5%) in alcohol. Absolute alcohol content about 65 per cent.

USES: Carminative and fugacious indirect circulatory stimulant.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Spiritus Aetheris Compositus (Sp. Aether. Co.), Compound Spirit of Ether, N. F. (Hoffmann's Anodyne).—Ether (32.5%), ethereal oil (2.5%) and alcohol. Absolute alcohol content about 60 per cent.

USES: A superfluous modification of spirit of ether.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Aether Aceticus (Aeth. Acet.), Acetic Ether, N. F. (Ethyl Acetate).

A colorless, fragrant liquid, with a burning taste.

ACTION AND USES: Carminative, similar to and without special advantage over ether. Also used pharmaceutically, as flavoring ingredient.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Aether Nitrosus, Nitrous Ether.—Ethyl nitrite.**

***Spiritus Aetheris Nitrosi (Sp. Aeth. Nitros.), Spirit of Nitrous Ether, U. S. P. (Sweet Spirit of Nitre).—**Ethyl nitrite (about 4%) in alcohol. Absolute alcohol content about 92 per cent.

ACTION AND USES: Popularly used as a weak diuretic and diaphoretic. Unstable and therefore unreliable. Has the physiologic action of a weak nitrite.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Aethylis Carbamas (Aethyl. Carbam.), Ethyl Carbamate, U. S. P. (Carbamic Acid Ethyl Ester, Urethane, Ethyl Urethane).

Colorless, odorless crystals or scales, with a cooling, saline taste. Freely soluble in water (1:0.45) and in alcohol (1:0.8). Incompatible with alkalis, acids and most other chemicals.

ACTION AND USES: Very mild and relatively harmless hypnotic.

DOSAGE: 1 Gm. or 15 grains (U. S. P.) in powder, cachets or solution.

***Aethylis Chloridum (Aethyl. Chlor.), Ethyl Chloride, U. S. P.—Monochlorethane, $\text{CH}_3\text{CH}_2\text{Cl}$.**

A colorless, very volatile liquid, with an agreeable odor and a sweetish, burning taste. The vapor is very inflammable. Slightly soluble in water and freely soluble in alcohol and in ether.

ACTION AND USES: Local anesthetic for minor operations. Used in the form of a spray to produce intense cold. Has also been used by inhalation as general anesthetic in short operations.

***Aethylmorphinae Hydrochloridum (Aethylmorph. Hydrochl.), Ethylmorphine Hydrochloride, U. S. P. (Ethylmorphine Chloride, Dionin).**

A white, odorless powder, with a slightly bitter taste. Very soluble in water (1:8) and soluble in alcohol (1:22).

ACTION AND USES: Systemically, intermediate between those of morphine and codeine. Probably possesses no

advantage over codeine. When applied to the eye it causes local hyperemia terminating in acute conjunctival edema.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.). Used as 10 per cent. ointment or solution in corneal opacity and similar conditions.

Agar (Agar), Agar, U. S. P. (Agar-Agar).—A mucilaginous substance extracted from certain seaweeds.

Nearly white, almost odorless and tasteless strips, shreds or powder, tough when damp, brittle when dry. Insoluble in cold water, but slowly soluble in hot water.

ACTION AND USES: Passes through the intestinal canal almost unchanged. Absorbs and retains moisture. Used in chronic constipation with intestinal atony; renders the feces soft and bulky and thus promotes peristalsis.

DOSAGE: 10 Gm. or $2\frac{1}{2}$ drachms (U. S. P.).

Agaricus (Agaric), Agaric, N. F. (White Agaric, Larch Agaric).—The fruit body of a fungus.

ACTION AND USES: Empirical and unreliable remedy for night sweats of phthisis.

DOSAGE: 0.6 Gm. or 10 grains (N. F.).

***Alcohol (Alcohol), Alcohol, U. S. P.**—Not less than 92.3 per cent. by weight or 94.9 per cent. by volume of C_2H_5OH .

A colorless, volatile, inflammable liquid, with a characteristic odor and burning taste. Freely miscible with water, ether or chloroform.

ACTION AND USES: Rubefacient, astringent, antiseptic and by its evaporation a refrigerant. Internally it is a narcotic widely used as a "stimulant" but may do more harm than good.

Alcohol Dehydratum (Alcohol Dehyd.), Dehydrated Alcohol, U. S. P. (Alcohol Absolutum, U. S. P. VIII).—Not less than 99 per cent. by weight of C_2H_5OH .

A liquid with the same color, odor and taste as alcohol.

Alcohol Dilutum (Alcohol Dil.), Diluted Alcohol, U. S. P.—About 41.5 per cent. by weight or 49 per cent. by volume of C_2H_5OH .

***Elixir Aromaticum (Elix. Arom.), Aromatic Elixir, U. S. P. (Simple Elixir).**—Compound spirit of orange in syrup, alcohol and distilled water. Absolute alcohol content about 22 per cent.

USES: Diluent and vehicle. The alcohol content of this preparation should not be overlooked.

Elixir Aromaticum Rubrum (Elix. Arom. Rub.), Red Aromatic Elixir, N. F. (Red Elixir).—Aromatic elixir colored with cudbear.

USES: Diluent and vehicle.

Elixir Glycyrrhizae (Elix. Glycyrrh.), Elixir of Glycyrrhiza, U. S. P. (Elixir Adjuvans, U. S. P. VIII, Elixir of Licorice).—Fluidextract of glycyrrhiza (12.5%) and aromatic elixir. Absolute alcohol content about 22 per cent.

USES: Diluent and vehicle. The alcohol content should not be overlooked.

Elixir Glycyrrhizae Aromaticum (Elix. Glycyrrh. Arom.), Aromatic Elixir of Glycyrrhiza, N. F. (Aromatic Elixir of Licorice).—Fluidextract of glycyrrhiza (12.5%) oil of clove, oil of cassia, oil of myristica, oil of fennel and aromatic elixir. Absolute alcohol content about 22 per cent.

USES: Flavoring vehicle.

DOSAGE: 8 Cc. or 2 fluidrachms.

Aletris (Aletr.), Aletris, N. F. (Unicorn Root, Colic Root, Star Grass).—Rhizome and root.

ACTION AND USES: Simple bitter, without advantage over gentian. Also used in proprietary "female remedies," without good evidence of value.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Aletridis (Fldext. Aletrid.), Fluidextract of Aletris, N. F.—Aletris (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Allium (Allium), Garlic, N. F.

ACTION AND USES: Irritant expectorant; without advantage over senega, terpin hydrate or asafetida.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Syrupus Allii (Syr. Allii), Syrup of Garlic, N. F.—Garlic (20%) in sugar and diluted acetic acid.

USES: Irritant expectorant.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Aloe (Aloe), Aloes, U. S. P.*—The inspissated juice of different species of aloes, yielding Socotrine, Curaçao and Cape aloes, respectively

ACTION AND USES: Moderately active cathartic used in the treatment of chronic constipation. Acts mostly on the large intestine.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), as pills.

**Extractum Aloes (Ext. Aloes), Extract of Aloes, N. F.* (Powdered Extract of Aloes).—An aqueous extract, 1 Gm. representing 2 Gm. of aloes; diluted with starch. No material advantage over aloes.

DOSAGE: 0.125 Gm. or 2 grains (N. F.).

Pilulae ad Prandium (Pil. ad Prand.), Dinner Pills, N. F.

1. For dinner pills without additional specifications *Pilulae Aloes et Mastiches* (which see), also called Lady Webster's Pills, should be dispensed.

2. CHAPMAN'S DINNER PILLS, N. F.—Each pill contains aloes and mastic (each 0.097 Gm. or $1\frac{1}{2}$ grains), and ipecac (0.065 Gm. or 1 grain), with oil of fennel.

3. COLE'S DINNER PILLS, N. F.—Each pill contains aloes, mass of mercury and jalap (each 0.078 Gm. or $1\frac{1}{4}$ grains), antimony and potassium tartrate (0.0013 Gm. or $\frac{1}{60}$ grain) and syrup.

4. HALL'S DINNER PILLS, N. F.—Each pill contains aloes (0.065 Gm. or 1 grain) with extract of glycyrrhiza and soap.

DOSAGE: 1 pill (N. F.).

Pilulae Aloes (Pil. Aloes), Pills of Aloes, U. S. P.—Each pill contains aloes (0.13 Gm. or 2 grains) with soap.

DOSAGE: 2 pills (U. S. P.).

Pilulae Aloes et Asafoetidae (Pil. Aloe. et Asafoet.), Pills of Aloes and Asafetida, N. F.—Each pill contains aloes and asafetida (each 0.09 Gm. or $1\frac{1}{2}$ grains) with soap.

DOSAGE: 1 pill (U. S. P.).

Pilulae Aloes et Ferri (Pil. Aloe. et Ferr.), Pills of Aloes and Iron, N. F.—Each pill contains aloes and exsiccated ferrous sulphate (each 0.07 Gm. or $1\frac{1}{6}$ grains), with aromatic powder and confection of rose.

DOSAGE: 2 pills (N. F.).

Pilulae Aloes et Mastiches (Pil. Aloe. et Mastich.), Pills of Aloes and Mastic, N. F. (Lady Webster's Dinner Pill).—Each pill contains aloes (0.13 Gm. or 2 grains) and mastic (0.04 Gm. or $\frac{2}{3}$ grain) with red rose.

DOSAGE: 2 pills (N. F.).

Pilulae Aloes et Myrrhae (Pil. Aloe. et Myrrh.), Pills of Aloes and Myrrh, N. F.—Each pill contains aloes (0.13 Gm. or 2 grains) and myrrh (0.06 Gm. or 1 grain), with aromatic powder and syrup.

DOSAGE: 2 pills (N. F.).

Pilulae Aloes et Podophylli Compositae (Pil. Aloes. et Podoph. Co.), Compound Pills of Aloes and Podophyllum, N. F. (Janeway's Pills).—Each pill contains aloes (0.065 Gm. or 1 grain), resin of podophyllum (0.0325 Gm. or $\frac{1}{2}$ grain), pilular extract of belladonna leaves and extract of nux vomica (each 0.016 Gm. or $\frac{1}{4}$ grain).

DOSAGE: 1 pill (N. F.).

Pilulae Aloes, Hydrargyri et Podophylli (Pil. Aloes. Hydrarg. et Podoph.), Pills of Aloes, Mercury and Podophyllum, N. F. (Pilulae Triplices, N. F. III, Triplex Pills, Pilula Triplex).—Each pill contains aloes (0.13 Gm. or 2 grains), mass of mercury (0.065 Gm. or 1 grain) and resin of podophyllum (0.016 Gm. or $\frac{1}{4}$ grain). These pills should not be taken habitually.

DOSAGE: 1 pill (N. F.).

Pilulae Aloes, Hydrargyri et Scammonii Compositae (Pil. Aloes. Hydrarg. et Scammon Co.), Compound Pills of Aloes, Mercury and Scammony, N. F. (Francis' Triplex Pills).—Each pill contains aloes, resin of scammony and mass of mercury (each 0.055 Gm. or $\frac{1}{2}$ grain), croton oil (0.0032 Cc. or $\frac{1}{20}$ minim), oil of caraway (0.016 Cc. or $\frac{1}{4}$ minim) and tincture of aloes and myrrh. Should not be taken habitually.

DOSAGE: 1 pill (N. F.).

Pulvis Aloes et Canellae (Pulv. Aloes. et Canell.), Powder of Aloes and Canella, N. F. (Hiera Picra).—Aloes (80%) and canella.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Tinctura Aloes (Tr. Aloes), Tincture of Aloes, U. S. P.—Aloes (10%) and glycyrrhiza in diluted alcohol. Absolute alcohol content about 47 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Aloes et Myrrhae (Tr. Aloes. et Myrrh.), Tincture of Aloes and Myrrh, N. F.—Represents aloes and myrrh (each 10%) and glycyrrhiza in alcohol and water. Absolute alcohol content about 70 per cent.

USES: Disagreeable cathartic.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Aloinum** (Aloin.), Aloin, U. S. P.—Obtained from aloes.

A yellow, odorless, intensely bitter powder. Slightly soluble in water and soluble in alcohol.

ACTION AND USES: Similar to those of aloes; is more easily absorbed, and has sometimes produced renal irritation.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.), as pills. In chronic constipation, frequently given in doses of from 0.006 Gm. to 0.02 Gm., or $\frac{1}{12}$ to $\frac{1}{8}$ grain in combination with extract of belladonna and strychnine.

Pilulae Aloini Compositae (Pil. Aloin. Co.), *Compound Pills of Aloin*, N. F.—Each pill contains aloin (0.0325 Gm. or $\frac{1}{2}$ grain), resin of podophyllum (0.008 Gm. or $\frac{1}{8}$ grain), and pilular extract of belladonna leaves (0.016 Gm. or $\frac{1}{4}$ grain).

DOSAGE: 1 pill (N. F.).

Pilulae Aloini, Strychninae et Belladonnae (Pil. A. S. et B.), *Pills of Aloin, Strychnine and Belladonna*, N. F.—Each pill contains aloin (0.013 Gm. or $\frac{1}{6}$ grain), strychnine (0.5 mg. or $\frac{1}{120}$ grain) and pilular extract of belladonna leaves (0.008 Gm. or $\frac{1}{8}$ grain).

DOSAGE: 1 pill (N. F.).

Pilulae Aloini, Strychninae et Belladonnae Compositae (Pil. A. S. et B. Co.), *Compound Pills of Aloin, Strychnine and Belladonna*, N. F.—Each pill contains aloin (0.013 Gm. or $\frac{1}{6}$ grain), strychnine (0.5 mg. or $\frac{1}{120}$ grain), pilular extract of belladonna leaves (0.008 Gm. or $\frac{1}{8}$ grain) and extract of cascara sagrada (0.0325 Gm. or $\frac{1}{2}$ grain).

USES: Needless complex mixture.

DOSAGE: 1 pill (N. F.).

Pilulae Laxativae Compositae (Pil. Lax. Co.), *Compound Laxative Pills*, N. F.—Each pill contains aloin (0.013 Gm. or $\frac{1}{6}$ grain), strychnine 0.5 mg. or $\frac{1}{120}$ grain), pilular extract of belladonna leaves (0.008 Gm. or $\frac{1}{8}$ grain) and ipecac (0.004 Gm. or $\frac{1}{15}$ grain) with glycyrrhiza and syrup.

USES: Complex and superfluous preparation of aloin.

DOSAGE: 2 pills (N. F.).

Althaea (Althaea), Althaea, U. S. P. (Marsh Mallow Root).

ACTION AND USES: Demulcent and emollient; without advantages over other drugs of the same kind, such as chondrus, tragacanth, slippery elm and flaxseed.

Species Pectorales (Spec. Pect.), *Pectoral Species*, N. F. (Species ad Infusum Pectorale, Breast Tea).—Althaea, coltsfoot, glycyrrhiza, anise, mullein flowers and orris root.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Syrupus Althaeae (Syr. Althae.), *Syrup of Althaea*, N. F.—Althaea (5%) with alcohol, glycerin and sugar in water. Absolute alcohol content about 2.5 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Althaeae Folia (Althae. Fol.), **Althaea Leaves**, N. F. (Marsh Mallow Leaves).

ACTION AND USES: Domestic demulcent and emollient; inferior for internal use to gums and for poultices to flaxseed.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Species Emollientes (*Spec. Emoll.*), *Emollient Species*, N. F. (Emollient Cataplasm).—Althaea leaves, mallow leaves, melilot, matricaria and flaxseed. Emollient poultice is made by adding a suitable quantity of hot water to this powder.

***Alumen** (Alum.), **Alum**, U. S. P.— $\text{AlNH}_4(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$ or $\text{AlK}(\text{SO}_4)_2 \cdot 12\text{H}_2\text{O}$. (The physician may indicate whether ammonium alum or potassium alum is desired.)

Colorless crystals or white powder, odorless and possessing a strongly astringent taste. Ammonium alum is somewhat less soluble in water than is potassium alum, which is freely soluble in water (1:7.2); both are insoluble in alcohol.

ACTION AND USES: Astringent, styptic and hemostatic. Seldom administered internally.

DOSAGE: As a gargle, in from 1 to 5 per cent. solution (somewhat injurious to the teeth); as an injection in gonorrhea, in from 0.5 to 1 per cent. solution; as a lotion in skin diseases, in 1 per cent. solution.

***Alumen Exsiccatum** (*Alum. Exsic.*), *Exsiccated Alum*, U. S. P. (Alumen Ustum, Dried Alum, Burnt Alum).—Anhydrous $\text{AlNH}_4(\text{SO}_4)_2$ or anhydrous $\text{AlK}(\text{SO}_4)_2$. (The composition of the desired salt may be indicated.)

White, odorless powder, with an astringent taste. Very slowly soluble in water (1:14); insoluble in alcohol.

ACTION AND USES: Used externally like alum, being more escharotic.

***Alumini Acetas**, **Aluminum Acetate**.—Normal aluminum acetate, $\text{Al}(\text{C}_2\text{H}_3\text{O}_2)_3$.

Liquor Alumini Acetatis (*Liq. Alumin. Acet.*), *Solution of Aluminum Acetate*, N. F. (Liquor Burowii, Burow's Solution).—Contains normal aluminum acetate, $\text{Al}(\text{C}_2\text{H}_3\text{O}_2)_3$ (about 5%).

USES: Popular astringent wash.

DOSAGE: Usually diluted with 15 parts of water.

Alumini Acetico-Tartras, Aluminum Acetico-Tartrate.

Liquor Alumini Acetico-Tartratis (Liq. Alumin. Acet. Tart.), Solution of Aluminum Acetico-Tartrate, N. F.—

A solution of aluminum acetate and tartrate obtained by dissolving aluminum hydroxide in a mixture of acetic and tartaric acids.

Alumini Chloridum (Alum. Chlor.), Aluminum Chloride, N. F.
— $\text{AlCl}_3 + 6\text{H}_2\text{O}$.

A nearly white deliquescent powder, with a sweetish, astringent taste. Freely soluble in water (1:1) and in alcohol (1:3).

ACTION AND USES: Antiseptic and astringent, without advantage over alum.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Alumini Hydroxidum (Alum. Hydrox.), Aluminum Hydroxide, U. S. P.— $\text{Al}(\text{OH})_3$.

A white, bulky, odorless and tasteless powder. Insoluble in water and in alcohol.

ACTION AND USES: Mild astringent and desiccant, sometimes used as a dusting powder.

Alumini Subacetat, Aluminum Subacetate.—Basic aluminum acetate, $\text{Al}(\text{C}_2\text{H}_3\text{O}_2)_2\text{OH}$.

**Liquor Alumini Subacetatis (Liq. Alumin. Subacet.), Solution of Aluminum Subacetate, N. F.* (Liquor Alumini Acetatis, N. F. III).—Contains basic aluminum acetate, $\text{Al}(\text{C}_2\text{H}_3\text{O}_2)_2\text{OH}$ (about 8%).

USES: Astringent wash.

DOSAGE: Usually diluted with ten parts of water.

Alumini Sulphas (Alum. Sulph.), Aluminum Sulphate, N. F.
— $\text{Al}_2(\text{SO}_4)_3 + 16\text{H}_2\text{O}$.

A white, odorless powder, with an astringent taste. Freely soluble in water (1:1), but insoluble in alcohol.

ACTION AND USES: Similar to and without advantage over alum.

***Ammonia (NH_3).**

**Aqua Ammoniae (Aq. Ammon.), Ammonia Water, U. S. P.*— NH_3 (about 10%) in water.

A liquid with a strong odor and a caustic, soapy taste. Loses NH_3 readily. Miscible with water or alcohol.

USES: Local irritant and antacid, preferably as a liniment. Fugacious indirect circulatory and respiratory stimulant.

Aqua Ammoniae Fortior (Aq. Ammon. Fort.), Stronger Ammonia Water, U. S. P.— NH_3 (about 28%) in water.

Loses NH_3 very readily. Miscible with water or alcohol.

**Linimentum Ammoniae (Lin. Ammon.), Ammonia Liniment, U. S. P.* (Volatile Liniment, Hartshorn Liniment).—Ammonia water (25%) in sesame oil.

Uses: Popularly used as counterirritant.

Lotio Ammoniacalis Camphorata (Lot. Ammon. Camph.), Ammoniated Camphor Wash, N. F. (Aqua Sedativa, N. F. III, Sedative Water, Eau Sédative de Raspail).—Ammonia water (6%) and spirit of camphor (1%), in sodium chloride and water.

Uses: Counterirritant.

Spiritus Ammoniae Anisatus (Sp. Ammon. Anis.), Anisated Spirit of Ammonia, N. F. (Liquor Ammoniae Anisatus, Anisated Solution of Ammonia).—Ammonia water (20%) with anethol, in alcohol. Absolute alcohol content about 70 per cent.

Uses: Aromatic carminative preparation.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Ammonii Acetas, Ammonium Acetate.— $\text{NH}_4\text{C}_2\text{H}_3\text{O}_2$.

Liquor Ammonii Acetatis (Liq. Ammon. Acet.), Solution of Ammonium Acetate, U. S. P. (Spirit of Mindererus).—Contains $\text{NH}_4\text{C}_2\text{H}_3\text{O}_2$ (not less than 7%).

Uses: Diaphoretic and diuretic in fevers; of doubtful value.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Liquor Ferri et Ammonii Acetatis (Liq. Ferr. et Ammon. Acet.), Solution of Iron and Ammonium Acetate, U. S. P. (Basham's Mixture).

Uses: An antiquated preparation, without the special advantages generally ascribed to it. Iron and acetates are better prescribed separately.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Ammonii Benzoas (Ammon. Benz.), Ammonium Benzoate, U. S. P.

A white, odorless or nearly odorless powder, with a bitterish taste. Very soluble in water (1:10) and soluble in alcohol (1:35). Incompatible with acids, fixed alkalies and ferric salts.

ACTION AND USES: Resembles sodium benzoate in its action but is more irritant and has no advantage over it.

DOSAGE: 1 Gm. or 15 grains (U. S. P.), in solution.

Ammonii Bromidum (Ammon. Brom.), Ammonium Bromide, U. S. P.— NH_4Br .

A white, odorless powder with a pungent saline taste. Incompatible with nitrous ether.

ACTION AND USES: Those of the bromides; more irritant than potassium bromide, over which it has no advantage.

DOSAGE: 1 Gm. or 15 grains (U. S. P.) in solution.

Elixir Ammonii Bromidi (Elix. Ammon. Brom.), Elixir of Ammonium Bromide, N. F.—Ammonium bromide (8.5%), syrup, distilled water and aromatic elixir. Absolute alcohol content about 5 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm.

Elixir Trium Bromidorum (Elix. Tri. Brom.), Elixir of Three Bromides, N. F.—Ammonium bromide, potassium bromide and sodium bromide (each 8%), with cudbear and compound elixir of almond. Absolute alcohol content about 4 per cent.

USES: An irrational bromide mixture, as there is no good evidence that the bases influence the bromide action materially.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Ammonii Carbonas (Ammon. Carb.), Ammonium Carbonate, U. S. P.**—A mixture of ammonium bicarbonate and ammonium carbamate, yielding about 31 per cent. of NH_3 .

White, hard, translucent masses, with a strong ammoniacal odor and a sharp, ammoniacal taste. Very soluble in water; alcohol dissolves the carbamate and leaves the bicarbonate.

ACTION AND USES: Alkaline and nauseant liquefying expectorant used in cough mixtures; also as reflex "diffusible" stimulant in smelling salts.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.) in solution.

Mistura Pectoralis Stokes (Mist. Pect. Stokes), Stokes' Expectorant, N. F.—Ammonium carbonate (1.75%), fluid-extracts of senega and squill (each 3.5%), camphorated tincture of opium (17.5%), water and syrup of tolu. Absolute alcohol content about 10 per cent.

USES: Irrational expectorant mixture.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Spiritus Ammoniae Aromaticus (Sp. Ammon. Arom.), Aromatic Spirit of Ammonia, U. S. P.**—Ammonium carbonate (3.4%), ammonia water (9%) and oils of lemon, lavender and myristica in alcohol and distilled water. Absolute alcohol content about 68 per cent.

USES: Useful aromatic alkali and carminative in flatulence, colic, etc.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

***Ammonii Chloridum (Ammon. Chlor.), Ammonium Chloride, U. S. P.— NH_4Cl .**

A white, odorless powder, with a saline taste. Freely soluble in water (1:2.6) and soluble in alcohol (1:100).

ACTION AND USES: Liquefying expectorant; used in cough mixtures, and in the treatment of inflammation of the air-passages.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.), in solution or lozenges.

Mistura Ammonii Chloridi (Mist. Ammon. Chlor.), Mixture of Ammonium Chloride, N. F. (Mistura Solvens Simplex).—Ammonium chloride (2.5%) and pure extract of glycyrrhiza in water.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Trochisci Ammonii Chloridi (Troch. Ammon. Chlor.), Troches of Ammonium Chloride, U. S. P.—Each troche contains ammonium chloride (0.1 Gm.) with extract of glycyrrhiza, tragacanth and sugar, flavored with syrup of tolu.

Ammonii Citras, Ammonium Citrate.

Liquor Ammonii Citratis (Liq. Ammon. Cit.), Solution of Ammonium Citrate, N. F.—Ammonium citrate (14.47%) in water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ammonii Hypophosphis (Ammon. Hypophos.), Ammonium Hypophosphite, N. F.— $\text{NH}_4\text{H}_2\text{PO}_3$.

Colorless plates or white powder, odorless, with a saline, bitter taste. Very soluble in water (1:1) and soluble in alcohol (1:20).

ACTION AND USES: Similar to but without material advantage over ammonium chloride.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Syrupus Ammonii Hypophosphitis (Syr. Ammon. Hypophos.), Syrup of Ammonium Hypophosphite, N. F.—Ammonium hypophosphite (3.5%) and diluted hypophosphorous acid, flavored with compound spirit of vanilla, in water, glycerin and syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ammonii Iodidum (Ammon. Iod.), Ammonium Iodide, U. S. P.—NH₄I.

White, odorless powder, with a saline taste. Freely soluble in water (1:0.6), in alcohol (1:3.7) and in glycerin (1:1.5).

ACTION AND USES: Those of the iodides. More irritant and less stable than sodium or potassium iodide, over which it has no advantage.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.), in solution.

Linimentum Ammonii Iodidi (Lin. Ammon. Iod.), Liniment of Ammonium Iodide, N. F.—Ammonium iodide (about 0.5%) with oil of rosemary, oil of lavender, camphor, ammonia water and alcohol.

USES: Irrational preparation.

Ammonii Phosphas (Ammon. Phos.), Ammonium Phosphate, N. F.—A mixture of $(\text{NH}_4)_2\text{HPO}_4$ and $\text{NH}_4\text{H}_2\text{PO}_4$.

Colorless crystals or white powder. Freely soluble in water (1:4), but insoluble in alcohol.

ACTION AND USES: Similar to but without advantage over sodium phosphate.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Ammonii Salicylas (Ammon. Salicyl.), Ammonium Salicylate, U. S. P.

White, odorless crystals or powder with a bitterish taste. Very soluble in water (1:1) and in alcohol (1:3). Incompatible with acids and fixed alkalies.

ACTION AND USES: Those of salicylates; without advantage over sodium salicylate.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in solution.

Ammonii Valeras (Ammon. Valer.), Ammonium Valerate, U. S. P. (Ammonium Valerianate).—Of variable composition.

White plates, with a valerian odor and a sharp, sweetish taste. Very soluble in water (1:0.3) and in alcohol (1:0.6).

ACTION AND USES: Sedative in hysteria, depending on the valeric odor. Not very effective.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Elixir Ammonii Valeratis (Elix. Ammon. Valer.), Elixir of Ammonium Valerate, N. F. (Elixir Ammonii Valerianatis, N. F. III).—Ammonium valerate (3.5%) in red elixir. Absolute alcohol content about 20 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm. The alcohol content should be borne in mind.

Amygdala Dulcis (Amygd. Dulc.), Sweet Almond, U. S. P.

ACTION AND USES: Demulcent, especially as cosmetic.

Emulsum Amygdalae (*Emuls. Amygd.*), *Emulsion of Almond*, U. S. P. (Milk of Almond).—Sweet almond, acacia, sugar and water.

***Amylis Nitris** (*Amyl. Nitris*), *Amyl Nitrite*, U. S. P.—Not less than 80 per cent. $C_6H_{11}NO_2$ (chiefly iso-amyl nitrite).

A yellowish liquid, with a fruity odor and a pungent, aromatic taste. Very volatile; almost insoluble in water, but miscible with alcohol or ether.

ACTION AND USES: Prompt vasodilator, used especially in spasms of angina pectoris, asthma and general convulsions.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.), by inhalation.

***Amylum** (*Amyl.*), *Starch*, U. S. P. (Corn Starch).

ACTION AND USES: Dusting powder and diluent.

Glyceritum Amyli (*Glycer. Amyl.*), *Glycerite of Starch*, U. S. P.—Starch (10%) and water (10%) in glycerin.

USES: Emollient and excipient.

Anethol (*Anethol*), *Anethol*, N. F.—The chief constituent of oil of anise and oil of fennel.

A colorless or yellowish liquid, with a sweet taste and an odor of anise; solid below 20 C. Almost insoluble in water, but freely soluble in alcohol or ether.

ACTION AND USES: Carminative, etc. Similar to but without advantage over oil of anise.

DOSAGE: 0.2 Cc. or 3 minims (N. F.).

Elixir Anisi (*Elix. Anis.*), *Elixir of Anise*, N. F.—Anethol, oil of fennel, spirit of bitter almond, alcohol, syrup and water. Absolute alcohol content about 23 per cent.

USES: Carminative and flavor.

DOSAGE: For infants, 1 Cc. or 15 minims (N. F.).

Angelicae Fructus (*Angel. Fruct.*), *Angelica Fruit*, N. F. (*Angelica Seed*).

ACTION AND USES: Aromatic stomachic and carminative; without advantage over gentian.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Angelicae Radix (*Angel. Rad.*), *Angelica Root*, N. F.

ACTION AND USES: Stimulating expectorant; without advantage over tolu.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Angelicae Radicis (*Fldext. Angel. Rad.*),
Fluidextract of Angelica Root, N. F.—Angelica root
(100%). Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Anisum (*Anis.*), *Anise*, U. S. P. (*Aniseed*).

ACTION AND USES: Carminative. (See *Oleum Anisi*).

***Antimonii et Potassii Tartras** (*Antim. et Pot. Tart.*), *Antimony and Potassium Tartrate*, U. S. P. (*Antimonyl Potassium Tartrate*, *Tartrated Antimony*, *Tartar Emetic*).

A white, odorless powder with a sweet, metallic taste. Soluble in water (1:12) and in glycerin (1:15); insoluble in alcohol.

ACTION AND USES: Nauseant expectorant and emetic. Emetic doses and the prolonged use of expectorant doses are somewhat dangerous. Externally, a pustulant.

DOSAGE: Expectorant, 0.005 Gm. or $\frac{1}{12}$ grain (U. S. P.), in solution. It is best to begin with small doses (0.001 Gm. or $\frac{1}{60}$ grain) which may be repeated hourly, care being taken to avoid too great depression.

Vinum Antimonii (*Vin. Antimon.*), *Wine of Antimony*, N. F.—Antimony and potassium tartrate (0.4%), in water and sherry wine. Absolute alcohol content about 20 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Antimonii Oxidum (*Antim. Oxid.*), *Antimony Oxide*, N. F.—Antimonous oxide, Sb_2O_3 .

A white or grayish-white, odorless, tasteless powder. Insoluble in water and in alcohol.

ACTION AND USES: Obsolete nauseant, emetic and cathartic. Action uncertain and possibly toxic.

DOSAGE: 0.06 Gm. or 1 grain (N. F.).

Pulvis Antimonialis (*Pulv. Antimon.*), *Antimonial Powder*, N. F. (*James' Powder*).—Antimonous oxide (33%) and precipitated calcium phosphate.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Antimonium Sulphuratum (*Antim. Sulphurat.*), *Sulphurated Antimony*, N. F. (*Antimonium Oxysulphuratum*, *Kermes Mineral*).—Chiefly antimony trisulphide, Sb_2S_3 .

A red-brown, odorless, tasteless powder. Insoluble in water and in alcohol.

ACTION AND USES: Mineral cathartic, without advantage over calomel. Has no special action on the bile, as was formerly supposed.

Pilulae Antimonii Compositae (*Pil. Antimon. Co.*), *Compound Pills of Antimony*, N. F. (Plummer's Pills).—Each pill contains sulphurated antimony and mild mercurous chloride (each 0.04 Gm. or $\frac{2}{3}$ grain), guaiac (0.08 Gm. or $1\frac{1}{2}$ grains), and castor oil.

DOSAGE: 1 pill (N. F.).

**Antipyrina* (*Antipyr.*), *Antipyrine*, U. S. P. (*Phenazone*, *Phenyldimethylpyrazolon*).

White, odorless powder, with a slightly bitter taste. Freely soluble in water (1:1) and in alcohol (1:1.3).

ACTION AND USES: Antipyretic and analgesic, similar to acetanilid.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.), in solution, given with even greater caution than acetanilid and acetphenetidin.

Apii Fructus (*Apii Fruct.*), *Celery Fruit*, N. F. (*Celery Seed*).

ACTION AND USES: Domestic emmenagogue, depending for its action on the volatile oil.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Apii Fructi (*Fldext. Apii Fruct.*), *Fluidextract of Celery Fruit*, N. F.—Celery fruit (100%). Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Apocynum (*Apocyn.*), *Apocynum*, N. F. (*Canadian Hemp*).—Rhizome and roots.

ACTION AND USES: Cardiac tonic of digitalis group; unreliable as to rate of absorption.

DOSAGE: 0.75 Gm. or 12 grains (N. F.).

Fluidextractum Apocyni (*Fldext. Apocyn.*), *Fluidextract of Apocynum*, N. F. (Fluidextract of Canadian Hemp).—Apocynum (100%). Absolute alcohol content about 50 per cent.

DOSAGE: 0.75 Cc. or 12 minims (N. F.).

**Apomorphinae Hydrochloridum* (*Apomorph. Hydrochl.*), *Apomorphine Hydrochloride*, U. S. P. (*Apomorphine Chloride*).

White powder, becoming greenish on exposure to light and air, odorless and with a slightly bitter taste. Soluble in water (1:50) and in alcohol (1:50).

ACTION AND USES: Prompt, centrally acting emetic, especially adapted for hypodermic administration. Rarely used as nauseant expectorant.

DOSAGE: Expectorant 0.003 Gm. or $\frac{1}{20}$ grain (U. S. P.). Probably too large; 0.004 Gm. or $\frac{1}{15}$ grain is said to have produced death in a person enfeebled by chronic bronchitis. From 0.001 Gm. to 0.002 Gm., or $\frac{1}{60}$ to $\frac{1}{30}$ grain, repeated once an hour or once in two hours, is safer. Emetic, by mouth (hypodermic administration is better), 0.01 Gm. or $\frac{1}{6}$ grain (U. S. P.). Emetic, hypodermic, 0.005 Gm. or $\frac{1}{12}$ grain (U. S. P.). This may be repeated at ten-minute intervals until effective, but it should be remembered that in some cases apomorphin produces toxic effects without causing vomiting.

***Aqua (Aqua), Water, U. S. P.—H₂O.**

ACTION AND USES: Solvent. The ingestion of large amounts is a very valuable eliminative measure.

***Aqua Destillata (Aq. Dest.), Distilled Water, U. S. P.—**Water obtained by distillation.

Aqua Destillata Sterilisata (Aq. Dest. Steril.), Sterilized Distilled Water, U. S. P.—Should be used within forty-eight hours after its preparation.

AQUAE AROMATICAE, AROMATIC WATERS.—The official aromatic waters are aqueous solutions of volatile substances, usually volatile oils. They are used as vehicles for the more active water-soluble drugs. For preparations included in this list see under the names of the principal constituents. For the following, see under the names of the respective oils: Aqua Amygdalae Amarae, Aqua Foeniculi, Aqua Menthae Piperitae and Aqua Menthae Viridis.

Aralia (Aralia), Aralia, N. F. (American Spikenard. Spignet).—Rhizome and roots.

ACTION AND USES: Obsolete irritant, diaphoretic and "alterative." Probably without value.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Araliae (Fldext. Aral.), Fluidextract of Aralia, N. F.—Aralia (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Argenti Nitras (Arg. Nit.), Silver Nitrate, U. S. P.—AgNO₃.**

Colorless plates, darkening on exposure to light in the presence of organic matter; odorless; strongly caustic and having a bitter, metallic taste. Very soluble in water (1: 0.4) and soluble in alcohol (1: 30). Incompatible especially with chlorides and organic matter.

ACTION AND USES: Externally as a caustic, antiseptic and germicide; internally as an astringent.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.), diluted or in kaolin pills. The concentrations of solutions for local use vary from 0.01 to 10 per cent., according to the sensitiveness of the surface and the degree of action desired. Distilled water should be used in making the solutions.

**Argenti Nitras Fusus* (*Arg. Nit. Fus.*), *Molded Silver Nitrate*, U. S. P. (Fused Silver Nitrate, Lunar Caustic).— AgNO_3 toughened by the addition of a small proportion of silver chloride.

White, hard pencils or cones.

Argenti Oxidum (*Arg. Oxid.*), **Silver Oxide**, U. S. P.— Ag_2O .

A heavy, odorless, brownish-black powder with a metallic taste. Only slightly soluble in water; insoluble in alcohol.

ACTION AND USES: Astringent in gastric ulcer. Probably inefficient.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Arnica (*Arnic.*), **Arnica**, U. S. P. (*Arnica Flowers*).

ACTION AND USES: Feeble rubefacient. Still used externally to some extent in the form of a tincture mixed with water as an evaporating lotion. The chief active ingredient of the tincture is probably its alcohol.

Fluidextractum Arnicae (*Fldest. Arnic.*), *Fluidextract of Arnica*, N. F.—Arnica (100%). Absolute alcohol content about 40 per cent.

Tinctura Arnicae (*Tr. Arnic.*), *Tincture of Arnica*, U. S. P.—Arnica (20%) in diluted alcohol. Absolute alcohol content about 46 per cent.

Arseni Iodidum (*Arsen. Iod.*), **Arsenous Iodide**, U. S. P. (*Arsenic Iodide*).— AsI_3 .

An orange-red, odorless powder. Soluble in water (1:12) with partial decomposition; soluble in alcohol.

ACTION AND USES: Similar to those of arsenic trioxide, over which it has no advantage.

DOSAGE: 0.005 Gm. or $\frac{1}{2}$ grain (U. S. P.).

**Liquor Arseni et Hydrargyri Iodidi* (*Liq. Arsen. et Hydrarg. Iod.*), *Solution of Arsenous and Mercuric Iodide*, U. S. P. (Donovan's Solution).—Arsenous iodide and red mercuric iodide (each 1%) in water.

DOSAGE: 0.1 Cc. or $1\frac{1}{2}$ minims (U. S. P.).

***Arseni Trioxidum** (*Arsen. Triox.*), **Arsenic Trioxide**, U. S. P. (*Arsenous Acid*, *Arsenous Oxide*, *White Arsenic*).— As_2O_3 .

Amorphous, glassy masses or white, crystalline masses or powder. Slowly soluble in water (about 1:100), slightly soluble in alcohol and ether and freely soluble in glycerine. Readily dissolved by hydrochloric acid or by alkaline solutions.

ACTION AND USES: Believed to modify nutrition. Used as tonic in anemias, nervous diseases, etc., in skin diseases and in some forms of malaria. Externally, a mild escharotic, but the danger of poisoning prevents its general use. Likely to produce nephritis. A very common poison.

DOSAGE: 0.002 Gm. or $\frac{1}{30}$ grain (U. S. P.), in pills or solutions.

***Liquor Acidi Arsenosi (Liq. Acid. Arsen.), Solution of Arsenous Acid, U. S. P.** (Hydrochloric Solution of Arsenic, "Solution of Arsenic Chloride").—Arsenic trioxide (1%) in diluted hydrochloric acid (5%) and water.

DOSAGE: 0.2 Cc. or 3 minims.

***Asafoetida (Asafoet.), Asafetida, U. S. P. (Gum Asafetida).**
—A gum resin.

Should be 60% soluble in alcohol.

ACTION AND USES: Used as a carminative and also in the treatment of hysteria, the action being psychic.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), in pills.

Emulsum Asafoetidae (Emuls. Asafoet.), Emulsion of Asafetida, U. S. P. (Milk of Asafetida).—Asafetida (4%) in water.

USES: Used especially as enema in tympanites.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Pilulae Asafoetidae (Pil. Asafoet.), Pills of Asafetida, U. S. P.—Each pill contains asafoetida (0.2 Gm. or 3 grains), and soap.

DOSAGE: 2 pills (U. S. P.).

Tinctura Asafoetidae (Tr. Asafoet.), Tincture of Asafetida, U. S. P.—Asafetida (20%) in alcohol. Absolute alcohol content about 81 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Asarum (Asar.) Asarum, N. F. (Canada Snake-Root. Wild Ginger).—Rhizome.

ACTION AND USES: Carminative and flavor; similar to but without advantage over ginger.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Syrupus Asari Compositus (Syr. Asar. Co.), Compound Syrup of Asarum, N. F.—Asarum (6.2%), fluidextract

of ipecac (0.3%), alcohol, cochineal and potassium carbonate in syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Asclepias (Asclep.), Asclepias, N. F. (Pleurisy Root).

ACTION AND USES: Obsolete domestic remedy for pleurisy, also used as diaphoretic. Probably without value.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Asclepiadis (Fldext. Asclepiad.), Fluidextract of Asclepias, N. F.—Asclepias (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Aspidium (Aspidium), Aspidium, U. S. P. (Male Fern).—**The rhizome of certain ferns.

ACTION AND USES: Used as a leniacide in the form of oleoresin.

**Oleoresina Aspidii (Oleores. Aspid.) Oleoresin of Aspidium, U. S. P. (Oleoresin of Male Fern).*

DOSAGE: 2 Gm. or 30 grains (total dose in a day, U. S. P.), which may be given singly or in repeated, divided doses, at intervals of fifteen minutes, in capsules containing 0.3 Gm. or 5 grains, or in emulsion. Larger doses may be given to robust persons. Special caution is necessary in giving it to anemic or debilitated persons. The patient should be prepared by a light diet or fasting for twenty-four hours. The drug should be given early in the morning, preceded by a saline cathartic, and the last dose followed in three hours by a saline laxative. Caution:—Castor oil or other fixed oils should not be used in connection with oleoresin of aspidium, as they favor the absorption of the active principle.

Aspidosperma (Aspidosp.), Aspidosperma, U. S. P. (Quebracho).—A bark.

ACTION AND USES: Has been recommended in dyspneic conditions. Value doubtful.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Fluidextractum Aspidospermatis (Fldext. Aspidosp.), Fluidextract of Aspidosperma, U. S. P. (Fluidextract of Quebracho).—Aspidosperma (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

***Atropina (Atrop.), Atropine, U. S. P.—**An alkaloid obtained from belladonna and some other plants of the nightshade family.

A white, odorless powder. Slightly soluble in water (1:455), but freely soluble in alcohol (1:2).

ACTION AND USES: Atropine and the related alkaloids paralyze the parasympathetic endings. They are used especially as mydriatics and cycloplegics, to suppress secretions, as in gastric hyperacidity, and to quicken the heart, regulate peristalsis, lessen bronchial spasm, etc.

DOSAGE: 0.5 mg. or $\frac{1}{120}$ grain (U. S. P.).

Oleatum Atropinae (*Oleat. Atrop.*), *Oleate of Atropine*, N. F.—Atropine (2%), in alcohol, oleic acid and olive oil.

***Atropinae Sulphas** (*Atrop. Sulph.*), *Atropine Sulphate*, U. S. P.

White, odorless powder. Very soluble in water (1:0.4) and in alcohol (1:5).

DOSAGE: 0.5 mg. or $\frac{1}{120}$ grain (U. S. P.).

Aurantii Amari Cortex (*Aurant. Amar. Cort.*), *Bitter Orange Peel*, U. S. P.

ACTION AND USES: Aromatic bitter stomachic, used principally as a flavoring.

Fluidextractum Aurantii Amari (*Fldext. Aurant. Amar.*), *Fluidextract of Bitter Orange Peel*, U. S. P.—Bitter orange peel (100%). Absolute alcohol content about 60 per cent.

Tinctura Aurantii Amari (*Tr. Aurant. Amar.*), *Tincture of Bitter Orange Peel*, U. S. P.—Bitter orange peel (20%) in alcohol and water. Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Vinum Aurantii Compositum (*Vin. Aurant. Co.*), *Compound Wine of Orange*, N. F. (*Elixir Aurantiorum Compositum*, *Compound Elixir of Orange*).—Bitter orange peel, absinthium, menyanthes, cascarilla, Saigon cinnamon and gentian, extracted with potassium carbonate in sherry wine. Absolute alcohol content about 20 per cent.

USES: Objectionable alcoholic "bitter."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Aurantii Dulcis Cortex (*Aurant. Dulc. Cort.*), *Sweet Orange Peel*, U. S. P.

ACTION AND USES: Flavoring agent.

Syrupus Aurantii (*Syr. Aurant.*), *Syrup of Orange*, U. S. P.—*Tincture of sweet orange peel in syrup.*

Tinctura Aurantii Dulcis (*Tr. Aurant. Dulc.*), *Tincture of Sweet Orange Peel*, U. S. P.—Sweet orange peel from fresh fruit (50%) in alcohol. Absolute alcohol content about 83 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Aurantii Flores, Bitter Orange Flowers.

ACTION AND USES: Flavoring agent.

Aqua Aurantii Florum (*Aq. Aurant. Flor.*), *Orange Flower Water*, U. S. P.—Stronger orange flower water in water.

Aqua Aurantii Florum Fortior (*Aq. Aurant. Flor. Fort.*), *Stronger Orange Flower Water*, U. S. P.—Prepared by distillation of bitter orange flowers with water. Must be free from mucilaginous growths.

Syrupus Aurantii Florum (*Syr. Aurant. Flor.*), *Syrup of Orange Flowers*, U. S. P.—Sugar in orange flower water.

Auri et Sodii Chloridum (*Aur. et Sod. Chlor.*), **Gold and Sodium Chloride**, U. S. P.—Equal parts of AuCl_3 and NaCl .

An orange-yellow, odorless, deliquescent powder with a saline, metallic taste. Very soluble in water.

ACTION AND USES: Preparations of gold have been employed in medicine for the most varied conditions without any real evidence of usefulness.

DOSAGE: 0.005 Gm. or $\frac{1}{12}$ grain (U. S. P.).

***Balsamum Peruvianum** (*Bals. Peruv.*), **Balsam of Peru**, U. S. P. (Peru Balsam).

Dark brown, viscid liquid, with a vanilla-like odor and a bitter, acrid taste. Soluble in alcohol or chloroform; only partially soluble in ether or petroleum benzin; practically insoluble in water.

ACTION AND USES: Externally in the form of ointments or alcoholic solutions as a stimulant to indolent wounds and ulcers and in the treatment of scabies.

***Balsamum Tolutanum** (*Bals. Tolu*), **Balsam of Tolu**, U. S. P. (Tolu Balsam).

Yellowish-brown or brown, plastic solid, becoming brittle when old, dried or exposed to cold, and having a vanilla-like odor and a mild, aromatic taste. Insoluble in water but soluble in alcohol, chloroform and ether.

ACTION AND USES: Has been recommended as an expectorant; used chiefly in the form of the syrup, which is little more than a pleasantly flavored vehicle.

**Syrupus Tolutanus* (*Syr. Tolu.*), *Syrup of Tolu*, U. S. P.—*Tincture of tolu* (5%) in syrup.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Tinctura Tolutana (*Tr. Tolu.*), *Tincture of Tolu*, U. S. P.—Balsam of tolu (20%) in alcohol. Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Baptisia (*Baptis.*), **Baptisia**, N. F. (Wild Indigo Root).

ACTION AND USES: Without any definite known action; formerly used in typhoid fever and diseases showing similar symptoms.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Baptisiae (*Fldext. Baptis.*), *Fluidextract of Baptisia*, N. F.—Baptisia (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Belladonnae Folia** (*Bellad. Fol.*), **Belladonna Leaves**, U. S. P. (Deadly Nightshade Leaves, *Belladonnae Folium*, P. I.).—Yield not less than 0.3 per cent. of mydriatic alkaloids.

ACTION AND USES: Those of the hyoscyamine and atropine (which see) contained therein.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

**Emplastrum Belladonnae* (*Emp. Bellad.*), *Belladonna Plaster*, U. S. P.—Extract of belladonna leaves (30%), yielding about 0.37 per cent. of alkaloids.

ACTION AND USES: Applied in the treatment of rheumatic and similar pains.

**Extractum Belladonnae Foliorum* (*Ext. Bellad. Fol.*), *Extract of Belladonna Leaves*, U. S. P.—One Gm. extract represents about 4 Gm. belladonna leaves and yields about 1.25 per cent. of alkaloids. Two forms: pilular extract of belladonna leaves and powdered extract of belladonna leaves.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.).

**Tinctura Belladonnae Foliorum* (*Tr. Bellad. Fol.*), *Tincture of Belladonna Leaves*, U. S. P. (*Belladonnae Tinctura*, P. I.).—Belladonna leaves (10%) yielding about 0.03 per cent. of alkaloids in alcohol. Absolute alcohol content about 47 per cent.

DOSAGE: 0.75 Cc. or 12 minims (U. S. P.).

**Unguentum Belladonnae* (*Ung. Bellad.*), *Belladonna Ointment*, U. S. P.—Pilular extract of belladonna leaves (10%) in diluted alcohol, hydrous wool fat and benzoinated lard.

Belladonnae Radix (Bellad. Rad.), **Belladonna Root**, U. S. P.
(Deadly Nightshade Root).—Yields not less than 0.45 per cent. of mydriatic alkaloids.

ACTION AND USES: Those of the hyoscyamine and atropine (which see) contained therein.

DOSAGE: 0.045 Gm. or $\frac{3}{4}$ grain (U. S. P.).

Fluidextractum Belladonnae Radicis (Fldext. Bellad. Rad.), *Fluidextract of Belladonna Root*, U. S. P.—Belladonna root (100%) yielding about 0.45 per cent. of alkaloids. Absolute alcohol content about 75 per cent.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.).

Linimentum Belladonnae (Lin. Bellad.), *Belladonna Liniment*, U. S. P.—Fluidextract of belladonna root (about 95%) and camphor (5%). Absolute alcohol content about 70 per cent.

USES: A mildly analgesic and rubefacient application.

Benzaldehydum (Benzaldehyd.), **Benzaldehyde**, U. S. P.—Produced synthetically or obtained from oil of bitter almonds.

A colorless or yellowish liquid, with a bitter-almond odor and a burning, aromatic taste. Slightly soluble in water; miscible with alcohol, ether, and fixed or volatile oils.

ACTION AND USES: Flavoring agent.

DOSAGE: 0.03 Cc. or $\frac{1}{2}$ minim (U. S. P.).

Benzinum Purificatum (Benzin. Purif.), **Purified Petroleum Benzin**, U. S. P. (Petroleum Ether).

A clear, colorless liquid, with an ethereal or faint petroleum odor; highly inflammable. Practically insoluble in water, freely soluble in alcohol and miscible with ether, chloroform, volatile oils and fixed oils with the exception of castor oil.

ACTION AND USES: Pharmaceutical solvent.

***Benzoinum** (Benzoin.), **Benzoin**, U. S. P. (Gum Benjamin).—A balsamic resin.

ACTION AND USES: Used locally to promote healing, and in inhalations as an expectorant.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Tinctura Benzoini (Tr. Benz.), *Tincture of Benzoin*, U. S. P.—Benzoin (20%) in alcohol. Absolute alcohol content about 83 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

***Tinctura Benzoini Composita** (Tr. Benz. Co.), *Compound Tincture of Benzoin*, U. S. P.—Benzoin (10%) aloes

(2%), storax (8%), balsam of tolu (4%), in alcohol. Absolute alcohol content about 80 per cent.

USES: Protective and local stimulant, especially by inhalation.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

***Benzosulphinidum (Benzosulphinid.), Benzosulphinide, U. S. P. (Saccharin, Glusidum).**

A white, nearly odorless powder with an intensely sweet taste even in dilute solutions. Slightly soluble (1:290) in water and soluble (1:31) in alcohol and readily dissolved by ammonia water or by a solution of alkali hydroxides or of sodium bicarbonate.

ACTION AND USES: Sweetening agent in diabetes mellitus; about 500 times as sweet as sugar, weight for weight.

Berberis (Berber.), Berberis, N. F. (Oregon Grape Root).—Rhizome and roots.

ACTION AND USES: Probably similar to, but without advantage over, hydrastis.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Berberidis (Fldext. Berberid.), Fluidextract of Berberis, N. F.—Berberis (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Betaeucainae Hydrochloridum (Betaeucain. Hydrochl.), Betaeucaine Hydrochloride, U. S. P. (Eucaine Chloride, Eucaine; also known as Benzamine Hydrochloride).

A white, odorless powder. Soluble in water (1:30) and in alcohol (1:35).

ACTION AND USES: Local anesthetic.

DOSAGE: In a 2 or 3 per cent. solution for the eye or 5 to 10 per cent. solution for the nose and throat.

***Betanaphthol (Betanaph.), Betanaphthol, U. S. P. (Naphthol).**

Colorless or pale buff crystalline laminae or white or yellowish powder, with a faint phenol odor and a pungent taste. Only slightly soluble in water (1:1000) and very soluble in alcohol (1:0.8).

ACTION AND USES: Antiseptic, somewhat stronger than phenol and less toxic. Used in skin affections, and as intestinal antiseptic and anthelmintic for hookworm.

DOSAGE: Internally, 0.25 Gm. or 4 grains (U. S. P.). Externally as a 3 to 5 per cent. ointment.

Pasta Betanaphtholis (Past. Betanaphthol.), Betanaphthol Paste, N. F. (Lassar's Naphthol Paste).—Betanaphthol

(10%) and precipitated sulphur (50%) in petrolatum and soft soap.

Petroxolinum Betanaphtholis (*Petrox. Betanaphthol.*), *Betanaphthol Petroxolin*, N. F. (*Betanaphthol Petrox*).—*Betanaphthol* (10%) in liquid petroxolin.

Bismuthi Betanaphtholas (**Bism. Betanaph.**), **Bismuth Betanaphthol**, U. S. P. (**Bismuth Betanaphtholate**; also sold as *Orphol*).—Of somewhat varying composition.

A buff-colored or grayish-brown, nearly odorless, tasteless powder. Nearly insoluble in water or alcohol.

ACTION AND USES: Used as an intestinal antiseptic, being decomposed into its constituents in the intestine, where it exerts the effects of bismuth salts and of betanaphthol.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Bismuthi et Ammonii Citras (**Bism. et Ammon. Cit.**), **Bismuth and Ammonium Citrate**, U. S. P. (**Bismuth Ammonio-citrate**).

Shining, pearly or translucent, odorless scales or white powder, with a metallic taste. Very soluble in water and sparingly soluble in alcohol.

ACTION AND USES: Somewhat astringent and irritating; should not be used as a substitute for the insoluble salts of bismuth. Its use in "bismuth elixirs" is irrational.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Bismuthi et Sodii Tartras, **Bismuth and Sodium Tartrate**.

USES: An irrational method of administering bismuth.

Elixir Bismuthi (*Elix. Bismuth.*), *Elixir of Bismuth*, N. F.—Glycerite of bismuth (12.5%), in glycerin, water and aromatic elixir. Absolute alcohol content about 10 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Cinchonae Alkaloidorum, Ferri et Bismuthi (*Elix. Cinchon. Ferr. et Bism.*), *Elixir of Cinchona Alkaloids, Iron and Bismuth*, N. F. (*Elixir Cinchonae, Ferri et Bismuthi*, N. F. III, *Elixir of Calisaya, Alkaloidal, with Iron and Bismuth*).—Glycerite of bismuth (6.5%), elixir of cinchona, alkaloids and iron (about 87%) and water. Absolute alcohol content about 17 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Glyceritum Bismuthi (*Glycer. Bismuth.*), *Glycerite of Bismuth*, N. F.—A solution of bismuth and sodium tartrate.

Liquor Bismuthi (Liq. Bism.), Solution of Bismuth, N. F.—Glycerite of bismuth (12.5%), in alcohol and water. Absolute alcohol content about 12 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm. (N. F.).

Bismuthi Oxidum, Bismuth Oxide.

Magma Bismuthi (Magma Bism.), Bismuth Magma, U. S. P. (Milk of Bismuth).—A suspension in water of bismuth hydroxide and bismuth subcarbonate, forming a thick, white opaque liquid.

USES: Used similarly to bismuth subcarbonate, over which it has no advantage.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

***Bismuthi Subcarbonas (Bism. Subcarb.), Bismuth Subcarbonate, U. S. P.**—Of varying chemical composition.

White or nearly white, odorless, tasteless powder. Insoluble in water or alcohol.

ACTION AND USES: Used like other insoluble salts of bismuth as a protective and healing agent against wounds, diarrheas, etc., and in x-ray work.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.) in powders or cachets or in suspension (best form).

***Bismuthi Subgallas (Bism. Subgall.), Bismuth Subgallate, U. S. P. (Dermatol).**—A basic bismuth gallate of varying chemical composition.

Bright yellow, odorless, tasteless powder. Insoluble in water or alcohol.

ACTION AND USES: Similar to those of bismuth subcarbonate, over which it has no advantage.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

***Bismuthi Subnitras (Bism. Subnit.), Bismuth Subnitrate, U. S. P.**—Basic bismuth nitrate of varying chemical composition.

A heavy, white, odorless, almost tasteless powder. Almost insoluble in water and insoluble in alcohol. In aqueous mixtures, incompatible with alkali carbonates because of the liberation of nitric acid.

ACTION AND USES: Somewhat more astringent than the subcarbonate, because of the liberation of nitric acid.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.). Preferably administered as a powder or as cachets.

Bismuthi Subsali-cylas (Bism. Subsali-cyl.), Bismuth Subsalicylate, U. S. P.—Basic bismuth salicylate, of varying chemical composition.

A white or nearly white, odorless, tasteless powder. Almost insoluble in water or alcohol.

ACTION AND USES: Protective and astringent like other insoluble salts of bismuth, over which it has no marked advantage.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Boldo (Boldo), Boldo, N. F. (Boldo Leaves).

ACTION AND USES: Aromatic, perhaps somewhat diuretic. No well defined indications for its use.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Fluidextractum Boldi (Fldext. Boldi), Fluidextract of Boldo, N. F.—Boldo (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

Brayera (Brayer.), Brayera, N. F. (Kousso, Cusso).—Flowers.

ACTION AND USES: Teniacide; without advantage over aspidium or pelletierine.

DOSAGE: 15 Gm. or 4 drachms (N. F.).

Infusum Brayerae (Inf. Brayer.), Infusion of Brayera, N. F.—Brayera (6%) in water.

DOSAGE: 250 Cc. or 8 fluidounces (N. F.).

Bromoformum (Bromof.), Bromoform, U. S. P.—CHBr₃.

A heavy, clear, colorless liquid, with an ethereal odor and a sweet taste resembling that of chloroform. Slightly soluble in water and miscible with alcohol, chloroform, ether or fixed or volatile oils.

ACTION AND USES: Has been recommended as a sedative in the treatment of whooping-cough. Opinions differ as to its value.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.), preferably dissolved in almond oil or dropped on sugar. Bromoform is too dense to be dispensed safely with aqueous vehicles.

Bromum (Bromum), Bromine, N. F.—Br.

Heavy, dark, brownish-red, volatile liquid, with a suffocating odor resembling that of chlorine. Soluble in water (1:90) and freely soluble in alcohol or ether.

ACTION AND USES: Violent irritant; seldom used in medicine.

Liquor Bromi (Liq. Brom.), Solution of Bromine, N. F., (Smith's Solution of Bromine).—Bromine (8.3%), potassium bromide and water.

USES: Antiseptic and irritant.

Bryonia (Bryon.), Bryonia, N. F. (Bryony).—Root.

ACTION AND USES: Drastic resinous cathartic; without advantage over jalap. Used in dropsies, pleurisy, neuralgia, etc.; of doubtful value.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Tinctura Bryoniae (Tr. Bryon.), Tincture of Bryonia, N. F.—Bryonia (10%) in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Buchu (Buchu), Buchu, U. S. P. (Buchu Leaves, Short Buchu, Long Buchu).

ACTION AND USES: At one time largely used as antiseptic and diuretic, especially in catarrhal cystitis. Value doubtful.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Elixir Buchu (Elix. Buchu), Elixir of Buchu, N. F.—Fluidextract of buchu (12.5%) in alcohol and aromatic elixir. Absolute alcohol content about 35 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm.

Elixir Buchu Compositum (Elix. Buchu Co.), Compound Elixir of Buchu, N. F.—Compound fluidextract of buchu (25%) in aromatic elixir. Absolute alcohol content about 30 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm.

Elixir Buchu et Potassii Acetatis (Elix. Buchu et Pot. Acet.), Elixir of Buchu and Potassium Acetate, N. F.—Potassium acetate (8.5%) and elixir of buchu. Absolute alcohol content about 30 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm.

Fluidextractum Buchu (Fldext. Buchu), Fluidextract of Buchu, U. S. P.—Buchu (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Fluidextractum Buchu Compositum (Fldext. Buchu Co.), Compound Fluidextract of Buchu, N. F.—Buchu (62.5%), cubeb, juniper and uva ursi (each 12.5%). Absolute alcohol content about 55 per cent.

USES: A needlessly complex diuretic mixture.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Cacao Praeparata (Cacao Praep.), Cocoa, N. F. (Prepared Cacao, Soluble Cocoa).

ACTION AND USES: Beverage and flavor.

Cactus Grandiflorus (Cact. Grand.), **Cactus Grandiflorus**, N. F. (Night Blooming Cereus).—Fresh stems.

ACTION AND USES: Used in neurotic heart disturbances; results probably imaginary. Drug apparently inactive.

Tinctura Cacti Grandiflori (Tr. Cact. Grand.), **Tincture of Cactus Grandiflorus**, N. F.—Cactus grandiflorus (50%) in alcohol. Absolute alcohol content about 65 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Caffeina** (Caffein.), **Caffeine**, U. S. P. (Theine).—An alkaloid obtained from coffee or tea.

White, silky, efflorescent needles, odorless and bitter. Soluble in water (1:46) and in alcohol (1:66).

ACTION AND USES: Diuretic; cardiac, respiratory and psychic stimulant. Used as a nerve stimulant in poisoning by narcotic drugs and for the relief of headache. Widely used to combat the circulatory depression of the acute infectious diseases but its value in such conditions is uncertain. As a diuretic probably inferior to theobromine.

DOSAGE: 0.15 Gm. or $2\frac{1}{2}$ grains (U. S. P.), in capsules.

Caffeina Citrata (Caffein Cit.), **Citrated Caffeine**, U. S. P.—Caffeine and citric acid equal parts.

White, odorless powder, with a slightly bitter, acid taste.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.), in solution or capsules.

Caffeina Citrata Effervescens (Caff. Cit. Eff.), **Effervescent Citrated Caffeine**, U. S. P.—Contains anhydrous caffeine (about 2%), with sodium tartrate and citrate.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.) in water.

***Caffeinae Sodio-Benzoeas** (Caff. Sod. Benz.), **Caffeine Sodio-Benzoeate**, U. S. P.—Caffeine and sodium benzoate about equal parts.

White, odorless powder, with a slightly bitter taste. Freely soluble in water (1:1.1) and soluble in alcohol (1:30); partly soluble in chloroform.

ACTION AND USES: The form of caffeine usually employed for hypodermic administration, since it is freely soluble.

DOSAGE: By mouth, 0.3 Gm. or 5 grains (U. S. P.); hypodermic, 0.2 Gm. or 3 grains (U. S. P.).

Caffeinae Sodio-Salicylas (Caff. Sod. Salicyl.), **Caffeine Sodio-Salicylate**, N. F.—Represents equal parts of caffeine and sodium salicylate.

White, odorless powder. Freely soluble in water (1:2).

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Calamina Praeparata (Calamin. Praep.), Prepared Calamine, N. F. (Lapis Calaminaris).—Native zinc carbonate.

A pink powder. Insoluble in water.

ACTION AND USES: Protective, similar to zinc oxide, over which it has no therapeutic advantage; has the cosmetic advantage of being slightly pink.

Unguentum Calaminae (Ung. Calamin.), Calamine Ointment, N. F., (*Unguentum Zinci Carbonatis Crudi*, *Unguentum Calaminare*, Turner's Cerate).—Prepared calamine (17%) in ointment.

Calcii Bromidum (Calc. Brom.), Calcium Bromide, U. S. P.—Hydrated calcium bromide (not less than 84 per cent. of CaBr_2).

A white, odorless, deliquescent powder, with a sharp, saline taste. Very soluble in water (1:0.7) and in alcohol (1:1.3). Incompatible with alkali carbonates and phosphates.

ACTION AND USES: Similar to but more irritant than sodium bromide, over which it has ordinarily no advantage, although the calcium perhaps increases the sedative action.

DOSAGE: 1 Gm. or 15 grains, preferably administered in aqueous solutions, being too deliquescent for use in powder or cachets.

Elixir Calcii Bromidi (Elix. Calc. Brom.), *Elixir of Calcium Bromide*, N. F.—Calcium bromide (8.5%) diluted hydrobromic acid (0.4%), syrup, distilled water and aromatic elixir. Absolute alcohol content about 7 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Calcii Carbonas Praecipitatus** (Calc. Carb. Praec.), Precipitated Calcium Carbonate, U. S. P. (Precipitated Chalk).— CaCO_3 .

A fine, white, odorless, tasteless powder. Nearly insoluble in water and insoluble in alcohol. Decomposed by acids with evolution of carbon dioxide.

ACTION AND USES: Chiefly used as a basis for tooth powder. Internally, employed as an antacid in diarrheas, etc.; externally in mixtures with cream or oil, has been used as an application to burns as an antacid and protective.

DOSAGE: 1 Gm. or 15 grains (U. S. P.), as powder or in suspension.

***Calcii Chloridum (Calc. Chlor.), Calcium Chloride, U. S. P.**
—Hydrated calcium chloride (not less than 75 per cent. of CaCl_2).

White, hard, deliquescent, odorless masses or granules, with a sharp, saline taste. Very soluble in water (1:0.62), and freely soluble in alcohol (1:10).

ACTION AND USES: Used for the oral administration of calcium. Employed in hemorrhagic conditions, in which it is probably useless. Has also been advised in inflammatory conditions of the skin and mucous membranes (urticaria, serum-rashes, hay-fever, etc.).

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in solution.

Calcii Glycerophosphas (Calc. Glycerophos.), Calcium Glycerophosphate, U. S. P. (Calcium Glycerinophosphate).
— $\text{CaC}_6\text{H}_5\text{O}_6\text{P}$.

White, odorless, almost tasteless powder. Soluble in water (1:50) and insoluble in alcohol.

ACTION AND USES: Formerly believed to be a tonic, and used especially in neurasthenia, to improve the nutrition of the nervous system. There is no reliable evidence that it has any value.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), in solution.

Elixir Calcii et Sodii Glycerophosphatum (Elix. Calc. et Sod. Glycerophos.), Elixir of Calcium and Sodium Glycerophosphates, N. F. (Elixir Glycerophosphatum N. F. III).—Solution of sodium glycerophosphate (2.5%), calcium glycerophosphate (0.875%), phosphoric acid (0.8%), in glycerin, aromatic elixir and distilled water. Absolute alcohol content about 7 per cent.

USES: An irrational and inefficient "tonic" mixture.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Calcii Hypophosphis (Calc. Hypophos.), Calcium Hypophosphite, U. S. P.— $\text{Ca}(\text{PH}_2\text{O}_2)_2$.

Colorless crystals or white powder, odorless and having a nauseous, bitter taste. Freely soluble in water (1:6.5) and insoluble in alcohol.

ACTION AND USES: The hypophosphites were introduced as of special value in tuberculosis; later they were used as general reconstructive tonics, but there is no experimental or reliable clinical evidence that the hypophosphite radical has any value as such. Calcium hypophosphite must be regarded merely as a means of administering calcium.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.) in syrups.

Elixir Calcii Hypophosphitis (*Elix. Calc. Hypophos.*), *Elixir of Calcium Hypophosphite*, N. F.—Calcium hypophosphite (3.5%) in aromatic elixir. Absolute alcohol content about 20 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Hypophosphitum (*Elix. Hypophos.*), *Elixir of Hypophosphites*, N. F.—Calcium hypophosphite (5.25%), sodium hypophosphite and potassium hypophosphite (each 1.75%) in water, glycerin, compound spirit of cardamom and aromatic elixir. Absolute alcohol content about 12 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Hypophosphitum et Ferri.—See under *Ferri Hypophosphis*.

Liquor Hypophosphitum (*Liq. Hypophos.*), *Solution of Hypophosphites*, N. F.—Calcium hypophosphite (3.5%), sodium hypophosphite (2%) and potassium hypophosphite (1.75%) in water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Liquor Hypophosphitum Compositus (*Liq. Hypophos. Co.*), *Compound Solution of Hypophosphites*, N. F.—Calcium hypophosphite and potassium hypophosphite (each 0.85%), sodium hypophosphite (0.22%), ferric hypophosphite (0.44%) and manganese hypophosphite (0.22%), quinine hypophosphite (0.22%), strychnine hypophosphite (0.007%) and potassium citrate in orange flower water, glycerin and water.

USES: Complex and irrational mixture.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Calcii et Sodii Hypophosphitum (*Syr. Calc. et Sod. Hypophos.*), *Syrup of Calcium and Sodium Hypophosphites*, N. F.—Calcium hypophosphite and sodium hypophosphite (each 3.5%), in sugar and water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Calcii Hypophosphitis (*Syr. Calc. Hypophos.*), *Syrup of Calcium Hypophosphite*, N. F.—Calcium hypophosphite (3.5%) in sugar and water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Hypophosphitum (*Syr. Hypophos.*), *Syrup of Hypophosphites*, U. S. P.—Calcium hypophosphite (4.5%), potassium hypophosphite and sodium hypophosphite (each 1.5%) in glycerin and syrup.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Syrupus Hypophosphitum Compositus (Syr. Hypophos. Co.), *Compound Syrup of Hypophosphites*, N. F.—Calcium hypophosphite (3.5%), potassium hypophosphite and sodium hypophosphite (each 1.75%), ferric hypophosphite and manganese hypophosphite (each 0.225%), quinine (0.11%), strychnine hypophosphite (0.0123%) and sodium citrate (0.375%) in glycerin and syrup.

USES: Antiquated, complex and irrational "tonic."

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Calcii Iodidum, Calcium Iodide.

Syrupus Calcii Iodidi (Syr. Calc. Iodid.), *Syrup of Calcium Iodide*, N. F.—Represents calcium iodide (8.75%) in syrup.

USES: Superfluous, whether regarded as a calcium preparation or as an iodide preparation.

DOSAGE: 2 Cc. or 30 minims.

***Calcii Lactas (Calc. Lact.), Calcium Lactate, U. S. P.—**
 $\text{Ca}(\text{C}_2\text{H}_3\text{O}_2)_2 + 5\text{H}_2\text{O}$.

White, odorless, practically tasteless masses or powder. Soluble in water (1:20) and almost insoluble in alcohol.

ACTION AND USES: Used for the characteristic action of calcium; less irritating and therefore better adapted to hypodermic administration than calcium chloride (which see).

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in solution.

Calcii Lactophosphas (Calc. Lactophos.), Calcium Lactophosphate, N. F.—A mixture of calcium lactate and calcium acid lactate with calcium acid phosphate.

White, odorless powder. Soluble in water; almost insoluble in alcohol.

ACTION AND USES: Similar to but without advantage over the lactate. Not adapted to hypodermic use.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Elixir Calcii Lactophosphatis (Elix. Calc. Lactophos.), *Elixir of Calcium Lactophosphate*, N. F.—Calcium lactophosphate (3%) in compound spirit of orange, syrup, alcohol and distilled water. Absolute alcohol content about 19 per cent. Small and probably inefficient quantities of calcium.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Cinchonae Alkaloidorum, Ferri et Calcii Lactophosphatis (Elix. Cinchon., Ferr. et Calc. Lactophos.), *Elixir of*

Cinchona Alkaloids, Iron and Calcium Lactophosphate, N. F.—(Elixir Cinchonae, Ferri et Calcii Lactophosphatis, N. F. III, Elixir of Calisaya, Iron and Lactophosphate of Lime).—Syrup of calcium lactophosphate (50%), potassium citrate (3%) and elixir of cinchona alkaloids and iron. Absolute alcohol content about 12 per cent.

USES: An irrational mixture, supposed to present seven active ingredients.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Syrupus Calcii Lactophosphatis (Syr. Calc. Lactophos.), Syrup of Calcium Lactophosphate, U. S. P.—Made by dissolving precipitated calcium carbonate (2.5%) in lactic acid (6%), phosphoric acid (3.6%) and a syrup containing orange flower water.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Syrupus Calcii Lactophosphatis et Ferri (Syr. Calc. Lactophos. et Ferr.), Syrup of Calcium Lactophosphate and Iron, N. F.—Ferrous lactate and potassium citrate (each 0.85%) and syrup of calcium lactophosphate (about 99%).

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Calcii Phosphas Praecipitatus (Calc. Phos. Praec.), Precipitated Calcium Phosphate, N. F.— $\text{Ca}_3(\text{PO}_4)_2$.

A bulky, white, odorless, tasteless powder. Practically insoluble in water and insoluble in alcohol.

ACTION AND USES: Precipitated calcium phosphate has been given in the form of powders or mixtures for its action as a calcium salt, which is similar to but without advantage over the action of the carbonate. Because of its insolubility it is now seldom used.

USES: Similar to but without advantage over the carbonate.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Syrupus Calcii Hydrochlorophosphatis (Syr. Calc. Hydrochlorophos.), Syrup of Calcium Hydrochlorophosphate, N. F. (Syrupus Calcii Chlorhydrophosphatis, N. F. III).—Precipitated calcium phosphate (1.75%) flavored with tincture of lemon peel and dissolved in hydrochloric acid, water and syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Calcii Sulphidum Crudum (Calc. Sulphid. Crud.), Crude Calcium Sulphide, U. S. P. (Calx Sulphurata, U. S. P. VIII, Sulphurated Lime).— CaS (not less than 55%).

A pale gray or yellowish powder, with a faint hydrogen sulphide odor and a nauseous, alkaline taste. Very slightly soluble in cold water, insoluble in alcohol.

ACTION AND USES: Externally, depilatory similar to barium sulphide. Internally has been used to arrest and prevent suppuration especially in boils, carbuncles and pustular acne. There is no reliable evidence of its therapeutic usefulness.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.), in solution.

Liquor Calcis Sulphuratae (Liq. Calc. Sulphurat.), *Solution of Sulphurated Lime*, N. F. (Solution of Oxysulphuret of Calcium, Fleming's Solution, Vleminckx' Solution, Vleminckx' Lotion).—A solution of calcium polysulphides and calcium thiosulphate produced by boiling together lime, sulphur and water.

USES: Depilatory.

Calendula (Calend.), **Calendula**, N. F. (Marigold, Calendula Flowers).

ACTION AND USES: Locally, mild irritant, without material advantage over diluted alcohol. Internally, no definite indications for its use.

Fluidextractum Calendulae (Fldext. Calend.), *Fluidextract of Calendula*, N. F.—Calendula (100%). Absolute alcohol content about 80 per cent.

Tinctura Calendulae (Tr. Calend.), *Tincture of Calendula*, N. F.—Calendula (20%) in alcohol. Absolute alcohol content about 90 per cent.

Calumba (Calumb.), **Calumba**, U. S. P. (Columba, Columbo, Colombo).—A root.

ACTION AND USES: Used as a simple bitter and stomachic. It appears to have no advantage over other bitters, for instance, gentian.

DOSAGE: 2 Gm. or 30 grains (U. S. P.); usually given in the form of the tincture.

Fluidextractum Calumbae (Fldext. Calumb.), *Fluidextract of Calumba*, N. F.—Calumba (100%). Absolute alcohol content about 65 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Calumbae (Tr. Calumb.), *Tincture of Calumba*, U. S. P.—Calumba (20%) in alcohol and water. Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

***Calx (Calx), Calcium Oxide, U. S. P. (Lime, Quicklime).—CaO.**

Hard, white, odorless masses with a caustic taste, forming with water calcium hydroxide, which is slightly soluble in water (1:84), insoluble in alcohol.

ACTION AND USES: As liquor calcis (an aqueous solution of calcium hydroxide), used as an antacid and mild alkali. As milk of lime (a mixture of calcium hydroxide and water), frequently used as a disinfectant.

***Linimentum Calcis (Lin. Calc.), Lime Liniment, U. S. P. (Carron Oil).—Lime water (50%) and linseed oil.**

USES: Topical application, used especially for burns.

***Liquor Calcis (Liq. Calc.), Solution of Calcium Hydroxide, U. S. P. (Lime Water).—Ca(OH)₂ (not less than 0.14%).**

DOSAGE: 15 Cc. or 4 fluidrachms.

***Calx Chlorinata (Calx Chlorin.), Chlorinated Lime, U. S. P. ("Chloride of Lime").—A compound of calcium chloride and hypochlorite containing not less than 30 per cent. of available chlorine.**

A white or grayish-white powder having a chlorine-like odor and a repulsive saline taste. Partially soluble in water or alcohol. Decomposed by acids with liberation of chlorine.

ACTION AND USES: Disinfectant, especially for excreta. Should be used liberally and left in contact for at least one hour. Widely used for treating contaminated drinking water.

Cambogia (Cambog.), Gamboge, U. S. P. (Pipe Gamboge).—A gum resin.

Odorless, grayish-orange-brown pieces or bright yellow powder, with a very acrid taste. Not less than 65 per cent. soluble in alcohol.

ACTION AND USES: A powerful irritant hydragogue cathartic; not often employed alone. Usually administered in combination with other drugs.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.), in pills.

***Camphora (Camph.), Camphor, U. S. P.—A volatile solid obtained from the camphor tree.**

Tough, white, translucent masses with a pungent odor and slightly soluble in water and freely soluble in alcohol, chloroform and fixed or volatile oils.

ACTION AND USES: Used hypodermically to stimulate circulation and respiration in collapse. Opinions

divided as to its efficiency. Locally, applied as a mild irritant and antiseptic.

DOSAGE: By mouth 0.2 Gm. or 3 grains (U. S. P.). Hypodermic, 0.1 Gm. or 1½ grains (U. S. P.), as 10 per cent. solution in oil.

**Aqua Camphorae* (*Aq. Camph.*), *Camphor Water*, U. S. P.—A saturated solution of camphor in water.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Ceratum Camphorae (*Cerat. Camph.*), *Camphor Cerate*, N. F.—Camphor (2%), with cottonseed oil, white wax, white petrolatum and benzoinated lard.

Emplastrum Fuscum Camphoratum (*Emp. Fusc. Camph.*), *Camphorated Brown Plaster*, N. F. (Camphorated Motherplaster).—Camphor (1%), red oxide of lead, olive oil and yellow wax.

**Linimentum Camphorae* (*Lin. Camph.*), *Camphor Liniment*, U. S. P. (Camphorated Oil).—Camphor (20%) in cottonseed oil.

USES: Mild counterirritant. Note: This preparation, as found in drug stores, should never be used hypodermically.

Mistura Camphora Acida.—See under *Opium*.

Mistura Camphorae Aromatica.—See under *Oleum Lavandulae*.

**Spiritus Camphorae* (*Sp. Camph.*), *Spirit of Camphor*, U. S. P.—Camphor (10%), in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Unguentum Camphorae (*Ung. Camph.*), *Camphor Ointment*, N. F.—Camphor (22%) in white wax and lard.

Unguentum Fuscum (*Ung. Fusc.*), *Brown Ointment*, N. F. (*Unguentum Matris*, *Mother's Salve*).—Camphorated brown plaster (50%) olive oil and prepared suet.

Camphora Monobromata (*Camph. Monobrom.*), **Monobromated Camphor**, U. S. P.

Colorless crystals or white powder, with a mild camphoraceous odor and taste. Freely soluble in alcohol (1:6.5); almost insoluble in water.

ACTION AND USES: Formerly recommended as a sedative and hypnotic but now seldom used. Value doubtful.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Canella (*Canell.*), **Canella**, N. F.—A bark.

ACTION AND USES: Obsolete aromatic. Without advantage over gentian.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Cannabis (Cannab.), Cannabis, U. S. P. (*Cannabis Indica*, U. S. P. VIII).—Flowering tops of *Cannabis sativa* (or var. *indica*).

ACTION AND USES: A narcotic poison, producing a mild delirium. Used in sedative mixtures, but of doubtful value. Also employed to color corn remedies.

Extractum Cannabis (Ext. Cannab.), Extract of Cannabis, U. S. P.—Assayed biologically by producing incoordination in dogs.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.). Physicians who prescribe extract of cannabis should specify the biologically assayed preparation.

Fluidextractum Cannabis (Fldext. Cannab.), Fluidextract of Cannabis, U. S. P.—Cannabis (100%). Assayed biologically by producing incoordination in dogs. Absolute alcohol content about 80 per cent.

DOSAGE: 0.1 Cc. or $1\frac{1}{2}$ minims (U. S. P.). Physicians who prescribe fluidextract of cannabis should specify the biologically assayed preparation.

Tinctura Cannabis (Tr. Cannab.), Tincture of Cannabis, U. S. P. (*Tinctura Cannabis Indicae*, U. S. P. VIII).—Cannabis (10%) in alcohol. Assayed biologically by producing incoordination in dogs. Absolute alcohol content about 90 per cent.

DOSAGE: 0.75 Cc. or 12 minims (U. S. P.). Those physicians who prescribe tincture of cannabis should specify the biologically assayed preparation.

***Cantharis (Canthar.), Cantharides, U. S. P.** (Spanish Flies, Russian Flies).—A beetle.

ACTION AND USES: Externally, rubefacient and vesicant. Internally, genito-urinary irritant, often producing serious nephritis. Its internal use is probably never justified.

***Ceratum Cantharidis (Cerat. Canthar.), Cantharides Cerate, U. S. P.** (Blistering Cerate).—Cantharides (35%) with turpentine, glacial acetic acid, yellow wax, rosin and benzoinated lard.

Collodium Cantharidatum (Collod. Canth.), Cantharidal Collodion, U. S. P. (Blistering Collodion, Vesicating Collodion).—The glacial acetic acid acetone extract from cantharides (60%) in flexible collodion.

Emplastrum Cantharidis (Emp. Canthar.), Cantharides Plaster, U. S. P.—Cantharides cerate spread on rosin

plaster. Each square centimeter of spread plaster contains 0.1 Gm. of cantharides cerate.

USES: Convenient form for applying cantharides as a vesicant. Requires from six to eight hours to produce a blister. May produce poisoning by absorption through the skin.

**Tinctura Cantharidis* (*Tr. Canthar.*), *Tincture of Cantharides*, U. S. P. (*Cantharidis Tinctura* P. I.).—Cantharides (10%) in alcohol. Absolute alcohol content about 90 per cent.

**Capsicum* (*Capsic.*), *Capsicum*, U. S. P. (Cayenne Pepper, African Chillies).—Fruit.

ACTION AND USES: Carminative and rubefacient; also stomachic, especially in alcoholic gastritis (as the tincture).

Emplastrum Capsici (*Emp. Capsic.*), *Capsicum Plaster*, U. S. P.—Oleoresin of capsicum spread on rubber plaster. Each 15 square cm. of spread plaster contains 0.25 Gm. of oleoresin of capsicum.

Oleoresina Capsici (*Oleores. Capsic.*), *Oleoresin of Capsicum*, U. S. P.—Highly irritant.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

**Tinctura Capsici* (*Tr. Capsic.*), *Tincture of Capsicum*, U. S. P.—Capsicum (10%) in alcohol and water. Absolute alcohol content about 88 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

Tinctura Capsici et Myrrhae (*Tr. Capsic. et Myrrh.*), *Tincture of Capsicum and Myrrh*, N. F. (Hot Drops, Thomsonian Number Six).—Capsicum (3%) and myrrh (12%) in alcohol and water. Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Caramel (*Caram.*), *Caramel*, N. F. (*Saccharum Ustum*, Burnt Sugar Coloring).

A dark brown, syrupy, somewhat bitter liquid. Soluble in water and in diluted alcohol; insoluble in 80 per cent. alcohol.

ACTION AND USES: Used in pharmacy as a brown coloring agent.

Tinctura Caramellis, *Tincture of Caramel*, N. F.—Caramel in alcohol and water. Absolute alcohol content about 20 per cent.

**Carbo Ligni* (*Carbo Lig.*), *Wood Charcoal*, U. S. P. (Coal).—Powdered wood charcoal.

ACTION AND USES: Absorbent of gases and dissolved substances. Used internally against digestive disorders and vegetable poisons but of doubtful value. Locally employed as a deodorant, for fetid ulcers, usually in the form of a poultice.

DOSAGE: 1 Gm. or 15 grains, in powder or suspension.

Trochisci Carbonis Ligni (*Troch. Carb. Lig.*), *Troches of Charcoal*, N. F.—Charcoal with tragacanth, sugar and vanillin.

DOSAGE: 1 troche (N. F.).

***Cardamomi Semen** (*Cardam. Sem.*), *Cardamom Seed*, U. S. P. (*Cardamomum*, U. S. P. VIII).

ACTION AND USES: Used as flavor and carminative. Frequently added to purgatives, as in the compound extract of colocynth, with the object of diminishing griping.

**Tinctura Cardamomi* (*Tr. Cardam.*), *Tincture of Cardamom*, U. S. P.—Cardamom seed (15%) in diluted alcohol. Absolute alcohol content about 46 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Cardamomi Composita (*Tr. Cardamom. Co.*), *Compound Tincture of Cardamom*, U. S. P.—Cardamom seed, Saigon cinnamon, caraway and cochineal in alcohol, glycerin and water. Absolute alcohol content about 46 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Carminum (*Carmin.*), *Carmine*, N. F.—From cochineal.

ACTION AND USES: Used in pharmacy as a red coloring agent.

Liquor Carmini (*Liq. Carmin.*), *Solution of Carmine*, N. F.—Carmine in ammonia water, glycerin and water.

Caro, Beef.

Extractum Carnis (*Ext. Carnis*), *Extract of Beef*, N. F. (*Beef Extract*).—Residue from evaporation of beef broth.

ACTION AND USES: Stimulant and flavor; not a nutrient tonic, as was believed at one time. The "wines" are without real value and can easily be misused as beverages.

Vinum Carnis (*Vin. Carn.*), *Wine of Beef*, N. F. (*Beef and Wine*).—Extract of beef (3%) in water, syrup, alcohol, compound spirit of orange and sherry wine. Absolute alcohol content about 20 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Vinum Carnis et Ferri (*Vin. Carn. et Ferr.*), *Wine of Beef and Iron*, N. F. (Beef, Wine and Iron).—Extract of beef (3%), iron and ammonium citrate, water, syrup, alcohol, compound spirit of orange and sherry wine.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Carum (**Carum**), Caraway, U. S. P. (Carawayseed).

ACTION AND USES: Aromatic carminative. Similar to anise and fennel.

***Caryophyllus** (**Caryoph.**), Clove, U. S. P. (Cloves).

ACTION AND USES: Aromatic carminative and counter-irritant.

Pulvis Aromaticus Rubefaciens (*Pulv. Arom. Rubefac.*), *Rubefacient Spice Powder*, N. F. (Replacing *Emplastrum Aromaticum*, N. F. III).—Clove, Saigon cinnamon, ginger and capsicum.

***Cascara Sagrada** (**Casc. Sagr.**), *Cascara Sagrada*, U. S. P. (*Rhamnus Purshiana*, U. S. P. VIII).—A bark.

ACTION AND USES: Laxative, acting mainly on the colon. Widely used for habitual constipation. Has little tendency to produce secondary constipation.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Cascarae Sagradae (*Elix. Cascar. Sagr.*), *Elixir of Cascara Sagrada*, N. F. (*Elixir Rhamni Purshianae*, N. F. III).—Aromatic fluidextract of cascara sagrada (50%) and aromatic elixir. Absolute alcohol content about 22 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Cascarae Sagradae Compositum (*Elix. Cascar. Sagr. Co.*), *Compound Elixir of Cascara Sagrada*, N. F. (*Elixir Rhamni Purshianae Co.*, N. F. III, Laxative Elixir).—Aromatic fluidextract of cascara sagrada (12.5%), fluidextract of senna (7.5%), fluidextract of juglans (6.5%) and aromatic elixir. Absolute alcohol content about 24 per cent.

USES: Needlessly complicated laxative.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Extractum Cascarae Sagradae** (**Ext. Cascar. Sagr.**), *Extract of Cascara Sagrada*, U. S. P. (*Extractum Rhamni Purshianae*, U. S. P. VIII, Powdered Extract of *Cascara Sagrada*).—One Gm. extract represents 3 Gm. cascara sagrada.

DOSAGE: 0.25 Gm. or 4 grains

**Fluidextractum Cascarae Sagradae* (*Fldext. Cascar. Sagr.*), *Fluidextract of Cascara Sagrada*, U. S. P. (Fluidextractum Rhamni Purshianae, U. S. P. VIII).—Cascara sagrada (100%). Absolute alcohol content about 20 per cent. Bitter, unpleasant taste.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

**Fluidextractum Cascarae Sagradae Aromaticum* (*Fldext. Cascar. Sagr. Arom.*), *Aromatic Fluidextract of Cascara Sagrada*, U. S. P. (Fluidextractum Rhamni Purshianae Aromaticum, U. S. P. VIII).—Sweetened and flavored fluidextract. Absolute alcohol content about 20 per cent. Not so bitter, but less effective than the plain fluidextract.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Fluidglyceratum Cascarae Sagradae (*Fldglycer. Cascar. Sagr.*), *Fluidglycerate of Cascara Sagrada*, N. F.—Cascara sagrada (100%) in water and glycerin.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Fluidglyceratum Cascarae Sagradae Aromaticum (*Fldglycer. Cascar. Sagr. Arom.*), *Aromatic Fluidglycerate of Cascara Sagrada*, N. F.—Cascara sagrada (75%) and fluidglycerate of glycyrrhiza (25%) with oils of fennel, clove and cinnamon, in water and glycerin.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Cascarilla (*Cascarill.*), **Cascarilla**, N. F. (Sweetwood Bark, Sweet Bark).

ACTION AND USES: Aromatic bitter, without advantage over gentian.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Cassia Fistula (*Cass. Fist.*), **Cassia Fistula**, N. F. (Purging Cassia).—Dried fruit.

ACTION AND USES: Obsolete laxative, without advantage over senna.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Castanea (*Castan.*), **Castanea**, N. F. (Chestnut Leaves).

ACTION AND USES: Astringent; no definite indications for its use.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Castaneae (*Fldext. Castan.*), *Fluidextract of Chestnut Leaves*, N. F.—Chestnut leaves (100%). Absolute alcohol content about 15 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

CATAPLASMATA, POULTICES.—Poultices are soft solid preparations used for the purpose of applying heat and moisture to localized areas of the body. Poultices may also be made the means of applying counterirritant drugs.

Cataria (Catar.), Catnep, N. F. (Catmint).—Leaves and flowering tops.

ACTION AND USES: Aromatic carminative, without advantage over peppermint.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Catariae (Fldext. Catar.), Fluidextract of Catnep, N. F.—Catnep (100%). Absolute alcohol content about 35 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Caulophyllum (Caulophyl.), Caulophyllum, N. F. (Blue Cohosh, Papoose Root, Squaw Root).—Rhizome and roots.

ACTION AND USES: Said to be diuretic and emmenagogue; has not been adequately studied.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Fluidextractum Caulophylli (Fldext. Caulophyl.), Fluidextract of Caulophyllum, N. F.—Caulophyllum (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

Centaurium (Centaur.), Centaury, N. F.—Flowering plant.

ACTION AND USES: Bitter stomachic; without advantage over gentian. Also used in "female remedies," but probably useless.

DOSAGE: 2 Gm. or 30 grains (N. F.).

***Cera Alba (Cer. Alb.), White Wax, U. S. P.**—Yellow wax bleached.

ACTION AND USES: Used in the preparation of emollient ointments, and to raise the melting point of mixtures of oils and fats.

Ceratum (Cerat.), Cerate, U. S. P. (Simple Cerate).—White wax (30%) and benzoinated lard.

***Cera Flava (Cer. Flav.), Yellow Wax, U. S. P. (Beeswax).**

ACTION AND USES: Used in the making of plasters and of ointments in which the color is not objectionable.

CERATAE, CERATES.—Cerates

preparations containing
becoming soft at or

like

Cerevisiae Fermentum Compressum (Cerev. Ferm. Compr.), Compressed Yeast, N. F.—The commercial compressed yeast.

ACTION AND USES: For use in making fermented milk (Lac Fermentatum), which see under **Lac Vaccinum**.

Cerii Oxalas (Cerii Oxal.), Cerium Oxalate, U. S. P.—A mixture of the oxalates of cerium, didymium, lanthanum, and other associated elements.

White or pinkish, odorless, tasteless powder. Insoluble in water, alcohol and ether.

ACTION AND USES: Formerly used for the treatment of chronic vomiting but probably without value.

DOSAGE: 0.2 Gm. or 3 grains (U. S. P.), in powders, cachets or capsules.

Cetaceum (Cetac.), Spermaceti, U. S. P.—A solid fat from the sperm whale.

White, nearly odorless, unctuous masses, with a bland, mild taste. Insoluble in water and nearly insoluble in cold alcohol.

ACTION AND USES: Used as a component of cerates and cold creams.

CHARTAE, PAPERS.—Papers are pieces of unsized paper saturated with or pieces of hard-surfaced papers coated with a medicinal substance.

Chimaphila (Chimaph.), Chimaphila, N. F. (Pipsissewa).—Leaves.

ACTION AND USES: Obsolete; no definite indications for its use. Has been described as "tonic, astringent, diuretic."

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Chimaphilae (Fldext. Chimaphil.), Fluid-extract of Chimaphila, N. F.—Chimaphila (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Chionanthus (Chionanth.), Chionanthus, N. F. (Fringe Tree Bark).—Bark of the root.

ACTION AND USES: Obsolete; no definite indications for its use. Formerly used by eclectics and homeopaths in hepatic disorders and syphilis.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Chionanthi (Fldext. Chionanth.), Fluid-extract of Chionanthus, N. F.—Chionanthus (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Chirata (Chirat.), Chirata, N. F. (Chiretta).—Plant.

ACTION AND USES: Bitter stomachic, similar to and without advantage over gentian.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Chiratae (Fldext. Chirat.), Fluidextract of Chirata, N. F.—Chirata (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Chloral Camphoratum (Chloral Camph.), Camphorated Chloral, N. F.—Equal parts of hydrated chloral and camphor.

An oily liquid with a camphoraceous odor.

ACTION AND USES: Local rubefacient and analgesic.

***Chloralum Hydratum (Chloral. Hydrat.), Hydrated Chloral, U. S. P. ("Chloral," Chloral Hydrate).— $\text{CCl}_3\text{CHO} + \text{H}_2\text{O}$.**

Colorless, transparent crystals, with an aromatic, penetrating odor and a bitterish, caustic taste. Very soluble in water (1:0.25) and in alcohol (1:1.3).

ACTION AND USES: A reliable hypnotic in moderate doses, inducing a condition closely resembling that of normal sleep. Used for the relief of insomnia and nervousness, and for the control of convulsions and spasms. Should be used with special caution in cases of weakened heart or blood vessels. Continued use is likely to produce a habit.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.), in solution. Decomposed by alkalies; should not be dispensed in liquids containing both bromides and alcohol.

Mistura Chloralis et Potassii Bromidi Composita (Mist. Chloral. et Pot. Brom. Co.), Compound Mixture of Chloral and Potassium Bromide, N. F. (Chloral and Bromide Compound).—Hydrated chloral (20%), potassium bromide (20%), extract of hyoscyamus (0.2%) in water. The formula calls for cannabis but this is filtered out.

USES: Similar in composition to the widely advertised nostrum Bromidia and equally irrational and objectionable.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Chloroformum (Chlorof.), Chloroform, U. S. P.— CHCl_3 .**

A clear, colorless liquid, with an ethereal odor and a burning, sweet taste. Slightly soluble (1:210) in water and miscible with alcohol, ether, petroleum benzin or fixed or volatile oils. Decomposed by alkalies.

ACTION AND USES: General anesthetic, administered by inhalation. Locally, a penetrating and fairly powerful irritant. Frequently used in the form of liniments. Taken by mouth in small doses, carminative, anodyne and antiseptic. Larger doses are anthelmintic but dangerous. A direct cardiac depressant.

DOSAGE: 0.3 Cc. or 5 minims (U. S. P.), diluted.

**Aqua Chloroformi (Aq. Chlorof.), Chloroform Water, U. S. P.*—A saturated solution of chloroform in water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

**Linimentum Chloroformi (Lin. Chlorof.), Chloroform Liniment, U. S. P.*—A mixture of chloroform (30%) and soap liniment. Fairly active counterirritant.

Mistura Chloroformi et Morphinae Composita (Mist. Chlorof. et Morph. Co.), Compound Mixture of Chloroform and Morphine, N. F. (Mistura Chloroformi et Cannabis Indicae Composita, N. F. III, Chloroform Anodyne).—Morphine sulphate (0.25%), chloroform (12.5%), ether (3.25%), tincture of cannabis (18.5%), tincture of capsicum, oil of peppermint, glycerin, water and alcohol. Absolute alcohol content about 60 per cent.

USES: An evident imitation of the nostrum sold as chloranodyne, and equally irrational and objectionable.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Petroxolinum Chloroformi Camphoratum (Petrox. Chlorof. Camph.), Camphorated Chloroform Petroxolin, N. F. (Camphor and Chloroform Petrox).—Chloroform (25%), camphor (25%) and liquid petroxolin.

**Spiritus Chloroformi (Sp. Chlorof.), Spirit of Chloroform, U. S. P.*—Chloroform (6%) in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Chlorum, Chlorine.—Cl.

Liquor Chlorig Compositus (Liq. Chlor. Co.), Compound Solution of Chlorine, N. F. (Chlorine Water).—A solution containing chlorine, oxides of chlorine and hydrochloric acid made by treating potassium chlorate with hydrochloric acid and dissolving the gases produced in water.

USES: Antiseptic.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Chondrus (Chondrus), Chondrus, U. S. P. (Irish Moss, Carrageen).—Plant.

ACTION AND USES: Demulcent and lubricant, possessing no advantages over other gums, acacia, for instance.

Gelatinum Chondri (*Gelat. Chondr.*), *Chondrus Gelatin*, N. F. (Irish Moss Gelatin).—Obtained by extraction with hot water and evaporation to semifluid consistency.

Mucilago Chondri (*Mucil. Chondr.*), *Mucilage of Chondrus*, N. F. (Mucilage of Irish Moss).—*Chondrus* (3%) in water.

***Chromii Trioxidum** (*Chrom. Triox.*), **Chromium Trioxide**, U. S. P. ("Chromic Acid," Chromic Anhydride).— CrO_3 .

Dark purplish-red, odorless, deliquescent crystals. Very soluble in water (1:0.6). Decomposes, sometimes with dangerous violence, when brought into contact with organic solvents like alcohol and glycerin. Incompatible with most organic substances.

ACTION AND USES: A powerful oxidizing agent and caustic. Its action is somewhat difficult to control.

***Chrysarobinum** (*Chrysarob.*), **Chrysarobin**, U. S. P.—Neutral principles from Goa powder.

Brownish to orange-yellow, tasteless, odorless powder, irritating to the mucous membrane. Slightly soluble in alcohol (1:385), soluble in chloroform (1:12.5) and very slightly soluble in water.

ACTION AND USES: Antiseptic and antiparasitic; a powerful irritant to the skin, used chiefly in the treatment of psoriasis, trichophytosis and alopecia areata.

**Unguentum Chrysarobini* (*Ung. Chrysarobin.*), *Chrysarobin Ointment*, U. S. P.—Chrysarobin (6%) in benzoinated lard.

Cimicifuga (*Cimicif.*), **Cimicifuga**, U. S. P. (Black Snake-root, Black Cohosh, *Macrotys*).—A rhizome.

ACTION AND USES: One of the class of "domestic medicines" that were tried for a great variety of conditions (used as "stomachic, antispasmodic, aphrodisiac, diaphoretic, diuretic and expectorant"), but which have not been found to possess definite value.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Extractum Cimicifugae (*Ext. Cimicif.*), *Extract of Cimicifuga*, U. S. P. (Powdered Extract of *Cimicifuga*).—One Gm. extract represents 4 Gm. *cimicifuga*.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Fluidextractum Cimicifugae (*Fldest. Cimicif.*), *Fluidextract of Cimicifuga*, U. S. P. (Fluidextract of ^P *Cimicifuga*, U. S. P. (Fluidextract of Black Snakeroot).—

Absolute alcohol content about 8

DOSAGE: 1 Cc. or 15 minims (1

Syrupus Cimicifugae Compositus (Syr. Cimicif. Co.), Compound Syrup of Cimicifuga, N. F. (Syrupus Actaeae Compositus, N. F. III, Compound Syrup of Actaea). Fluidextract of cimicifuga (4%), fluidextract of senega (2%), fluidextract of ipecac (1%), fluidextract of glycyrrhiza, wild cherry, sugar and water. Absolute alcohol content about 5 per cent.

USES: Complex expectorant.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Tinctura Cimicifugae (Tr. Cimicif.), Tincture of Cimicifuga, N. F.—Cimicifuga (20%) in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Cinchona* (Cinch.), Cinchona, U. S. P. (Yellow Cinchona, Calisaya Bark, Yellow Peruvian Bark).—Yields not less than 5 per cent. of alkaloids.

ACTION AND USES: Astringent bitter tonic.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Cinchonae Alkaloidorum and *Elixir Cinchonae Alkaloidorum et Hypophosphitum*.—See under Quinine Sulphas.

Elixir Cinchonae Alkaloidorum et Ferri.—See under Ferric Phosphas.

Elixir Cinchonae Alkaloidorum, Ferri, Bismuthi et Strychninae and *Elixir Cinchonae Alkaloidorum, Ferri et Strychninae*.—See under Strychninae Sulphas.

Elixir Cinchonae Alkaloidorum, Ferri et Bismuthi.—See under Bismuthi et Sodii Tartras.

Elixir Cinchonae Alkaloidorum, Ferri et Calcii Lactophosphatis.—See under Calcii Lactophosphas.

Elixir Cinchonae Alkaloidorum, Ferri et Pepsini.—See under Pepsinum.

Extractum Cinchonae (Ext. Cinchon.), Extract of Cinchona, N. F.—A hydroalcoholic extract diluted with glucose, yielding about 24 per cent. of cinchona alkaloids.

DOSAGE: 0.25 Gm. or 4 grains (N. F.).

Fluidextractum Cinchonae (Fldext. Cinchon.), Fluidextract of Cinchona, U. S. P.—Cinchona (100%) yielding about 4.5 per cent. of alkaloids. Absolute alcohol content about 65 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Fluidextractum Cinchonae Aquosum (Fldext. Cinchon. Aq.), *Aqueous Fluidextract of Cinchona*, N. F.—Cinchona (100%), yielding about 5 per cent. of alkaloids. Absolute alcohol content about 12 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Infusum Cinchonae (Inf. Cinchon.), *Infusion of Cinchona*, N. F.—Cinchona (6%), aromatic sulphuric acid (1%) and water.

DOSAGE: 50 Cc. or 12 fluidrachms (N. F.).

**Tinctura Cinchonae* (Tr. Cinch.), *Tincture of Cinchona*, U. S. P.—Cinchona (20%), yielding about 0.9 per cent. of alkaloids; in alcohol, glycerin and water. Absolute alcohol content about 63 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Cinchona Rubra (Cinch. Rub.), **Red Cinchona**, U. S. P. (Red Peruvian Bark).—Yields not less than 5 per cent. of alkaloids.

ACTION AND USES: Identical with those of cinchona. Used as ingredient of the compound tincture of cinchona.

**Tinctura Cinchonae Composita* (Tr. Cinch. Co.), *Compound Tincture of Cinchona*, U. S. P.—Red cinchona (10%), yielding about 0.45 per cent. of cinchona alkaloids, serpentaria (2%) and bitter orange peel, in glycerin, alcohol and water. Absolute alcohol content about 64 per cent.

USES: Aromatic astringent bitter, of high alcoholic content; superfluous.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Cinchonidinae Sulphas (Cinchonid. Sulph.), **Cinchonidine Sulphate**, U. S. P.—The sulphate of the alkaloid cinchonidine, obtained from cinchona bark.

White, glistening, odorless needles with a very bitter taste. Soluble in water (1:65) and in alcohol (1:90).

ACTION AND USES: Formerly used as a cheap substitute for quinine sulphate, over which it possesses no advantage, being less efficient and more convulsant.

DOSAGE: 0.15 Gm. or 2½ grains (U. S. P.).

Cinchoninae Sulphas (Cinchonin. Sulph.), **Cinchonine Sulphate**, U. S. P.—The sulphate of the alkaloid cinchonine, obtained from cinchona bark.

Crystals with a very bitter taste. Soluble in water (1:12.5).

ACTION AND USES: Similar to those of cinchonidine sulphate.

DOSAGE: 0.15 Gm. or 2½ grains (U. S. P.).

***Cinnamomum Saigonicum** (Cinnam. Saigon.), Saigon Cinnamon, U. S. P.—A bark.

ACTION AND USES: Carminative, antiseptic, somewhat astringent, occasionally administered in diarrhea.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), in powder.

Fluidextractum Aromaticum (Fldext. Aromat.), *Aromatic Fluidextract*, U. S. P.—Aromatic powder (100%), extracted with alcohol. Absolute alcohol content about 80 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Pulvis Aromaticus (Pulv. Arom.), *Aromatic Powder*, U. S. P.—Saigon cinnamon, Jamaica ginger, cardamom seed and nutmeg.

USES: Carminative.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Pulvis Cretae Aromaticus (Pulv. Cret. Arom.), *Aromatic Powder of Chalk*, N. F.—Saigon cinnamon, nutmeg, clove, cardamom seed, prepared chalk and sugar.

USES: Antidiarrheic.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Syrupus Cinnamomi (Syr. Cinnam.), *Syrup of Cinnamon*, N. F.—Saigon cinnamon, alcohol and syrup.

USES: Flavoring vehicle.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Tinctura Aromatica (Tr. Arom.), *Aromatic Tincture*, N. F.—Saigon cinnamon, Jamaica ginger, galangal, clove, cardamom seed, extracted with a mixture of alcohol and water. Absolute alcohol content about 63 per cent.

USES: Aromatic carminative, complex and without special advantage over other carminatives, as, for instance, tincture of cardamom.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Cinnamomi (Tr. Cinnam.), *Tincture of Cinnamon*, U. S. P.—Saigon cinnamon in alcohol, glycerin and water. Absolute alcohol content about 63 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Cinnamomum Zeylanicum (Cinnam. Zeylan.), Ceylon Cinnamon, U. S. P.—A bark.

ACTION AND USES: Same as those of Saigon cinnamon.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

- ***Cocaina (Cocain.), Cocaine, U. S. P.**—An alkaloid obtained from coca leaves.

Large colorless prisms or a white powder; odorless. Slightly soluble in water (1: 600) and freely soluble in alcohol (1: 6.5) and in chloroform (1: 0.7).

ACTION AND USES: A local anesthetic, paralyzing the peripheral sensory nerves and also contracting the blood vessels. Acute systemic poisoning is often caused by its local use on mucous membranes. Stimulant to the central nervous system, but such use is inadvisable because of danger of habit-formation.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.).

Oleatum Cocainae (Oleat. Cocain.), Oleate of Cocaine, N. F.
—Cocaine (5%) in alcohol, oleic acid and olive oil.

- ***Cocainae Hydrochloridum (Cocain. Hydrochl.), Cocaine Hydrochloride, U. S. P. (Cocaine Chloride, Cocainum Hydrochloricum, P. I.).**

Colorless crystals, flaky leaflets or a white powder; odorless. Freely soluble in water (1: 0.4) and in alcohol (1: 3.2), and soluble in chloroform (1: 12.5). Incompatible with borax, mercuric chloride, alkalis and alkali carbonates, phenol, tannic acid, mercuric oxide, and soluble silver salts.

ACTION AND USES: Local anesthetic.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.). For local anesthesia in 0.2 to 1.0 per cent. solution when used hypodermatically, and in 1 to 4 per cent. solution when applied to mucous membranes.

- Cocculus Indicus (Coccul. Ind.), Cocculus Indicus, N. F. (Fish Berry, Indian Berry).**—A fruit.

ACTION AND USES: Source of picrotoxin. An alcoholic extract has been used for destroying pediculi.

Tinctura Cocculi Indici (Tr. Coccul. Ind.), Tincture of Cocculus Indicus, N. F.—Cocculus indicus (10%) in diluted alcohol. Absolute alcohol content about 45 per cent.

- Coccus (Coccus), Cochineal, U. S. P.**—A dried insect.

ACTION AND USES: Coloring agent.

Liquor Cocci (Liq. Cocci), Cochineal Color, N. F.—A glycerin-alcohol-water solution of cochineal.

- Cocillana (Cocillan.), Cocillana, N. F.**—A bark.

ACTION AND USES: Nauseant expectorant, similar to but without advantage over ipecac. The proprietary syrup contains heroin.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Cocillanae (*Fldext. Cocillan.*), *Fluidextract of Cocillana*, N. F.—Cocillana (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

- ***Codeina** (Codein.), **Codeine**, U. S. P. (Methylmorphine).—An alkaloid obtained from opium, or prepared from morphine by methylation.

Colorless, translucent prisms or a white powder; odorless. Slightly soluble in water (1:120) and freely soluble in alcohol (1:2) and in chloroform (1:0.5).

ACTION AND USES: Analgesic, hypnotic and sedative. It is perhaps the best modification of morphine for use in coughs. As compared with morphine it is less constipating and less apt to induce a drug habit. The phosphate and sulphate are preferred for hypodermic use.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

Elixir Terpini Hydratis et Codeinae (*Elix. Terpin. Hyd. et Codein.*), *Elixir of Terpin Hydrate and Codeine*, N. F.—Codeine (0.2%) and elixir of terpin hydrate. Absolute alcohol content about 42 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.). This dose contains about 0.008 Gm. or $\frac{1}{8}$ grain of codeine, and about 0.07 Gm. or $1\frac{1}{6}$ grains of terpin hydrate.

- ***Codeinae Phosphas** (Codein. Phos.), **Codeine Phosphate**, U. S. P.

White, odorless, efflorescent powder. Freely soluble in water (1:2.3) and slightly soluble in alcohol (1:325).

ACTION AND USES: Same as those of codeine; preferred for hypodermic use.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

- ***Codeinae Sulphas** (Codein. Sulph.), **Codeine Sulphate**, U. S. P.

White, odorless, efflorescent crystals or powder. Soluble in water (1:30) and very slightly soluble in alcohol (1:1280).

ACTION AND USES: Same as those of codeine. Preferred for hypodermic injections.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

Syrupus Codeinae (*Syr. Codein.*), *Syrup of Codeine*, N. F.—Codeine sulphate (0.2%) and syrup.

USES: Superfluous.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Coffea Tosta (Coff. Tost.), Coffee, N. F. (Roasted Coffee).

ACTION AND USES: Beverage and flavor.

Fluidextractum Coffeae (Fldext. Coff.), *Fluidextract of Coffee*, N. F. (Fluidextractum Coffeae Tostae, N. F. III).—Coffee (100%). Absolute alcohol content about 20 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Colchici Cormus (Colch. Corm.), *Colchicum Corm*, U. S. P. (Colchicum Root).—Yields not less than 0.35 per cent. of colchicine.

ACTION AND USES: Said to be antineuralgic and analgesic; frequently used in the treatment of acute gouty attacks and to some extent to ward off such attacks. Full doses produce watery stools and abdominal pains.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), in pills; or preferably as the tincture.

Extractum Colchici Cormi (Ext. Colch. Corm.), *Extract of Colchicum Corm*, U. S. P. (Powdered Extract of Colchicum Corm).—One Gm. extract represents 4 Gm. colchicum corm and yields about 1.4 per cent. of colchicine.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Fluidextractum Colchici Cormi (Fldext. Colch. Corm.), *Fluidextract of Colchicum Corm*, N. F.—Colchicum corm (100%), yielding about 0.35 per cent. of colchicine. Absolute alcohol content about 55 per cent.

DOSAGE: 0.2 Cc. or 3 minims (N. F.).

Vinum Colchici Cormi (Vin. Colchic. Corm.), *Wine of Colchicum Corm*, N. F.—Colchicum corm (40%) yielding about 0.14 per cent. of colchicine, in alcohol and sherry wine. Absolute alcohol content about 30 per cent.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

***Colchici Semen** (Colch. Sem.), *Colchicum Seed*, U. S. P. (Colchici Semen P. I.).—Yields not less than 0.45 per cent. of colchicine.

ACTION AND USES: Same as those of colchicum corm.

DOSAGE: 0.2 Gm. or 3 grains (U. S. P.).

Fluidextractum Colchici Seminis (Fldext. Colch. Sem.), *Fluidextract of Colchicum Seed*, U. S. P.—Colchicum seed (100%), yielding about 0.4 per cent. of colchicine. Absolute alcohol content about 55 per cent.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

**Tinctura Colchici Seminis* (*Tr. Colch. Sem.*), *Tincture of Colchicum Seed*, U. S. P. (*Colchici Tinctura* P. I.).—Colchicum seed (10%), yielding about 0.04 per cent. of colchicine; in alcohol and water. Absolute alcohol content about 56 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Vinum Colchici Seminis (*Vin. Colch. Sem.*), *Wine of Colchicum Seed*, N. F.—Fluidextract of colchicum seed (10%), yielding about 0.04 per cent. of colchicine; in alcohol and sherry wine. Absolute alcohol content about 35 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Colchicina (*Colchicina*), **Colchicine**, U. S. P.—An alkaloid obtained from colchicum.

Pale yellow, odorless scales or powder. Freely soluble in alcohol, soluble in water (1:22) and slightly soluble in ether (1:220).

ACTION AND USES: Same as those of colchicum corm.

DOSAGE: 0.5 mg. or $\frac{1}{120}$ grain.

COLLODIA, COLLODIONS.—Collodions are solutions of pyroxylin or soluble gun cotton with medicinal agents, for external application. For the official collodions see under the names of the principal ingredients. In particular, for Collodium, see under **Pyroxylin**; for Collodium Flexible, under **Oleum Ricini**; and for Collodium Stypticum, under **Acidum Tannicum**.

***Colocynthis** (*Colocyn.*), **Colocynth**, U. S. P. (*Colocynth Pulp*, *Bitter Apple*, *Colocynth Apple*).

ACTION AND USES: A powerful irritant hydragogue cathartic. Because of its drastic nature it is rarely used alone. The various compound pills listed below are needlessly complex and therefore irrational. Because of their irritant character, they are especially unsuited for continued use.

DOSAGE: 0.06 Gm. or 1 grain.

**Extractum Colocynthis* (*Ext. Colocynth.*), *Extract of Colocynth*, U. S. P. (*Powdered Extract of Colocynth*).—One Gm. of extract represents 4 Gm. of colocynth.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.).

**Extractum Colocynthis Compositum* (*Ext. Colocynth. Co.*), *Compound Extract of Colocynth*, U. S. P. (*Powdered Compound Extract of Colocynth*).—Extract of colocynth (16%), aloes (50%), resin of scammony (14%), cardamom seed and soap.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Pilulae Catharticae Compositae (Pil. Cathart. Co.), *Compound Cathartic Pills*, U. S. P.—Each pill contains compound extract of colocynth (0.08 Gm. or $1\frac{1}{3}$ grains), mild mercurous chloride (0.06 Gm. or 1 grain), resin of jalap (0.02 Gm. or $\frac{1}{3}$ grain) and gamboge (0.015 Gm. or $\frac{1}{4}$ grain).

DOSAGE: 2 pills (U. S. P.).

Pilulae Catharticae Vegetabiles (Pil. Cathart. Veget.), *Vegetable Cathartic Pills*, N. F.—Each pill contains compound extract of colocynth (0.06 Gm. or 1 grain), extract of hyoscyamus (0.03 Gm. or $\frac{1}{2}$ grain), resin of jalap (0.02 Gm. or $\frac{1}{3}$ grain), extract of leptandra and resin of podophyllum (each 0.015 Gm. or $\frac{1}{4}$ grain) with oil of peppermint.

DOSAGE: 2 pills (N. F.).

Pilulae Colocynthis Compositae (Pil. Colocynth. Co.), *Compound Pills of Colocynth*, N. F. (Pilulae Cocciae, *Cochia Pills*).—Each pill contains extract of colocynth (0.011 Gm. or $\frac{1}{6}$ grain), aloes and resin of scammony (each 0.13 Gm. or 2 grains), with oil of clove.

DOSAGE: 1 pill (N. F.).

Pilulae Colocynthis et Hyoscyami (Pil. Colocynth. et Hyoscy.), *Pills of Colocynth and Hyoscyamus*, N. F.—Each pill contains extract of colocynth (0.0065 Gm. or $\frac{1}{40}$ grain), aloes, resin of scammony and extract of hyoscyamus (each 0.097 Gm. or $1\frac{1}{2}$ grains) and oil of clove.

DOSAGE: 1 pill (N. F.).

Pilulae Colocynthis et Podophylli (Pil. Colocynth. et Podoph.), *Pills of Colocynth and Podophyllum*, N. F.—Each pill contains compound extract of colocynth (0.162 Gm. or $2\frac{1}{2}$ grains) and resin of podophyllum (0.016 Gm. or $\frac{1}{4}$ grain) with syrup.

DOSAGE: 1 pill (N. F.).

Pilulae Laxativae Post Partum (Pil. Lax. Post Part.), *Post Partum Laxative Pills*, N. F. (Barker's Post Partum Pills).—Each pill contains compound extract of colocynth (0.11 Gm. or $1\frac{2}{3}$ grains), Socotrine aloes (0.055 Gm. or $\frac{7}{8}$ grain), extract of nux vomica (0.025 Gm. or $\frac{3}{8}$ grain), resin of podophyllum and ipecac (each 0.005 Gm. or $\frac{1}{42}$ grain) and extract of hyoscyamus (0.008 Gm. or $\frac{1}{8}$ grain).

DOSAGE: 1 pill (N. F.).

Condurango (Conduran.), Condurango, N. F.—A bark.

ACTION AND USES: Bitter stomachic, introduced especially for cancer cachexia, but without advantage over gentian.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Condurango (Fldext. Condurango), Fluidextract of Condurango, N. F.—Condurango (100%).

Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

CONFECTIONES, CONFECTIONS (Conserves, Electuaries).—Soft, pasty solids consisting of active drugs mixed with sugar and honey.

Conium (Conium) Conium, N. F. (Poison Hemlock, *Conium maculatum*).—Unripe fruit.

ACTION AND USES: Source of coniine. Depresses the motor system, but has no definite therapeutic indication.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Extractum Conii (Ext. Conii), Extract of Conium, N. F.—A hydro-alcoholic extract diluted with glucose, yielding about 2 per cent. of coniine.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (N. F.).

Fluidextractum Conii (Fldext. Conii), Fluidextract of Conium, N. F.—Conium (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 0.2 Cc. or 3 minims (N. F.).

Convallariae Flores (Convallar. Flor.), Convallaria Flowers, N. F. (Lily-of-the-Valley Flowers).

ACTION AND USES: Believed to act as a cardiac tonic, similar to digitalis, but unreliable.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Fluidextractum Convallariae Florum (Fldext. Convall. Flor.), Fluidextract of Convallaria Flowers, N. F. (Fluidextractum Convallariae, N. F. III).—Convallaria flowers (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

Convallariae Radix (Convallar. Rad.), Convallaria Root, N. F. (Lily-of-the-Valley Root).—Rhizome and roots.

ACTION AND USES: Same as the flowers.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Fluidextractum Convallariae Radicis (*Fldext. Conval. Rad.*), *Fluidextract of Convallaria Root*, N. F.—Convallaria root (100%). Absolute alcohol content about 60 per cent. DOSAGE: 0.5 Cc. or 8 minims (N. F.).

**Copaiba* (*Copaib.*), *Copaiba*, U. S. P. (Balsam of Copaiba, *Copaiva*).—An oleoresin.

Pale yellow or brownish-yellow, viscid liquid, with an aromatic odor and a persistent, bitter, acrid taste. Insoluble in water and partly soluble in alcohol.

ACTION AND USES: A mild but disagreeable irritant and diuretic. Used especially in the treatment of gonorrhea, but of doubtful efficiency. The copaiba mixtures as listed below were formerly widely used, but are disagreeable, irrational and usually inefficient.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Massa Copaibae (*Mass. Copaib.*), *Mass of Copaiba*, N. F. (Solidified Copaiba).—Copaiba solidified by combination with magnesium oxide.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Mistura Copaibae (*Mist. Copaib.*), *Copaiba Mixture*, N. F. (Lafayette Mixture).—Copaiba (12.5%), spirit of nitrous ether (12.5%), compound tincture of lavender, solution of potassium hydroxide in syrup and mucilage of acacia. Absolute alcohol content about 20 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Mistura Copaibae et Opü (*Mist. Copaib. et Opü*), *Mixture of Copaiba and Opium*, N. F. (Chapman's Mixture).—Copaiba (25%), tincture of opium (3.2%), spirit of nitrous ether (25%), and compound tincture of lavender in mucilage of acacia and water. Absolute alcohol content about 29 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Coptis (*Coptis*), *Coptis*, N. F. (Goldthread).—Plant.

ACTION AND USES: Bitter stomachic, possessing no advantage over gentian.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Coptis (*Fldext. Copt.*), *Fluidextract of Coptis*, N. F.—Coptis (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

CORDIALES, CORDIALS.—Cordials are sweetened alcoholic liquids.

Coriandrum (Coriand.), Coriander, U. S. P. (Coriander Seeds).

ACTION AND USES: Aromatic and carminative.

Cornus (Corn.), Cornus, N. F. (Dogwood Bark).—Bark of the root.

ACTION AND USES: Without definite indications for its use; practically obsolete.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Corni (Fldext. Corni), Fluidextract of Cornus, N. F.—Cornus (100%). Absolute alcohol content about 35 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Corydalis (Coryd.), Corydalis, N. F. (Turkey Corn, Squirrel Corn).—Tubers.

ACTION AND USES: Formerly exploited as antiluetic, but useless and therefore harmful.

DOSAGE: 0.65 Gm. or 10 grains (N. F.).

Elixir Corydalis Compositum (Elix. Coryd. Co.), Compound Elixir of Corydalis, N. F.—Fluidextract of corydalis, (6%), fluidextract of stillingia (6%), fluidextract of xanthoxylum (3%), fluidextract of blue flag (9%), potassium iodide (5%), alcohol and aromatic elixir. Absolute alcohol content about 40 per cent.

ACTION AND USES: An irrational and harmful "alterative."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Fluidextractum Corydalis (Fldext. Corydal.), Fluidextract of Corydalis, N. F.—Corydalis (100%). Absolute alcohol content about 65 per cent.

DOSAGE: 0.65 Cc. or 10 minims (N. F.).

Cotarninae Hydrochloridum (Cotarn. Hydrochl.), Cotarnine Hydrochloride, U. S. P. (Cotarnine Chloride; Stypticin).

Yellow, odorless powder. Freely soluble in water and alcohol, yielding yellow solutions.

ACTION AND USES: Used to check functional uterine hemorrhage. The mechanism of its action has not been explained, and its efficiency is somewhat questionable.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Coumarinum (Coumar.), Coumarin, N. F.

Colorless, fragrant crystals, with a bitter, burning taste. Sparingly soluble in cold water; freely soluble in alcohol.

ACTION AND USES: Perfume and flavor.

Creosoti Carbonas (Creosot. Carb.), Creosote Carbonate, U. S. P.—A mixture of the carbonates of various constituents of creosote, chiefly guaiacol and cresol.

An almost colorless, almost tasteless, odorless viscid liquid. Insoluble in water; freely soluble in alcohol and miscible with chloroform and benzin.

ACTION AND USES: Passes stomach unchanged; hence is not a gastric irritant; decomposed in the intestines; used as an intestinal antiseptic.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

***Creosotum (Creosot.), Creosote, U. S. P. (Creasote).**—A mixture chiefly of guaiacol and cresol, obtained from wood-tar.

An almost colorless, oily liquid, with a penetrating, smoky odor, and a burning, caustic taste. Slightly soluble in water, and miscible with alcohol, ether or fixed or volatile oils.

ACTION AND USES: Antiseptic; formerly widely used as an intestinal antiseptic and externally to some extent as an antiseptic dressing.

DOSAGE: 0.25 Cc. or 4 minims (U. S. P.), in capsules.

Aqua Creosoti (Aq. Creosot.), Creosote Water, U. S. P.—Creosote (1%) in water.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Petroxolinum Creosoti (Petrox. Creosot.), Creosote Petroxolin, N. F. (Creosote Petrox).—Creosote (20%), oleic acid and liquid petroxolin.

***Cresol (Cresol), Cresol, U. S. P.**—A mixture of isomeric cresols, $C_6H_4(CH_3)OH$. Obtained from coal tar.

Colorless, yellowish or brownish liquid, darkening with age and exposure to light, and having a phenol-like odor. Soluble in water (1:50), usually forming a cloudy solution, and miscible with alcohol, ether, petroleum benzin or glycerin.

ACTION AND USES: Action similar to phenol. Used as a disinfectant and antiseptic. Approximately four times as germicidal as phenol, and no more toxic.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.).

***Liquor Cresolis Compositus (Liq. Cresol. Co.), Compound Solution of Cresol, U. S. P.**—Cresol (50%) with a potassium soap solution.

Creta Praeparata (Cret. Praep.), Prepared Chalk, U. S. P. (Drop Chalk).—A native calcium carbonate.

White, amorphous, odorless, tasteless powder. Almost insoluble in water, insoluble in alcohol. Decomposed and dissolved by dilute acids.

ACTION AND USES: Used internally as a mild alkali, as antacid and against diarrhea.

DOSAGE: 1 Gm. or 15 grains (U. S. P.) as a powder or in suspension.

Mistura Cretae (*Mist. Cret.*), *Chalk Mixture*, U. S. P.—Compound chalk powder (20%) with cinnamon water and water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Pulvis Cretae Aromaticus.—See under *Cinnamomum Saigonum*.

Pulvis Cretae Compositus (*Pulv. Cret. Co.*), *Compound Chalk Powder*, U. S. P.—Prepared chalk (30%), with acacia and sugar.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Crocus (Croc.), Crocus, N. F. (Saffron).—Stigmas of the flowers.

ACTION AND USES: Yellow coloring agent without definite therapeutic indications. Has been used as "stimulant and antispasmodic."

Tinctura Croci (*Tr. Croc.*), *Tincture of Saffron*, N. F.—Saffron (10%) in diluted alcohol. Absolute alcohol content about 45 per cent.

Cubeba (Cubeb.), Cubeb, U. S. P. (Cubebs).—Fruit.

ACTION AND USES: Formerly much used internally as antiseptic, diuretic and as a stimulant to the genito-urinary membranes; also to some extent as a stimulating expectorant. Probably of little value.

DOSAGE: 1 Gm. or 15 grains (U. S. P.). The oleoresin is preferred.

Fluidextractum Cubebae (*Fldext. Cubeb.*), *Fluidextract of Cubeb*, N. F.—Cubeb (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Oleoresina Cubebae (*Oleores. Cubeb.*), *Oleoresin of Cubeb*, U. S. P.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Tinctura Cubebae (*Tr. Cubeb.*), *Tincture of Cubeb*, N. F.—Cubeb (20%) in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Trochisci Cubebae (*Troch. Cubeb.*), *Troches of Cubeb*, U. S. P.—Each troche contains oleoresin of cubeb

(0.02 Gm.) with oil of sassafras, extract of glycyrrhiza and acacia, flavored with syrup of tolu.

***Cupri Sulphas (Cupr. Sulph.), Copper Sulphate, U. S. P. (Cupric Sulphate).— $\text{CuSO}_4 + 5\text{H}_2\text{O}$.**

Blue, odorless crystals or powder with a metallic taste. Freely soluble in water (1:2.5), and slightly soluble in alcohol (1:500).

ACTION AND USES: Astringent and in strong solutions corrosive. In large doses a gastric and intestinal irritant producing violent vomiting and purging. Considered one of the best antidotes in phosphorus poisoning.

DOSAGE: Emetic, 0.25 Gm. or 4 grains (U. S. P.).

Mistura Adstringens (Mist. Adstring.), Astringent Mixture, N. F. (Mistura Adstringens et Escharotica, N. F. III, Villate's Mixture).—A needlessly complex mixture containing the acetates and sulphates of copper and zinc in dilute acetic acid solution with lead sulphate in suspension.

Cypripedium (Cypriped.), Cypripedium, N. F. (Lady Slipper Root).—Rhizome and roots.

ACTION AND USES: No definite indications for its use. Has been described as "tonic, stimulant and diaphoretic."

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Fluidextractum Cypripedii (Fldext. Cypriped.), Fluidextract of Cypripedium, N. F.—Cypripedium (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Damiana (Damiana), Damiana, N. F. (Turnera).—Leaves.

ACTION AND USES: Ingredient of "lost manhood" nostrums; mildly irritant, but otherwise probably inert.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Damianae (Fldext. Damian.), Fluidextract of Damiana, N. F. (Fluidextractum Turnerae, N. F. III).—Damiana (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

DECOCTA, DECOCTIONS.—Aqueous preparations made by boiling drugs in water and straining. They must be freshly made from the drugs.

Delphinium (Delphin.), Larkspur Seed, N. F.

ACTION AND USES: Preparations are used externally to destroy pediculi of the hair and of the pubic region.

Tinctura Delphinii (*Tr. Delphin.*), *Tincture of Larkspur*, N. F.—Larkspur seed (10%) in alcohol. Absolute alcohol content about 90 per cent.

Dextrinum Album (*Dext. Alb.*), **White Dextrin**, N. F.—A variable mixture of soluble carbohydrates with unconverted starch.

White, odorless powder, with a sweetish taste. Partly soluble in cold water; completely soluble in three parts of water on boiling.

ACTION AND USES: Substitute for acacia and other gums. *Pasta Dextrinata* (*Past. Dextrin.*), *Dextrinated Paste*, N. F.—White dextrin (33%) in glycerin (33%) and water.

USES: Basis for dermatologic pastes.

Diacetylmorphina (*Diacetylmorph.*), **Diacetylmorphine**, U. S. P. (*Acetomorphine*, *Heroin*).—An alkaloid prepared from morphine by acetylation.

White, odorless powder. Very slightly soluble in water (1:1700) and soluble in alcohol (1:31) and in chloroform (1:1.4).

ACTION AND USES: Used as a respiratory sedative to relieve cough. With few exceptions, generally contraindicated in pulmonary tuberculosis. Like morphine, but more toxic and less valuable for the relief of pain. Is habit-forming. See **Morphina**.

DOSAGE: 0.003 Gm. or $\frac{1}{20}$ gr. (U. S. P.).

Diacetylmorphinae Hydrochloridum (*Diacetylmorph. Hydrochl.*), **Diacetylmorphine Hydrochloride**, U. S. P. (*Diacetylmorphine Chloride*; *Heroin Hydrochloride*).

White, odorless, bitter powder. Freely soluble (1:2) in water and soluble in alcohol (1:11).

DOSAGE: 0.003 Gm. or $\frac{1}{20}$ grain (U. S. P.).

Elixir Terpini Hydratis et Diacetylmorphinae (*Elix. Terpin. Hyd. et Diacetylmorph.*), *Elixir of Terpin Hydrate and Diacetylmorphine*, N. F. (*Elixir Terpini Hydratis cum Heroina* N. F. III).—Diacetylmorphine hydrochloride (0.027%) and elixir of terpin hydrate. Absolute alcohol content about 42 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.). This dose contains about 0.001 Gm. or $\frac{1}{60}$ grain of diacetylmorphine hydrochloride, and about 0.07 Gm. or $\frac{1}{6}$ grains of terpin hydrate.

Diastasum (*Diastasum*), **Diastase**, U. S. P.—A mixture containing amylolytic enzymes from malt.

Yellowish-white, odorless, tasteless powder or scales. Soluble in water; almost insoluble in alcohol.

ACTION AND USES: May be mixed with starchy foods to assist their conversion into sugar.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

***Digitalis (Digit.), Digitalis, U. S. P.** (Foxglove, Digitalis Leaves, Digitalis Folium P. I.).

ACTION AND USES: Direct cardiac stimulant and diuretic. Used in cardiac decompensation. Most effective when the heart action is rapid and very irregular (auricular fibrillation). Effective in the treatment of edema only when the latter is due to impaired circulation. Very likely to cause disturbance of gastro-intestinal tract.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.). Physicians are advised to specify preparations which have been biologically assayed.

Fluidextractum Digitalis (Fldext. Digital.), Fluidextract of Digitalis, U. S. P.—Digitalis (100%). May be biologically assayed by producing death in the frog. Absolute alcohol content about 70 per cent.

DOSAGE: 0.05 C.c. or 1 minim (U. S. P.).—Physicians should specify the biologically assayed fluidextract.

***Infusum Digitalis (Inf. Digit.), Infusion of Digitalis, U. S. P.**—Digitalis (1.5%), cinnamon water and water. This is an effective form of digitalis, but has no advantage over tinctures of proper potency, and is not so conveniently standardized.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Pilulae Digitalis, Scillae et Hydrargyri (Pil. Digit. Scill. et Hydrarg.), Pills of Digitalis, Squill and Mercury, N. F. (Niemeyer Pills for Dropsy, Guy's Pills).—Each pill contains digitalis, squill and mass of mercury (each 0.065 gm. or 1 grain) with clarified honey.

USES: Popular but needlessly complex cardiac diuretic.

DOSAGE: 1 pill (N. F.).

***Tinctura Digitalis (Tr. Digit.), Tincture of Digitalis, U. S. P.** (Digitalis Tinctura P. I.).—Digitalis (10%) in alcohol and water. May be biologically assayed by producing death in frogs. Absolute alcohol content about 70 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.). Physicians should specify the biologically assayed tincture.

Dioscorea (Dioscor.), Dioscorea, N. F. (Wild Yam Root, Colic Root).—Rhizome.

ACTION AND USES: No established indications for its use; formerly used in the treatment of "abdominal irritation."

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Dioscoreae (*Fldext. Dioscor.*), *Fluidextract of Dioscorea*, N. F.—Dioscorea (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Drosera (*Droser.*), *Drosera*, N. F. (Sundew).—Flowering plant.

ACTION AND USES: No established indications for its use (formerly used in respiratory diseases).

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Droserae (*Fldext. Droser.*), *Fluidextract of Drosera*, N. F.—Drosera (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Dulcamara (*Dulcam.*), *Bittersweet*, N. F.—Stems and branches.

ACTION AND USES: No established indications for its use. Formerly used as "alterative."

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Dulcamarae (*Fldext. Dulcam.*), *Fluidextract of Bittersweet*, N. F.—Bittersweet (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Echinacea (*Echin.*), *Echinacea*, N. F.—Rhizome and roots.

ACTION AND USES: The claims for this drug as an "alterative" and antisyphilitic are extravagant and unwarranted. There are no established indications for its use.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Echinaceae (*Fldext. Echin.*), *Fluidextract of Echinacea*, N. F.—Echinacea (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Elaterinum** (*Elaterin.*), *Elaterin*, U. S. P.—A neutral principle obtained from the squirting cucumber. The commercial product is variable and may at times be practically inert.

White, odorless powder, with a slightly acid, bitter taste. Insoluble in water; slightly soluble in alcohol (1:325).

ACTION AND USES: One of the most powerful and irritant of hydragogue cathartics, causing profuse watery evacuations.

DOSAGE: 0.003 Gm. or $\frac{1}{20}$ grain.

Trituratio Elaterini (*Trit. Elaterin.*), *Trituration of Elaterin*, U. S. P.—Elaterin (10%).

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

ELIXIRIA, ELIXIRS.—Elixirs are sweetened, aromatic, alcoholic liquids similar to cordials. In the administration of an elixir, the effects due to its alcohol content should be borne in mind. The statements of alcohol content of elixirs in this volume refer to percentage of absolute alcohol by volume.

For the various elixirs, see under the names of the principal constituents. For Elixir Aromaticum, Elixir Aromaticum Rubrum and Elixir Glycyrrhizae (Elixir Adjuvans), see under **Alcohol**; for Elixir Catharticum Compositum, see under **Frangula**; for Elixir Cinchonae Alkaloidorum and Elixir Cinchonae Alkaloidorum et Hypophosphitum, see under **Quininae Sulphas**; for Elixir Cinchonae Alkaloidorum et Ferri, see under **Ferri Phosphas**; for Elixir Cinchonae Alkaloidorum, Ferri, Bismuthi et Strychninae and Elixir Cinchonae Alkaloidorum, Ferri et Strychninae, see under **Strychninae Sulphas**; for Elixir Cinchonae Alkaloidorum, Ferri et Bismuthi, see under **Bismuthi et Sodii Tartras**; for Elixir Cinchonae Alkaloidorum, Ferri et Calcii Lactophosphatis, see under **Calcii Lactophosphas**; for Elixir Cinchonae Alkaloidorum, Ferri et Pepsini, see under **Pepsinum**; for Elixir Ferri, Quininae et Strychninae, and Elixir Gentianae et Ferri, see under **Ferri Citro-Chloridum**; for Elixir Formatum and Elixir Formatum Compositum, see under **Acidum Formicum**; for Elixir Gentianae et Ferri Phosphatis, see under **Ferri Phosphas**; for Elixir Glycerophosphatum Compositum, see under **Sodii Glycerophosphas**; for Elixir Hypophosphitum, see under **Calcii Hypophosphis**; for Elixir Hypophosphitum et Ferri, see under **Ferri Hypophosphis**; for Elixir Pepsini, Bismuthi et Strychninae, see under **Strychnina**; for Elixir Terpini Hydratis et Codeinae, see under **Codeina**; for Elixir Terpini Hydratis et Diacetylmorphinae, see under **Diacetylmorphinae Hydrochloridum**; for Elixir Trium Bromidorum, see under **Ammonii Bromidum**.

***Emetinae Hydrochloridum** (**Emet. Hydrochl.**), **Emetine Hydrochloride**, U. S. P.—The hydrochloride of the alkaloid emetine, obtained from ipecac.

White, or nearly white, odorless powder, darkening on exposure to light. Freely soluble in water or alcohol.

ACTION AND USES: Given hypodermically. An effective treatment of amebic dysentery and other amebic diseases.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain.

EMPLASTRA, PLASTERS.—Plasters are solid preparations for external use, serving either as simple adhesives or for the conveyance of drugs. The commercial sprays and plasters have largely displaced the extemporaneous prepared product. For the various plasters see under the names of the principal constituents. For *Emplastrum Elasticum*, see under *Resina*; for *Emplastrum Fuscum*, see under *Camphora*; for *Emplastrum Plumbi*, see under *Plumbi Oxidum*.

EMULSA, EMULSIONES, EMULSIONS.—Emulsions are aqueous preparations in which oils or resins are suspended by means of mucilaginous substances.

For the official emulsions see under the names of the respective oils and resins.

***Ergota (Ergot.), Ergot, U. S. P.** (*Secale Cornutum*, P. *Ergot of Rye, Spurred Rye*).—

ACTION AND USES: Causes powerful tonic, sometimes tetanic contraction of the uterus and of the blood vessel walls. Checks postpartum hemorrhage by contracting the uterus. Used for excessive menstrual flow and as a circulatory stimulant. Has also been tried but without success in a great variety of other diseases.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Extractum Ergotae (Ext. Ergot.), Extract of Ergot, U. S. P.—A pilular extract of ergot. Probably does not contain some of the valuable constituents of ergot (the alkaloids).

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Extractum Ergotae Aquosum (Ext. Ergot. Aq.), Aqueous Extract of Ergot, N. F. (*Ergoti Extractum P. I.*).—A purified aqueous extract.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

***Fluidextractum Ergotae (Fldext. Ergot.), Fluidextract of Ergot, U. S. P.** (*Secalis Cornuti Extractum Fluidum P. I.*).—Ergot (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Ergotae Ammoniata (Tr. Ergot. Ammoniated Tincture of Ergot, N. F.)—Ergot (25%) in ammonia water (10%) in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Eriodictyon (Eriodict.), **Eriodictyon**, U. S. P. (Yerba Santa).—Yerba santa leaves.

ACTION AND USES: Said to be expectorant. Its most remarkable property is its power of masking the taste of quinine.

DOSAGE: 1 Gm. or 15 grains.

Elixir Eriodictyi Aromaticum (Elix. Eriodict. Arom.), *Aromatic Elixir of Eriodictyon*, N. F. (Aromatic Elixir of Yerba Santa, Elixir Corrigen).—Fluidextract of eriodictyon (6%), compound elixir of taraxacum (about 44%) and syrup. Absolute alcohol content about 16 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Fluidextractum Eriodictyi (Fldext. Eriodict.), *Fluidextract of Eriodictyon*, U. S. P.—Eriodictyon (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Syrupus Eriodictyi Aromaticus (Syr. Eriodict. Arom.), *Aromatic Syrup of Eriodictyon*, N. F. (Aromatic Syrup of Yerba Santa, Syrupus Corrigen).—Fluidextract of eriodictyon, solution of potassium hydroxide, compound tincture of cardamom, oils of sassafras, lemon and clove, alcohol and syrup. Absolute alcohol content about 8 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

***Eucalyptol** (Eucalyptol), **Eucalyptol**, U. S. P. (Cineol).—A constituent of the volatile oil of eucalyptus.

A colorless liquid with a characteristic odor and a pungent taste, producing a cooling sensation in the mouth. Very slightly soluble in water but miscible with alcohol and chloroform and fixed or volatile oils.

ACTION AND USES: Employed as a local stimulant, antiseptic and deodorant, largely as a constituent of inhalations, particularly oily sprays.

DOSAGE: 0.3 Cc. or 5 minims (U. S. P.).

Nebula Eucalyptolis (Nebul. Eucalyptol.), *Eucalyptol Spray*, N. F.—Eucalyptol (5%) in light liquid petrolatum.

Petroxolinum Eucalyptolis (Petrox. Eucalyptol.), *Eucalyptol Petroxolin*, N. F. (Eucalyptol Petrox).—Eucalyptol (20%) in liquid petrolatum.

***Eucalyptus** (Eucalypt.), **Eucalyptus**, U. S. P. (Eucalyptus Leaves, Blue Gum Leaves).

ACTION AND USES: Principally those of the volatile oil. Not used as such.

Fluidextractum Eucalypti (*Fldext. Eucalypt.*), *Fluidextract of Eucalyptus*, U. S. P.—Eucalyptus (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Eugenol (Eugenol), Eugenol, U. S. P.—A constituent of oil of clove.

A nearly colorless, thin liquid, with a strong odor of cloves and a pungent, spicy taste.

ACTION AND USES: Same as those of oil of cloves, over which it has no material advantage.

DOSAGE: 0.2 Cc. or 3 minims.

Euonymus (Euonym.), Euonymus, N. F. (Wahoo Bark).—Bark of the root.

ACTION AND USES: Obsolete cathartic; toxic digitalis effects. Caution: The uncertain absorption of this drug makes its use inadvisable.

DOSAGE: 0.5 Gm. or 8 grains.

Extractum Euonymi (*Ext. Euonym.*), *Extract of Euonymus*, N. F. (Powdered Extract of Euonymus).—A hydro-alcoholic extract, 1 Gm. representing the water-alcohol-soluble constituents of 4 Gm. of euonymus.

DOSAGE: 0.125 Gm. or 2 grains (N. F.).

Fluidextractum Euonymi (*Fldext. Euonym.*), *Fluidextract of Euonymus*, N. F.—Euonymus (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 0.5 Cc. or 8 minims. (N. F.).

Eupatorium (Eupator), Eupatorium, N. F. (Boneset, Thoroughwort).—Leaves and flowering tops.

ACTION AND USES: Domestic diaphoretic "tea." No established indications for its use.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Eupatorii (*Fldext. Eupator.*), *Fluidextract of Eupatorium*, N. F.—Eupatorium (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Euphorbia Pilulifera (Euphorb. Pilul.), Euphorbia Pilulifera, N. F. (Pill Bearing Spurge).—Entire flowering plant.

ACTION AND USES: Irritant; formerly used in respiratory diseases, but without established indications.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Euphorbiae Piluliferae (*Fldext. Euphorb. Pilul.*), *Fluidextract of Euphorbia Pilulifera*, N. F.—

Euphorbia pilulifera (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

EXTRACTA, EXTRACTS (Solid Extracts).—Soft, solid (pilular extracts) or powdered (powdered extracts) preparations made by evaporating a solution of the soluble ingredients of vegetable or animal drugs at a low temperature. Powdered extracts are often made from the solid extracts by the addition of inert material. The strength of most extracts is four times that of the drug. For the various official extracts see under the names of the respective drugs.

Farfara (Farfar.), Coltsfoot, N. F. (Coltsfoot Leaves, Tussilago Leaves).

ACTION AND USES: Demulcent expectorant; no advantage over linseed.

DOSAGE: 4 Gm. or 1 drachm.

***Fel Bovis** (Fel Bovis), Oxgall, U. S. P.

A brownish-green or dark green, somewhat viscid liquid, with a peculiar odor and a disagreeable, bitter taste.

ACTION AND USES: Used in the form of the extract as intestinal antiseptic, cholagogue and laxative.

***Extractum Fellis Bovis** (Ext. Fel. Bov.), *Extract of Oxgall*, U. S. P. (Powdered Extract of Oxgall).—One Gm. of extract represents 8 Gm. of oxgall.

DOSAGE: 0.1 Gm. or 1½ grains.

Ferri Acetas, Ferric Acetate.— $\text{Fe}(\text{C}_2\text{H}_3\text{O}_2)_3$.

Liquor Ferri Acetatis (Liq. Ferr. Acet.), *Solution of Ferric Acetate*, N. F.—Contains anhydrous ferric acetate $\text{Fe}(\text{C}_2\text{H}_3\text{O}_2)_3$ (about 31%), corresponding to about 7.5 per cent. metallic iron, Fe.

DOSAGE: 0.3 Cc. of 5 minims (N. F.).

Liquor Ferri et Ammonii Acetatis.—See under **Ammonii Acetas**.

***Ferri Carbonas, Ferrous Carbonate.**— FeCO_3 .

***Massa Ferri Carbonatis** (Mass. Ferr. Carb.), *Mass of Ferrous Carbonate*, U. S. P. (Vallet's Mass).—Contains FeCO_3 (35%) with sugar and honey.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

NOTE:—The prescribing of iron can be greatly simplified without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

Mistura Ferri Composita (*Mist. Ferr. Co.*), *Compound Iron Mixture*, U. S. P. (Griffith's Mixture).—Ferrous carbonate (obtained by decomposing 0.6% ferrous sulphate with potassium carbonate), sugar, myrrh, spirit of lavender and rose water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

**Pilulae Ferri Carbonatis* (*Pil. Ferr. Carb.*), *Pills of Ferrous Carbonate*, U. S. P. (Chalybeate Pills, Blau's Pills, Ferruginous Pills).—Each pill contains FeCO_3 (0.06 Gm. or 1 grain).

USES: Convenient and effective form of iron.

DOSAGE: 2 pills (U. S. P.).

Ferri Carbonas Saccharatus (*Ferr. Carb. Sacch.*), *Saccharated Ferrous Carbonate*, U. S. P.—Ferrous carbonate FeCO_3 (not less than 15%) preserved with sugar.

A greenish-brown, odorless powder, with a taste at first sweet afterward ferruginous.

ACTION AND USES: Hematinic, like other iron salts, practically non-astringent and non-irritating.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

**Ferri Chloridum* (*Ferr. Chlor.*), *Ferric Chloride*, U. S. P. (Iron Perchloride, Sesquichloride of Iron).—Ferric chloride, FeCl_3 , with water of hydration.

Orange-yellow crystalline deliquescent pieces, odorless, and with a strongly styptic taste. Very soluble in water (1:0.2); freely soluble in alcohol.

ACTION AND USES: Used in the form of the tincture of ferric chloride as an astringent, especially as an application to the throat. Also hematinic, but relatively irritating.

Gossypium Stypticum (*Gossyp. Stypt.*), *Styptic Cotton*, N. F.—Purified cotton, impregnated with ferric chloride.

Liquor Ferri Chloridi (*Liq. Ferr. Chlor.*), *Solution of Ferric Chloride*, U. S. P. (Solution of Iron Perchloride).— FeCl_3 (about 10.5%).

DOSAGE: 0.1 Cc. or 1½ minims.

**Tinctura Ferri Chloridi* (*Tr. Ferr. Chlor.*), *Tincture of Ferric Chloride*, U. S. P.—Ferric chloride (about 13% corresponding to not less than 4.48 per cent. of iron). Made by diluting a solution of ferric chloride (35%) with alcohol. Absolute alcoholic content about 63 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

Tinctura Ferri Chloridi Aetherea (*Tr. Ferr. Chlor. Aeth.*), *Ethereal Tincture of Ferric Chloride*, N. F. (Bestuscheff's Tincture, Lamotte's Drops).—Solution of ferric chloride (6%) and ether (25%) in alcohol. Absolute alcohol content about 65 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Ferri Citras, Ferric Citrate

Liquor Ferri Citratis (*Liq. Ferr. Cit.*), *Solution of Ferric Citrate*, N. F.—Contains ferric citrate corresponding to not less than 7.25 per cent. metallic iron, Fe.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

Ferri Citro-Chloridum, Iron Citro-Chloride

Elixir Ferri, Quininae et Strychninae (*Elix. Ferr. Quin. et Strych.*), *Elixir of Iron, Quinine and Strychnine*, N. F.—Tincture of ferric citro-chloride (12.5%), quinine hydrochloride (0.875%), strychnine sulphate (0.0175%), compound spirit of orange, glycerin and water. Absolute alcohol content about 25 per cent.

USES: A popular but irrational "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Gentianae et Ferri (*Elix. Gent. et Ferr.*), *Elixir of Gentian and Iron*, N. F. (Elixir of Gentian with Tincture of Ferric Citro-Chloride).—Tincture of ferric citro-chloride (10%) and elixir of gentian. Absolute alcohol content about 20 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Tinctura Ferri Citro-Chloridi (*Tr. Ferr. Citro-Chlor.*), *Tincture of Ferric Citro-Chloride*, N. F. (Tasteless Tincture of Ferric Chloride, Tasteless Tincture of Iron).—A tincture of a complex iron salt, containing about 4 per cent. of iron (Fe) and rendered non-styptic by the presence of a citrate. Made from solution of ferric chloride (35%), sodium citrate (44%), alcohol and water. Absolute alcohol content about 15 per cent.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

***Ferri et Ammonii Citras** (*Ferr. et Ammon. Cit.*), **Iron and Ammonium Citrate**, U. S. P. (Soluble Ferric Citrate, Ammonio-Ferric Citrate).—Iron citrate rendered more readily soluble by the presence of ammonium citrate. Contains about 17 per cent. of iron, Fe.

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

Thin, transparent, garnet-red, odorless scales, with a saline, mildly ferruginous taste. Readily and completely soluble in water; insoluble in alcohol.

ACTION AND USES: Hematinic, practically non-astringent. Has the general properties of iron salts.

DOSAGE: 0.25 Gm. or 4 grains.

Vinum Ferri (Vin. Ferr.); *Wine of Iron, N. F.* (Wine of Citrate of Iron).—Iron and ammonium citrate (4%) in tincture of sweet orange peel, syrup, and sherry wine. Absolute alcohol content about 20 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Ferri et Quininae Citras (Ferr. et Quin. Cit.), Iron and Quinine Citrate, U. S. P. (Ferri et Quininae Citras Solubilis, U. S. P. VIII, Soluble Iron and Quinine Citrate).—Iron citrate and quinine citrate rendered more soluble by the presence of ammonium citrate. Contains not less than 13 per cent. of iron, Fe.

Thin, transparent, deliquescent, greenish or golden yellow, odorless scales, with a bitter, mildly ferruginous taste. Rapidly and completely soluble in cold water, partly soluble in alcohol.

ACTION AND USES: Used as a bitter iron tonic. Of doubtful value.

DOSAGE: 0.25 Gm. or 4 grains.

Vinum Ferri Amarum (Vin. Ferr. Amar.), *Bitter Wine of Iron, N. F.*—Iron and quinine citrate (5%) in tincture of sweet orange peel, syrup and sherry wine. Absolute alcohol content about 17 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Ferri Glycerophosphas (Ferr. Glycerophos.), Ferric Glycero-phosphate, N. F. (Ferric Glycerinophosphate).

Yellowish-green scales or powder, odorless and nearly tasteless. Freely soluble in water (1:2) and insoluble in alcohol.

ACTION AND USES: Reputed "tonic," but without advantage over the ordinary iron salts.

DOSAGE: 0.2 Gm. or 3 grains.

Ferri Hydroxidum, Ferric Hydroxide.

Ferri Hydroxidum cum Magnesii Oxido (Ferr. Hydrox. cum Mag. Oxid.), *Ferric Hydroxide with Magnesium Oxide*, U. S. P. (Arsenic Antidote, Ferric Hydrate with Magnesia).—A dilute solution of ferric sulphate and a mixture of magnesium oxide with water, kept in separate bottles. When mixed, they yield ferric hydroxide.

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

USES: Antidote for arsenic.

DOSAGE: 120 Cc. or 4 fluidounces.

Magma Ferri Hydroxidi (*Magn. Ferr. Hydrox.*), *Ferric Hydroxide Magma*, N. F. (*Ferri Hydroxidum*, U. S. P. VIII, *Ferric Hydroxide*).—A suspension of ferric hydroxide in water.

Ferri Hypophosphis (Ferr. Hypophos.), Ferric Hypophosphate, N. F.

White or grayish white, odorless and nearly tasteless powder. Very slightly soluble in water (1:2300); insoluble in alcohol.

ACTION AND USES: Reputed "tonic," but without advantage over the ordinary iron salts.

DOSAGE: 0.2 Gm. or 3 grains.

Elixir Ferri Hypophosphitis (*Elix. Ferr. Hypophos.*), *Elixir of Ferric Hypophosphate*, N. F.—Ferric hypophosphate (1.65%), potassium citrate (2.15%), water and aromatic elixir. Absolute alcohol content about 21 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Hypophosphitum et Ferri (*Elix. Hypophos. et Ferr.*), *Elixir of Hypophosphites and Iron*, N. F.—Calcium hypophosphate and sodium hypophosphate (each 1.75%), potassium hypophosphate and ferric hypophosphate (each 0.875%), potassium citrate, hypophosphorous acid, water, syrup and aromatic elixir. Absolute alcohol content about 10 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Liquor Ferri Hypophosphitis (*Liq. Ferr. Hypophos.*), *Solution of Ferric Hypophosphate*, N. F. (*Solution of Hypophosphate of Iron*).—Ferric hypophosphate (16.5%) dissolved by means of potassium citrate (21.5%) in glycerin and water.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Syrupus Ferri Hypophosphitis (*Syr. Ferr. Hypophos.*), *Syrup of Ferric Hypophosphate*, N. F.—Ferric hypophosphate (1.75%), dissolved by means of potassium citrate (2.5%), and made into a syrup flavored with orange flower water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

***Ferri Iodidum, Ferrous Iodide.— FeI_2 .**

Pilulae Ferri Iodidi (Pil. Ferr. Iod.), *Pills of Ferrous Iodide*, U. S. P.—Each pill represents ferrous iodide, FeI_2 (about 0.06 Gm. or 1 grain).

DOSAGE: 2 pills (U. S. P.).

Syrupus Ferri et Mangani Iodidi (Syr. Ferr. et Mangan. Iod.), *Syrup of Iron and Manganese Iodide*, N. F.—Ferrous iodide (10%) and manganese iodide (5%) in syrup.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Syrupus Ferri Iodidi** (Syr. Ferr. Iod.), *Syrup of Ferrous Iodide*, U. S. P. (Ferri Iodidi Syrupus P. I.).—Ferrous iodide, FeI_2 (about 5%).

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Ferri Lactas (Ferr. Lact.), Ferrous Lactate, N. F. (Iron Lactate).

Greenish white powder or crystalline masses, with a slight, characteristic odor and a mild, sweet ferruginous taste. Slowly soluble in water (1:40), freely soluble in solutions of alkali citrates, but almost insoluble in alcohol.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Elixir Ferri Lactatis (Elix. Ferr. Lact.), *Elixir of Iron Lactate*, N. F.—Iron lactate (1.75%), potassium citrate (5.25%) in water and aromatic elixir. Absolute alcohol content about 18 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Lactophosphas, Iron Lactophosphate.

Syrupus Ferri Lactophosphatis (Syr. Ferr. Lactophos.), *Syrup of Iron Lactophosphate*, N. F.—Iron lactophosphate (2.5%) in syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Malas, Ferrous Malate (Iron Malate).

Extractum Ferri Pomatum (Ext. Ferr. Pomat.), *Ferrated Extract of Apples*, N. F. (Ferri Malas Crudus, Crude Malate of Iron).—A pilular extract containing ferrous malate.

DOSAGE: 0.65 Gm. or 10 grains (N. F.).

Tinctura Ferri Pomata (Tr. Ferr. Pomat.), *Tincture of Ferrated Extract of Apples*, N. F. (Tinctura Ferri Malatis Crudi, Tincture of Crude Malate of Iron).—

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

Ferrated extract of apples (10%) in alcohol and cinnamon water. Absolute alcohol content about 10 per cent.
DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Nitrates, Ferric Nitrate.— $\text{Fe}(\text{NO}_3)_3$.

Liquor Ferri Nitratis (Liq. Ferr. Nit.), Solution of Ferric Nitrate, N. F.—Ferric nitrate, $\text{Fe}(\text{NO}_3)_3$, corresponding to not less than 1.3 per cent. of metallic iron, Fe.

USES: Astringent and styptic.

DOSAGE: 0.3 Cc. or 5 minims (N. F.).

Ferri Oxidum Saccharatum (Ferr. Oxid. Sacch.), Saccharated Ferric Oxide, N. F. (Soluble Ferric Oxide, Ferrum Oxydatum Saccharatum, "Eisenzucker").—Saccharated ferric oxide representing not less than 2.8 per cent. of Fe.

Syrupus Ferri Saccharati Solubilis (Syr. Ferr. Sacch. Sol.), Syrup of Soluble Saccharated Iron, N. F. (Syrupus Ferri Oxydati Solubilis, Syrup of Saccharated Oxide of Iron, Syrup of Soluble Oxide of Iron).—Saccharated ferric oxide (41.5%), syrup and water.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Oxychloridum, Ferric Oxychloride.

Liquor Ferri Oxychloridi (Liq. Ferr. Oxychlor.), Solution of Ferric Oxychloride, N. F.—Ferric oxychloride corresponding to about 3 per cent. of metallic iron, Fe, dissolved in a mixture of glycerin and water.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Ferri Oxysulphas, Ferric Oxysulphate.

Liquor Ferri Oxysulphatis (Liq. Ferr. Oxysulph.), Solution of Ferric Oxysulphate, N. F.—A solution of basic ferric sulphate.

USES: Superfluous astringent iron solution.

***Ferri Phosphas (Ferr. Phos.), Ferric Phosphate, U. S. P.** (Ferri Phosphas Solubilis, U. S. P. VIII, Soluble Ferric Phosphate).—Ferric phosphate rendered soluble by sodium citrate. Contains iron, Fe (not less than 12%).

Thin, bright green, transparent, odorless scales, with an acidulous, slightly saline taste. Freely and completely soluble in water; insoluble in alcohol.

ACTION AND USES: Hematinic. Only slightly astringent; no special advantage over other iron salts.

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Elixir Cinchonae Alkaloidorum et Ferri (Elix. Cinchon. et Ferr.), *Elixir of Cinchona Alkaloids and Iron*, N. F. (Elixir Cinchonae et Ferri N. F. III, Ferrated Elixir of Calisaya, Alkaloidal).—Ferric phosphate (3.5%), elixir of cinchona alkaloids (about 92%) and water. Absolute alcohol content about 20 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Ferri Phosphatis (Elix. Ferr. Phos.), *Elixir of Ferric Phosphate*, N. F.—Ferric phosphate (3.5%), water and aromatic elixir. Absolute alcohol content about 20 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Gentianae et Ferri Phosphatis (Elix. Gent. et Ferr. Phosph.), *Elixir of Gentian and Ferric Phosphate*, N. F. (Elixir Gentianae Ferratum).—Ferric phosphate (1.75%), elixir of gentian (about 95%) and water. Absolute alcohol content about 20 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Ferri, Quininae et Strychninae Phosphatum (Syr. Ferr. Quin. et Strych. Phos.), *Syrup of the Phosphates of Iron, Quinine and Strychnine*, N. F.—Ferric phosphate (2%), quinine (2.6%), strychnine (0.02%), phosphoric acid (5%) in glycerin, water and syrup.

USES: Needlessly complex "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Protochloridum, Ferrous Chloride (Iron Protochloride).

Liquor Ferri Protochloridi (Liq. Ferr. Protochlor.), *Solution of Ferrous Chloride*, N. F. (Solution of Protochloride of Iron).—A solution of ferrous chloride in glycerin and water.

USES: Superfluous astringent iron solution.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

Syrupus Ferri Protochloridi (Syr. Ferr. Protochlor.), *Syrup of Ferrous Chloride*, N. F. (Syrup of Protochloride of Iron).—Solution of ferrous chloride (5%) in glycerin, orange flower water and syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Pyrophosphas (Ferr. Pyrophos.), Ferric Pyrophosphate, N. F. (Soluble Ferric Pyrophosphate).

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

Thin, apple-green, transparent, odorless scales, with an acidulous, slightly saline taste. Freely and completely soluble in water and insoluble in alcohol.

ACTION AND USES: Used in iron elixirs; no special advantage over other soluble salts of iron.

DOSAGE: 0.25 Gm. or 4 grains.

Elixir Ferri Pyrophosphatis (Elix. Ferr. Pyrophos.), *Elixir of Ferric Pyrophosphate*, N. F. (Elixir of Pyrophosphate of Iron).—Ferric pyrophosphate (3.5%), water and aromatic elixir. Absolute alcohol content about 20 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Ferri Pyrophosphatis, Quininae et Strychninae (Elix. Ferr. Pyrophos. Quin. et Strych.), *Elixir of Ferric Pyrophosphate, Quinine and Strychnine*, N. F.—Ferric pyrophosphate (3.5%), quinine sulphate (0.875%), strychnine citrate (0.014%), ammonium citrate, oil of orange, alcohol, syrup and water. Absolute alcohol content about 23 per cent.

USES: A needlessly complex "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Ferri Subsulphas, Ferric Subsulphate.

Liquor Ferri Subsulphatis (Liq. Ferr. Subsulph.), *Solution of Ferric Subsulphate*, U. S. P. (Monsel's Solution, Solution of Basic Ferric Sulphate).

USES: Local styptic and astringent.

***Ferri Sulphas (Ferr. Sulph.), Ferrous Sulphate, U. S. P.** (Iron Protosulphate).— $\text{FeSO}_4 + 7\text{H}_2\text{O}$. (An impure product is commercially known as copperas).

Pale bluish-green, odorless crystals, with a saline styptic taste. Freely soluble (1:1.4) in water, insoluble in alcohol.

ACTION AND USES: Hematinic, especially in pills; but mainly used in making pharmaceutical preparations. Formerly used extensively as a disinfectant, but is only deodorant.

DOSAGE: 0.1 Gm. or 1½ grains (U. S. P.).

***Ferri Sulphas Exsiccatus (Ferr. Sulph. Exsic.), Exsiccated Ferrous Sulphate, U. S. P.** (Dried Ferrous Sulphate).—Contains not less than 80 per cent. of FeSO_4 .

Grayish-white powder.

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.) as pills.

Ferri Sulphas Granulatus (*Ferr. Sulph. Gran.*), *Granulated Ferrous Sulphate*, U. S. P. (Precipitated Ferrous Sulphate).—Purified ferrous sulphate.

Very pale, bluish-green, crystalline powder.

DOSAGE: 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.).

Pilulae Ferri, Quininae, Aloes et Nucis Vomicae (*Pil. Ferr. Quin. Aloe. et Nuc. Vom.*), *Pills of Iron, Quinine, Aloes and Nux Vomica*, N. F. (*Pilulae Quadruplices*, N. F. III, *Quadruplex Pills*, *Quatuor Pills*, *Pilulae Ferri et Quininae Compositae*).—Each pill contains exsiccated ferrous sulphate, quinine sulphate and aloes (each 0.065 Gm. or 1 grain), extract of nux vomica (0.016 Gm. or $\frac{1}{4}$ grain) and extract of gentian.

USES: Antiquated, complex and irrational "tonic."

DOSAGE: 1 pill (N. F.).

Ferri Tersulphas, *Ferric Sulphate* (Iron Tersulphate).

Liquor Ferri Tersulphatis (*Liq. Ferr. Tersulph.*), *Solution of Ferric Sulphate* U. S. P. (Solution of Iron Tersulphate).

USES: For making pharmaceutical preparations.

**Ferrum* (*Ferr.*), *Iron*, U. S. P.—Metallic iron, Fe, in the form of fine, bright, and non-elastic wire.

USES: For making the salts.

Ferrum Albuminatum, *Iron Albuminate*.

Liquor Ferri Albuminati (*Liq. Ferr. Albumin.*), *Solution of Albuminate of Iron*, N. F.—Contains albuminate of iron equivalent to about 0.4% of iron, Fe, in aromatic elixir, alcohol and water. Absolute alcohol content about 20 per cent.

USES: Non-astringent iron preparation; converted into simple ionic iron by dilute acids.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Ferrum Peptonatum, *Iron Peptonate*.

Liquor Ferri Peptonati (*Liq. Ferr. Pepton.*), *Solution of Peptonate of Iron*, N. F.—Contains a compound of iron and peptonized fresh egg albumin representing 0.4 per cent. metallic iron, Fe; flavored with oil of orange.

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

acetic ether and vanillin in alcohol and water. Absolute alcohol content about 14 per cent.

USES: Non-astringent inorganic iron preparation.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Liquor Ferri Peptonati et Mangani (Liq. Ferr. Pepton. et Mangani.), Solution of Peptonate of Iron and Manganese, N. F. (Liquor Ferri Peptonati cum Mangano, N. F. III).

—Solution of peptonate of iron with soluble manganese citrate (1%). Absolute alcohol content about 14 per cent.

USES: Combination of iron and manganese, of doubtful advantage over the solution of peptonate of iron.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

**Ferrum Reductum* (Ferr. Reduct.), Reduced Iron, U. S. P.

(Ferrum Redactum, Iron by Hydrogen, Quevenne's Iron).

—Metallic iron, Fe (not less than 90%), obtained by reduction of iron oxide by hydrogen.

Very fine, grayish-black, lusterless, odorless and tasteless powder. Insoluble in water or alcohol.

ACTION AND USES: A hematinic, relatively non-irritant; given preferably just before meals in the form of pills or powder enclosed in capsules or cachets.

DOSAGE: 0.06 Gm. or 1 grain.

Pilulae Ferri, Quininae, Strychninae et Arseni Fortiores (Pil. Ferr., Quin., Strych. et Arsen. Fort.), Stronger Pills of Iron, Quinine, Strychnine, and Arsenic, N. F. (*Pilulae Metallorum*, N. F. III, Metallic Pills, *Pilulae Metallorum Amarae*, Bitter Metallic Pills).—Each pill contains reduced iron and quinine sulphate (each 0.065 Gm. or 1 grain), strychnine and arsenic trioxide (each 0.0032 Gm. or $\frac{1}{20}$ grain) with clarified honey.

USES: Complex and irrational tonic.

DOSAGE: 1 pill (N. F.).

Pilulae Ferri, Quininae, Strychninae et Arseni Mites (Pil. Ferr., Quin., Strych. et Arsen. Mit.), Mild Pills of Iron, Quinine, Strychnine and Arsenic, N. F. (Aitken Tonic Pills).—Each pill contains reduced iron (0.045 Gm. or $\frac{1}{3}$ grain), quinine sulphate (0.065 Gm. or 1 grain), strychnine and arsenic trioxide (each 0.0013 Gm. or $\frac{1}{60}$ grain) with clarified honey.

USES: Complex and irrational tonic.

DOSAGE: 1 pill (N. F.).

NOTE:—The prescribing of iron can be greatly simplified, without sacrificing anything of real value, if physicians will confine themselves to the starred preparations.

Ficus (Ficus), Fig, N. F.

ACTION AND USES: Mild laxative.

FLUIDEXTRACTA, FLUIDEXTRACTS.—Fluidextracts are concentrated liquid preparations of vegetable drugs, generally containing alcohol either as a solvent or as a preservative. One Cc. represents the extractives from 1 Gm. of drug.

The statements in this book regarding alcohol content are approximate and are based on the alcohol contents of the menstruum used. For the official preparations see under the names of the respective drugs. For Fluidextractum Aromaticum, see under Cinnamomum Saigonicum.

FLUIDGLYCERATA, FLUIDGLYCERATES.—Preparations intended to be of the same drug strength as fluidextracts—that is, representing 100 Gm. of drug in 100 Cc. of preparation—but containing about 50 per cent. by volume of glycerin and no alcohol. For the various fluidglycerates, see under the names of the respective drugs.

Foeniculum (Foenic.), Fennel, U. S. P. (Fennel Seed).

ACTION AND USES: Aromatic and carminative.

DOSAGE: 1 Gm. or 15 grains.

***Formaldehydum, Formaldehyde.—CH₂O.**

**Liquor Formaldehydi (Liq. Formaldehyd.), Solution of Formaldehyde, U. S. P.—CH₂O (not less than 37%) with methyl alcohol and water.*

ACTION AND USES: A powerful germicide, especially valuable in the form of gas for its penetrating power. Its irritant actions generally preclude its use on the body.

For disinfecting rooms, from 150 Cc. or 5 fluidounces to 1 quart or 1 liter of the official formaldehyde solution are allowed to each 1,000 cubic feet of space by various authorities. Various methods have been employed. One is to allow the solution to evaporate from sheets suspended in the tightly closed room. Another is to generate the gas in the room by treating the solution with potassium permanganate, allowing about 110 Gm. or 3½ ounces of potassium permanganate to 500 Cc. or 1 pint of formaldehyde solution. A third method utilizes the reaction between formaldehyde solution and sodium dichromate in the presence of sulphuric acid, used in the proportion of 45 Cc.

or 1½ fluidounces of sulphuric acid and 280 Gm. or 9 ounces of sodium dichromate to 500 Cc. or 1 pint of formaldehyde solution. These methods are not effective if the temperature of the room be below 60 F. or the relative humidity below 65 per cent.

Frangula (Frang.), Frangula, U. S. P. (Buckthorn Bark).

ACTION AND USES: Similar to those of cascara sagrada; used principally as fluidextract.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Catharticum Compositum (Elix. Cathart. Co.), Compound Cathartic Elixir, N. F.—Fluidextract of frangula (12.5%), fluidextract of senna (10%), fluidextract of rhubarb (6.2%), spirit of peppermint, solution of potassium hydroxide and aromatic elixir. Absolute alcohol content about 25 per cent.

USES: Needlessly complicated cathartic.

DOSAGE: Aperient, 4 Cc. or 1 fluidrachm (N. F.); cathartic, 12 Cc. or 3 fluidrachms (N. F.).

Fluidextractum Frangulae (Fldext. Frangul.), Fluidextract of Frangula, U. S. P. (Fluidextract of Buckthorn Bark).—Frangula (100%). Absolute alcohol content about 20 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Fraxinus (Fraxin.), White Ash Bark, N. F.

ACTION AND USES: Formerly used as tonic, astringent and purgative. No established indications for its use.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Vinum Fraxini (Vin. Frax.), Wine of White Ash, N. F.—White ash bark (50%), in alcohol and sherry wine. Absolute alcohol content about 30 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Fucus (Fucus), Fucus, N. F. (Bladderwrack).—A sea weed.

ACTION AND USES: Formerly used against goiters and tuberculous glands; may be of some value through its iodine content, but has no special advantage over other iodine preparations.

DOSAGE: 0.65 Gm. or 10 grains (U. S. P.).

Fluidextractum Fuci (Fldext. Fuci), Fluidextract of Fucus, N. F.—Fucus (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 0.65 Cc. or 10 minims (N. F.).

Galangal (Galang.), Galangal, N. F.—Rhizome.

ACTION AND USES: Aromatic carminative, resembling ginger and without advantage over the latter.

DOSAGE: 1 Gm. or 15 grains.

Galega (Galeg.), Galega, N. F. (European Goat's Rue).—Flowering tops.

ACTION AND USES: Exploited as a galactagogue and nerve stimulant; has no established value.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Galegae (Fldext. Galeg.), Fluidextract of Galega, N. F.—Galega (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Galla (Gall.), Nutgall, U. S. P. (Aleppo Galls, Smyrna Galls).

Incompatible with the salts of iron, lead, copper or silver.

ACTION AND USES: Astringent. Used especially as ointment, but has no advantage over tannic acid.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Tinctura Gallae (Tr. Gall.), Tincture of Nutgall, N. F.—Nutmall (20%) in glycerin and alcohol. Absolute alcohol content about 85 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Unguentum Gallae (Ung. Gall.), Nutgall Ointment, U. S. P.—Nutmall (20%) in ointment.

Gambir (Gambir), Gambir, U. S. P. (Pale Catechu).—A dried extract.

ACTION AND USES: Astringent; used against diarrhea.

DOSAGE: 1 Gm. or 15 grains (in form of tincture).

Pulvis Gambir Compositus (Pulv. Gambir Co.), Compound Powder of Gambir, N. F. (Pulvis Catechu Compositus, N. F. III).—Gambir (40%), kino (20%) and krameria (20%), with Saigon cinnamon and nutmeg.

USES: Needlessly complex intestinal astringent.

DOSAGE: 1.3 Gm. or 20 grains (N. F.).

Tinctura Gambir Composita (Tr. Gambir Co.), Compound Tincture of Gambir, U. S. P. (Compound Tincture of Pale Catechu).—Gambir (5%) with Saigon cinnamon in diluted alcohol. Absolute alcohol content about 47 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Trochisci Gambir (*Troch. Gambir*), *Troches of Gambir*, N. F.—Each troche contains gambir (0.06 Gm. or 1 grain) with tragacanth, sugar and oil of cinnamon.

DOSAGE: 1 troche (N. F.).

***Gelatinum (Gelatin), Gelatin, U. S. P.**—Obtained from skin, ligaments, bones, etc.

Insoluble in cold water but swells and softens when immersed in it; soluble in hot water and in glycerin; insoluble in alcohol and similar solvents.

ACTION AND USES: Used as a food product. Also, by mouth or by injection, with the intention of increasing the coagulability of the blood; of doubtful value. May contain tetanus spores.

Gelatinum Glycerinatum (*Gelat. Glycerin.*), *Glycerinated Gelatin, U. S. P.*—Consists of equal parts of gelatin and glycerin.

USES: Base for suppositories, bougies, etc.

Gelsemium (Gelsem.), Gelsemium, U. S. P. (Yellow Jasmine Root, Yellow Jessamine).

ACTION AND USES: Used in migraine and neuralgia, and in the treatment of ovarian, rheumatic and uterine pain. Efficiency uncertain. Untoward symptoms sometimes result from comparatively small doses.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.). (Not used as such).

Extractum Gelsemii (*Ext. Gelsem.*), *Extract of Gelsemium, U. S. P.* (Powdered Extract of Gelsemium).—One Gm. of extract represents 4 Gm. of gelsemium.

DOSAGE: 0.01 Gm. or $\frac{1}{6}$ grain (U. S. P.).

Fluidextractum Gelsemii (*Fldext. Gelsem.*), *Fluidextract of Gelsemium, U. S. P.*—Gelsemium (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 0.03 Cc. or $\frac{1}{2}$ minim (U. S. P.).

Tinctura Gelsemii (*Tr. Gelsem.*), *Tincture of Gelsemium, U. S. P.*—Gelsemium (10%) in alcohol and water. Absolute alcohol content about 62 per cent.

DOSAGE: 0.25 Cc. or 4 minims (U. S. P.).

***Gentiana (Gentian.), Gentian, U. S. P.** (Yellow Gentian Root).—Rhizome and roots.

ACTION AND USES: Probably the most widely used of the simple bitters.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Gentianae (*Elix. Gent.*), *Elixir of Gentian*, N. F.—Fluidextract of gentian (3.5%), compound spirit of cardamom, sodium citrate, glycerin, syrup, alcohol and water. Absolute alcohol content about 21 per cent.

USES: An agreeable aromatic bitter stomachic.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Gentianae et Ferri, N. F.—See under *Ferri Citro-Chloridum*.

Elixir Gentianae et Ferri Phosphatis, N. F.—See under *Ferri Phosphas*.

Elixir Gentianae Glycerinatum (*Elix. Gent. Glycerin.*), *Glycerinated Elixir of Gentian*, N. F.—Fluidextract of gentian (1%), fluidextract of taraxacum (1.5%), acetic ether, phosphoric acid, tincture of sweet orange peel, compound tincture of cardamom, glycerin, sugar and sherry wine. Absolute alcohol content about 11 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms.

**Extractum Gentianae* (*Ext. Gentian.*), *Extract of Gentian*, U. S. P.—A pilular extract.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Fluidextractum Gentianae (*Fldext. Gentian.*), *Fluidextract of Gentian*, U. S. P.—Gentian (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Infusum Gentianae Compositum (*Inf. Gent. Co.*), *Compound Infusion of Gentian*, N. F.—Gentian (3%), coriander and bitter orange peel in diluted alcohol and water. Absolute alcohol content about 12 per cent.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Tinctura Amara (*Tr. Amar.*), *Bitter Tincture*, N. F. (Stomachic Tincture, Bitter Stomachic Drops, Stomach Drops).—Gentian (6%), centaury (6%), zedoary (2%) and bitter orange peel in alcohol and water. Absolute alcohol content about 60 per cent.

USES: Bitter stomachic, complex and without special value.

DOSAGE: 2 Cc. or 30 minims (N. F.).

**Tinctura Gentianae Composita* (*Tr. Gentian. Co.*), *Compound Tincture of Gentian*, U. S. P.—Gentian (10%), bitter orange peel and cardamom seed in glycerin, alcohol and water. Absolute alcohol content about 46 per cent.

USES: Aromatic bitter.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Geranium (Geran.), Geranium, N. F. (Cranesbill).—Rhizome.

ACTION AND USES: Obsolete intestinal astringent, without special advantage over krameria, etc.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Geranii (Fldext. Geran.), Fluidextract of Geranium, N. F.—Geranium (100%). Absolute alcohol content about 50 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Glucosum (Glucos.), Glucose, U. S. P. (Syrupy Glucose, Liquid Glucose).—Chiefly dextrose (d-glucose) and dextrins.

Colorless or nearly colorless, odorless or nearly odorless, thick, syrupy, sweet liquid. Soluble in water, sparingly soluble in alcohol.

ACTION AND USES: Pill excipient. Pure dextrose is also employed in saline injections.

***Glycerinum (Glycerin.), Glycerin, U. S. P. (Glycerol).**— $C_3H_5(OH)_3$.

A colorless, syrupy, practically odorless liquid, with a sweet taste, and producing a sensation of warmth in the mouth. Miscible with water and alcohol; insoluble in chloroform or ether.

ACTION AND USES: Used as solvent, sweetening agent, demulcent and emollient. The enema or suppositories are promptly evacuant.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

**Suppositoria Glycerini (Supp. Glycerin.), Suppositories of Glycerin, U. S. P.*—Glycerin with a stearic acid soap.

USES: Rectal evacuant.

Suppositoria Boroglycerini (Suppos. Boroglycer.), Suppositories of Boroglycerin, N. F.—Glycerite of boroglycerin (7.5%) with glycerinated gelatin and glycerin.

USES: Antiseptic evacuant.

GLYCERITA, GLYCERITES.—Solutions of medicinal substances in glycerin. For the various glycerites, see under the names of the principal ingredients. For Glyceritum Boroglycerini, see under **Acidum Boricum**; for Glyceritum Vitelli, see under **Ovi Vitellum Recens**.

***Glycerylis Nitras, Glyceryl Trinitrate (Nitroglycerin, Trinitrin, Glonoin).**— $C_3H_5(NO_3)_3$.

ACTION AND USES: Vasodilator, acting more slowly than amyl nitrite.

Pilulae Glycerylis Nitratis (*Pil. Glycer. Nit.*), *Pills of Nitroglycerin*, N. F. (*Pilulae Glonoini*, N. F. III, *Pills of Glonoin*).—Each pill contains spirit of glyceryl trinitrate and althaea (each 0.065 Gm. or 1 grain) with confection of rose.

USES: Inferior to the tablets.

DOSAGE: 1 pill (N. F.).

**Spiritus Glycerylis Nitratis* (*Sp. Glyceryl. Nit.*), *Spirit of Glyceryl Trinitrate*, U. S. P. (*Spirit of Glonoin*, *Spirit of Nitroglycerin*).—Glyceryl trinitrate (about 1%) in alcohol.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.) dropped on the tongue.

**Glycyrrhiza* (*Glycyrrh.*), *Glycyrrhiza*, U. S. P. (*Licorice*, *Liquorice Root*).

ACTION AND USES: Used to disguise the taste of drugs and as demulcent expectorant. The preparations are incompatible with acids.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Elixir Glycyrrhizae and *Elixir Glycyrrhizae Aromaticum*.—See under *Alcohol*.

Elixir Glycyrrhizae Aquosum (*Elix. Glycyrrh. Aq.*), *Aqueous Elixir of Glycyrrhiza*, N. F. (*Aqueous Elixir of Licorice*).—Fluidextract of glycyrrhiza (15%), compound spirit of cardamom, stronger orange flower water, glycerin, syrup and distilled water. Absolute alcohol content about 3 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms.

Extractum Glycyrrhizae (*Ext. Glycyrrh.*), *Extract of Glycyrrhiza*, U. S. P. (*Extract of Licorice*).—The commercial extract of glycyrrhiza. Incompletely soluble in water (60%).

Extractum Glycyrrhizae Purum (*Ext. Glycyrrh. Pur.*), *Pure Extract of Glycyrrhiza*, U. S. P.—A pilular extract.

**Fluidextractum Glycyrrhizae* (*Fldext. Glycyrrh.*), *Fluidextract of Glycyrrhiza*, U. S. P.—Glycyrrhiza (100%). Absolute alcohol content about 20 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Fluidglyceratum Glycyrrhizae (*Fldglycer. Glycyrrh.*), *Fluidglycerate of Glycyrrhiza*, N. F. (*Fluidglycerate of Licorice*).—Glycyrrhiza (100%) in glycerin and water, made alkaline with ammonia water.

DOSAGE: 2 Cc. or 30 minims.

Mistura Glycyrrhizae Composita, U. S. P.—See under Opium.

Pulvis Glycyrrhizae Compositus, U. S. P.—See under Senna.

Syrupus Glycyrrhizae (Syr. *Glycyrrhiz.*), *Syrup of Glycyrrhiza*, N. F. (Syrup of Licorice).—Fluidglycerate of glycyrrhiza (25%) in syrup.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Glycyrrhizinum Ammoniatum (Glycyrrh. Ammon.), *Ammoniated Glycyrrhizin*, U. S. P.—The sweet principle of glycyrrhiza, combined with ammonia.

Dark brown or brownish-red, odorless scales, with a very sweet taste. Freely soluble in water and soluble in alcohol. Incompatible with acids.

ACTION AND USES: Intended as substitute for extract of glycyrrhiza, over which it presents no advantage.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Gossypii Cortex (Gossyp. Cort.), *Cotton Root Bark*, N. F.—Recently gathered or dried bark of the root.

ACTION AND USES: Exploited as emmenagogue, but without established value.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Gossypii Corticis (Fldext. *Gossyp. Cort.*), *Fluidextract of Cotton Root Bark*, N. F.—Cotton root bark (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Gossypium Purificatum (Gossyp. Purif.), *Purified Cotton*, U. S. P. (Absorbent Cotton).

ACTION AND USES: Used as an absorbent surgical dressing and as a filtering agent.

Gossypium Stypticum.—See under *Ferri Chloridum*.

Granatum (Granat.), *Pomegranate*, U. S. P. (Pomegranate Bark).

ACTION AND USES: The decoction or fluidextract expels tapeworm. Both are strongly astringent and often produce vomiting.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Fluidextractum Granati (Fldext. *Granat.*), *Fluidextract of Pomegranate*, U. S. P.—Pomegranate (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Grindelia (Grindel.), Grindelia, U. S. P.—Leaves and flowering tops.

ACTION AND USES: Used with doubtful results in spasmodic asthma, whooping cough, bronchitis, and hay fever, and locally against ivy poisoning. It is a feeble drug and its action has not been clearly defined.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Fluidextractum Grindeliae (Fldext. Grindel.), Fluidextract of Grindelia, U. S. P.—Grindelia (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Guaiaci Lignum (Guaiac. Lig.), Guaiac Wood, N. F. (Lignum Vitae).

ACTION AND USES: Reputed "alterative," but without established value. No advantage over guaiac resin.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

***Guaiacol (Guaiacol), Guaiacol, U. S. P.**—Obtained from wood-tar creosote or prepared synthetically.

Colorless or yellowish solid or liquid, with an aromatic odor and a strong phenolic taste. Soluble in water (1:53), freely soluble in glycerin (1:0.8) and miscible with alcohol, chloroform or ether.

ACTION AND USES: Used as intestinal antiseptic and bronchial stimulant, similar to creosote; less irritant but also less active. Inunction with 50 per cent. solution in oil is antipyretic but it is less powerful and more uncertain than acetphenetidin, etc. Action is difficult to control and collapse may occur.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

***Guaiacolis Carbonas (Guaiacol. Carb.), Guaiacol Carbonate, U. S. P.**

White, nearly odorless and nearly tasteless powder. Soluble in alcohol (1:60) and in ether (1:18); insoluble in water.

ACTION AND USES: Being insoluble in water, it is less disagreeable and less irritating than guaiacol, but liberates this substance in the intestines. Used as intestinal antiseptic and against bronchitis.

DOSAGE: 1 Gm. or 15 grains (U. S. P.). in powders or cachets.

Guaiacum (Guaiac.), Guaiac, U. S. P. (Guaiac Resin).

ACTION AND USES: Formerly used as an "alterative," against syphilis, chronic rheumatism and gout; and as throat lozenges for its mildly irritant action. It produces

no definite systemic effects, except as a mild laxative and possibly diuretic. Its value is very doubtful. The ammoniated tincture is the most active preparation.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Gargarisma Guaiaci Compositum (*Garg. Guaiac. Co.*), *Compound Gargle of Guaiac, N. F.*—Ammoniated tincture of guaiac (10%), compound tincture of cinchona (10%), potassium chlorate (4%), with clarified honey and oil of peppermint in water. Absolute alcohol content about 12 per cent.

USES: Complex and irrational mixture.

Glyceritum Guaiaci (*Glycer. Guaiac.*), *Glycerite of Guaiac, N. F.*—Guaiac (8.5%) and solution of potassium hydroxide in glycerin and water.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Mistura Guaiaci (*Mist. Guaiac.*), *Mixture of Guaiac, N. F.*—Tincture of guaiac (12.5%) in clarified honey and cinnamon water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Petroxolinum Guaiacolis (*Petrox. Guaiacol.*), *Guaiacol Petroxolin, N. F.* (Guaiacol Petrox).—Guaiacol (20%) in oleic acid and liquid petroxolin.

Tinctura Guaiaci (*Tr. Guaiac.*), *Tincture of Guaiac, U. S. P.*—Guaiac (20%) in alcohol. Absolute alcohol content about 80 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Tinctura Guaiaci Ammoniata (*Tr. Guaiac. Ammon.*), *Ammoniated Tincture of Guaiac, U. S. P.*—Guaiac (20%) in aromatic spirit of ammonia. Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Guaiaci Composita (*Tr. Guaiac. Co.*), *Compound Tincture of Guaiac, N. F.* (Dewees' Tincture of Guaiac).—Guaiac (12.5%), potassium carbonate and pimenta in diluted alcohol. Absolute alcohol content about 45 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Guarana (*Guarana*), *Guarana, U. S. P.*—Paste consisting chiefly of crushed seeds yielding not less than 4 per cent. of caffeine.

ACTION AND USES: Used for its caffeine. There is no reason for using it instead of caffeine.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Elixir Guaranae (*Elix. Guar.*), *Elixir of Guarana*, N. F.—Fluidextract of guarana (20%), compound elixir of taraxacum (60%) and aromatic elixir. Absolute alcohol content about 34 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Fluidextractum Guaranae (*Fldest. Guaran.*), *Fluidextract of Guarana*, U. S. P.—Guarana (100%) yielding about 4 per cent. of caffeine. Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Gutta Percha (*Gutta Percha*), *Gutta Percha*, N. F.—A purified, coagulated tree sap.

ACTION AND USES: Used for technical purposes in dentistry.

Liquor Guttae Perchae (*Liq. Gut. Perch.*), *Solution of Gutta Percha*, N. F.—A clarified solution of gutta percha in chloroform.

Haematoxylon (*Haematox.*), *Haematoxylon*, N. F. (Logwood).

ACTION AND USES: Intestinal astringent, without advantage over krameria, etc.

Extractum Haematoxyli (*Ext. Haematox.*), *Extract of Hematoxylon*, N. F.—An aqueous extract.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Hamamelidis Folia (*Hamamel. Fol.*), *Hamamelis Leaves*, N. F. (Witch Hazel Leaves).

ACTION AND USES: Astringent, without advantage over other tannin-bearing drugs (krameria, etc.).

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Hamamelidis Foliorum (*Fldest. Hamamel. Fol.*), *Fluidextract of Hamamelis Leaves*, N. F.—Hamamelis leaves (100%). Absolute alcohol content about 25 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Hamamelis (*Witch Hazel*).

Aqua Hamamelidis (*Aq. Hamam.*), *Hamamelis Water*, U. S. P. (Witch Hazel Water, Distilled Extract of Witch Hazel).—Witch hazel bark, twigs, smaller stems or entire shrub, distilled with water and preserved with about 14 per cent. of alcohol.

USES: Employed externally, for contusions. It owes its feeble activity mainly to the alcohol.

Helianthemum (Helianth.), Helianthemum, N. F. (Rock-Rose, Frost-Weed).—The dried herb.

ACTION AND USES: Obsolete astringent, tonic, alterative and emetic; unreliable and without established value.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Helianthemis (Fldext. Helianth.), Fluidextract of Helianthemum, N. F.—Helianthemum (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Helonias (Helon.), Helonias, N. F. (False Unicorn).—Rhizome and roots.

ACTION AND USES: Ingredient of "female remedies"; without established value.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Heloniatis (Fldext. Helon.), Fluidextract of Helonias, N. F.—Helonias (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Hexamethylenamina (Hexam.), Hexamethylenamine, U. S. P.** (Hexamethylene-tetramine; also sold as urotropine, aminoform, formamin, formin, cystamin, cystogen, urisol and uritone).— $(\text{CH}_2)_6\text{N}_4$, a condensation product of ammonia and formaldehyde.

Colorless crystals or white powder, odorless and with a sweetish taste. Freely soluble in water (1:1.5) and in alcohol (1:12.5). Incompatible with acids, with ammonium salts, with tannin and with mercuric chloride.

ACTION AND USES: Useful urinary antiseptic, liberating formaldehyde in the presence of acids (not otherwise). Also used as prophylactic against infection in bile and in other body fluids, but inefficient since these fluids are not acid.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), in solution. When sodium acid phosphate is given to render the urine acid, hexamethylenamine should not be taken until the phosphate has left the stomach.

***Homatropinae Hydrobromidum (Homatrop. Hydrobr.), Homatropine Hydrobromide, U. S. P.** (Homatropine Bromide).

White, odorless powder. Freely soluble in water (1:6) and soluble in alcohol (1:40).

ACTION AND USES: Used as mydriatic and cycloplegic. Its effects resemble those of atropine but occur more promptly and disappear in shorter time.

DOSAGE: 0.5 mg. or $\frac{1}{120}$ grain (U. S. P.). May be used in aqueous solution of 2 per cent. strength, or a drop of a 1:500 solution may be introduced into the conjunctival sac every five minutes for five times to produce a maximum dilatation in three quarters of an hour.

Humulus (Humul.), Hops, U. S. P.

ACTION AND USES: Formerly used as aromatic bitter stomachic.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Elixir Humuli (Elix. Humul.), Elixir of Hops, N. F.—Fluidextract of hops (12.5%), compound elixir of taraxacum, tincture of vanilla and aromatic elixir. Absolute alcohol content about 25 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Fluidextractum Humuli (Fldext. Humul.), Fluidextract of Hops, N. F.—Hops (100%). Absolute alcohol content about 50 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Humuli (Tr. Humul.), Tincture of Hops, N. F.—Hops (20%) in diluted alcohol. Absolute alcohol content about 45 per cent.

USES: Bitter stomachic.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Hydrangea (Hydrang.), Hydrangea, N. F. (Seven-Barks).—Rhizome and roots.

ACTION AND USES: Obsolete cathartic, diuretic and diaphoretic; without established value.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Hydrangeae (Fldext. Hydrang.), Fluidextract of Hydrangea, N. F.—Hydrangea (100%). Absolute alcohol content about 50 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Hydrargyri Chloridum Corrosivum (Hydrarg. Chlor. Corr.), Corrosive Mercuric Chloride, U. S. P. (Bichloride of Mercury, Corrosive Sublimate, Mercuric Chloride, Perchloride of Mercury).—HgCl.**

Heavy, colorless crystals, crystalline masses or white powder, odorless and with a characteristic metallic taste. Slowly soluble in water (1:13.5) and freely soluble in alcohol (1:3.8). Its solubility in water is increased by the addition of sodium or ammonium chloride. Incompatible with soluble carbonates and hydroxides, with iodides and with alkaloids and other organic compounds. Reduced to calomel or metallic mercury by iron, zinc and other reducing agents.

ACTION AND USES: Chiefly as a germicide and antiseptic; also as a specific antisyphilitic agent. Acute mercuric chloride poisoning is treated by raw eggs and lavage.

DOSAGE: 0.003 Gm. or $\frac{1}{20}$ grain (U. S. P.) in solution or pills. For hypodermic use in syphilis 0.01 Gm. or $\frac{1}{6}$ grain daily (produces marked pain and irritation). As antiseptic externally in solution from 1:20,000 to 1:2,000. As disinfectant of clothing, in 1:1,000 solution.

Mulla Hydrargyri Chloridi Corrosivi (Mull. Hydrarg. Chlor. Corros.), *Corrosive Mercuric Chloride* Mull, N. F. (Unguentum Hydrargyri Chloridi Corrosivi Extensum, N. F. III).—Corrosive mercuric chloride (0.2%), alcohol, benzoinated suet and benzoinated lard.

Toxibellae Hydrargyri Chloridi Corrosivi (Toxibell. Hydrarg. Chlor. Corr.), *Poison Tablets of Corrosive Mercuric Chloride*, U. S. P. (Corrosive Sublimate Tablets, Bichloride Tablets).—Tablets of an angular shape colored blue, each having the word "Poison" and the skull and cross bones design distinctly stamped on it. Each 1 Gm. tablet contains about 0.5 Gm. or 8 grains corrosive mercuric chloride and 0.5 Gm. or 8 grains sodium chloride.

**Hydrargyri Chloridum Mite* (Hydrarg. Chlor. Mit.), *Mild Mercurous Chloride*, U. S. P. (Mercurous Chloride, Calomel, Protochloride of Mercury, Subchloride of Mercury).— HgCl .

White or nearly white, odorless, tasteless powder. Insoluble in water, alcohol or ether. Incompatible with alkalies, with oxidizing acids like nitric acid, and with soluble bromides and iodides.

ACTION AND USES: A slow cathartic, fairly powerful and relatively non-irritant. Should usually be followed by a saline laxative. Should not be taken continuously as it may salivate. Also employed as a diuretic in cardiac dropsies. Intramuscular injections are very efficiently antisyphilitic, but painful.

DOSAGE: Laxative, 0.15 Gm. or $2\frac{1}{2}$ grains (U. S. P.); alternative, 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.). From 0.005 to 0.02 Gm., or from $\frac{1}{42}$ to $\frac{1}{3}$ grain, may be given every half hour or hour until from 0.1 to 0.2 Gm., or from $1\frac{1}{2}$ to 3 grains have been given. In the treatment of syphilis 0.1 Gm. or $\frac{1}{2}$ grain in oily suspension is injected about once a week.

Lotio Nigra (Lot. Nigr.), *Black Lotion*, N. F. (Black Wash, Aqua Phagedaenica Nigra).—A suspension of mercurous oxide, Hg_2O , produced by the action of lime water on mild mercurous chloride.

USES: Mercurial antiseptic.

Pulvis Hydrargyri Chloridi Mitis et Jalapae (*Pulv. Hydrarg. Chlor. Mit. et Jalap.*), Powder of Mild Mercurous Chloride and Jalap, N. F. (Calomel and Jalap).—Mild mercurous chloride (34%) and jalap (66%).

USES: Drastic purgative.

DOSAGE: 0.65 Gm. or 10 grains (N. F.).

***Hydrargyri Iodidum Flavum** (*Hydrarg. Iod. Flav.*), Yellow Mercurous Iodide, U. S. P. (Mercurous Iodide, Protoiodide of Mercury, Yellow Iodide of Mercury).— HgI .

Bright yellow, odorless, tasteless powder. Almost insoluble in water and wholly insoluble in alcohol or ether. Decomposed by oxidizing agents, alkalis and the haloid salts of the alkali metals (chlorides, bromides or iodides).

ACTION AND USES: Used principally as an antisymphilitic; generally given in the form of pills. Like all ingestion methods of treating syphilis, much less reliable than the intramuscular administration of mercurials and is falling into disuse on account of relative inefficiency. Has proved of value in lichen ruber planus.

DOSAGE: 0.01 Gm. or $\frac{1}{6}$ grain (U. S. P.).

***Hydrargyri Iodidum Rubrum** (*Hydrarg. Iod. Rub.*), Red Mercuric Iodide, U. S. P. (Biniodide of Mercury, Mercuric Iodide, Red Iodide of Mercury).— HgI_2 .

Scarlet-red, odorless, nearly tasteless powder. Nearly insoluble in water and slightly soluble in alcohol (1:115); soluble in solutions of soluble iodides, sodium thiosulphate and hot solutions of alkali chlorides.

ACTION AND USES: Mercuric iodide, dissolved with potassium or sodium iodide, is used as an antiseptic, germicide and antisymphilitic. It is practically as effective as mercuric chloride and less irritant or corrosive.

DOSAGE: 0.003 Gm. or $\frac{1}{20}$ grain (U. S. P.).

Liquor Hydrargyri et Potassii Iodidi (*Liq. Hydrarg. et Pot. Iod.*), Solution of Mercury and Potassium Iodide, N. F. (Solution of Potassium Iodohydrargyrate, Channing's Solution).—Red mercuric iodide (1%), potassium iodide (0.8%), and water.

DOSAGE: 0.2 Cc. or 3 minims (N. F.).

Hydrargyri Nitras, Mercuric Nitrate.— $\text{Hg}(\text{NO}_3)_2$.

Liquor Hydrargyri Nitratis (*Liq. Hydrarg. Nit.*), Solution of Mercuric Nitrate, N. F.— $\text{Hg}(\text{NO}_3)_2$ (60%).

USES: Caustic.

Unguentum Hydrargyri Nitratis (Ung. Hydrarg. Nit.), *Ointment of Mercuric Nitrate*, U. S. P. (Citrine Ointment).—Mercuric nitrate (equivalent to mercury 7%) in lard.

***Hydrargyri Oxidum Flavum** (Hydrarg. Oxid. Flav.), **Yellow Mercuric Oxide**, U. S. P.—HgO.

Light orange-yellow, heavy, odorless powder, with a somewhat metallic taste. Almost insoluble in water, insoluble in alcohol, readily dissolved by dilute hydrochloric acid and nitric acid.

ACTION AND USES: The ointment is used, especially on the eye, as an antiseptic and stimulant; also to some extent in impetiginous affections.

DOSAGE: For the eye, 2 per cent.; for the skin, from 5 to 10 per cent.

Lotio Flava (Lot. Flav.), *Yellow Lotion*, N. F. (Yellow Wash, *Aqua Phagedaenica Flava*).—A suspension of mercuric oxide produced by the action of lime water on mercuric chloride.

USES: Mercurial antiseptic.

***Unguentum Hydrargyri Oxidi Flavi** (Ung. Hydrarg. Oxid. Flav.), *Ointment of Yellow Mercuric Oxide*, U. S. P.—Yellow mercuric oxide (10%) in water, hydrous wool fat and petrolatum.

Hydrargyri Oxidum Rubrum (Hydrarg. Oxid. Rub.), **Red Mercuric Oxide**, U. S. P. (Red Precipitate).—HgO.

Heavy, orange-red scales or powder, odorless and with a somewhat metallic taste. Almost insoluble in water; insoluble in alcohol, readily dissolved by nitric acid.

ACTION AND USES: Occasionally used externally in the form of an ointment and as a parasiticide. Without advantage over yellow oxide.

Unguentum Hydrargyri Oxidi Rubri (Ung. Hydrarg. Oxid. Rub.), *Ointment of Red Mercuric Oxide*, N. F.—Red mercuric oxide (10%) in water, hydrous wool fat and petrolatum.

***Hydrargyri Salicylas** (Hydrarg. Salicyl.), **Mercuric Salicylate**, U. S. P. (Mercuric Subsalicylate).—Contains Hg (about 57%).

White or nearly white, odorless, tasteless powder. Nearly insoluble in water or alcohol. Dissolved by solutions of the fixed alkalis or their carbonates.

ACTION AND USES: Used as antisyphilitic especially by intramuscular injection of oily suspensions; also as local antiseptic.

DOSAGE: 0.004 Gm. or $\frac{1}{45}$ grain (U. S. P.). By intramuscular injection, 0.6 Cc. or 10 minims of a 10 per cent. suspension in liquid petrolatum once in four days.

***Hydrargyrum (Hydrarg.), Mercury, U. S. P. (Quicksilver).—Hg.**

Shining, silver-white, odorless, tasteless metal, liquid at ordinary temperatures. Insoluble in the ordinary solvents.

ACTION AND USES: Its preparations are used to secure the systemic effects of mercury and locally against parasites.

***Hydrargyrum cum Creta (Hydrarg. cum Cret.), Mercury with Chalk, U. S. P. (Gray Powder).—Mercury (38%), with prepared chalk (57%), honey and water.**

Light gray, odorless, rather damp powder, with a slightly sweet taste.

ACTION AND USES: A mild cathartic acting like calomel. Also used in the treatment of infantile and hereditary syphilis and of syphilis in aged patients.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.), as powder.

***Massa Hydrargyri (Mass. Hydrarg.), Mass of Mercury, U. S. P. (Blue Mass, Blue Pill).—Hg (about 33%) in pill mass.**

USES: Mainly as cathartic.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Oleatum Hydrargyri (Oleat. Hydrarg.), Oleate of Mercury, U. S. P.—A mixture of mercury oleate and oleic acid equivalent to 25 per cent. of HgO.

USES: Alterative, antisyphilitic; rarely used.

Petroxolinum Hydrargyri (Petrox. Hydrarg.), Mercury Petroxolin, N. F. (Mercury Petrox).—Mercury (30%) in hydrous wool fat, oleic acid and solid petroxolin.

***Unguentum Hydrargyri (Ung. Hydrarg.), Mercurial Ointment, U. S. P.—Metallic mercury (about 50%) and oleate of mercury (2%) with prepared suet and benzoinated lard.**

USES: Especially to secure the systemic effects of mercury by inunction.

***Unguentum Hydrargyri Dilutum (Ung. Hydrarg. Dil.), Diluted Mercurial Ointment, U. S. P. (Blue Ointment, Hydrargyri Unguentum P. I.).—Mercurial ointment (60%) equivalent to about 30 per cent. of metallic mercury, with petrolatum.**

USES: Especially in pediculosis.

***Hydrargyrum Ammoniatum (Hydrarg. Ammon.), Ammoniated Mercury, U. S. P.** (White Precipitate).—Mercurammonium chlorid, HgNH_2Cl , corresponding to about 79 per cent. of Hg.

White, odorless lumps or powder, with an earthy, afterward styptic and metallic taste. Insoluble in water or alcohol.

ACTION AND USES: The ointment is extensively used as a cutaneous antiseptic especially for impetigo contagiosa and other pustular dermatoses and ringworm of the scalp. Also used for removing scales of psoriasis. It may cause salivation.

DOSAGE: From 2 to 10 per cent. ointment.

***Unguentum Hydrargyri Ammoniatum (Ung. Hydrarg. Ammon.), Ointment of Ammoniated Mercury, U. S. P.** (White Precipitate Ointment).—Ammoniated mercury (10%) in white petrolatum and hydrous wool fat.

Hydrastina (Hydrastina), Hydrastine, U. S. P.—An alkaloid obtained from hydrastis or prepared synthetically.

White crystals or powder. Almost insoluble in water, soluble in alcohol (1:170).

ACTION AND USES: Used in the treatment of urethral and vesical catarrh and as a systemic hemostatic, especially in excessive menstruation. Its efficiency and the mechanism of its action are doubtful.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.).

Hydrastinae Hydrochloridum (Hydrastin. Hydrochl.), Hydrastine Hydrochloride, U. S. P. (Hydrastine Chloride).

White, odorless, bitter powder. Very soluble in water and in alcohol.

ACTION AND USES: Those of hydrastine.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.).

Liquor Hydrastinae Compositus (Liq. Hydrastin. Co.), Compound Solution of Hydrastine, N. F. (Colorless Hydrastine Solution).—Hydrastine hydrochloride (0.3%), aluminum chloride, calcium chloride, and magnesium chloride (each 0.3%), potassium chloride (0.1%) in glycerin and water.

USES: Complex and irrational astringent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Hydrastininae Hydrochloridum (Hydrastinin. Hydrochl.), Hydrastinine Hydrochloride, U. S. P. (Hydrastinine

Chloride).—The hydrochloride of hydrastinine, an alkaloid obtained synthetically from hydrastine.

Yellowish, odorless crystals or powder. Soluble in chloroform (1:195); very soluble in water and alcohol.

ACTION AND USES: Hydrastinine resembles hydrastine in its action but is said to be less depressing. Hydrastinine has a pronounced pressor effect on peripheral vessels, causing a rise of blood pressure.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

***Hydrastis (Hydrastis), Hydrastis, U. S. P. (Golden Seal).**

—Rhizome and roots, yielding not less than 2.5 per cent. of ether-soluble alkaloids, chiefly hydrastin.

ACTION AND USES: Hydrastis is an unpleasant bitter; it possesses no advantages over other simple bitters (gentian). It has been asserted to have a special influence on mucous membranes favoring their nutrition and return to a normal condition. There is little or no experimental evidence to bear out the claim for any special action.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Extractum Hydrastis (Ext. Hydrast.), Extract of Hydrastis, U. S. P. (Extract of Golden Seal, Powdered Extract of Hydrastis).—One Gm. extract represents about 4 Gm. hydrastis. Yields about 10 per cent. of alkaloids.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

***Fluidextractum Hydrastis (Fldext. Hydrast.), Fluidextract of Hydrastis, U. S. P.**—Hydrastis (100%), yielding about 2 per cent. of alkaloids. Absolute alcohol content about 55 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Glyceritum Hydrastis (Glycer. Hydrast.), Glycerite of Hydrastis, U. S. P. (Glycerite of Golden Seal).—Hydrastis (100%) yielding not less than 1.245 per cent. of alkaloids.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Hydrastis (Tr. Hydrast.), Tincture of Hydrastis, U. S. P. (Tincture of Golden Seal).—Hydrastis (20%), yielding about 0.4 per cent. of hydrastin; in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

***Hydrogenii Dioxidum, Hydrogen Dioxide (Hydrogen Peroxide).— H_2O_2 .**

**Liquor Hydrogenii Dioxidi (Liq. Hydrog. Diox.), Solution of Hydrogen Dioxide, U. S. P. (Aqua Hydrogenii Dioxidii, U. S. P. VIII, Solution of Hydrogen Peroxide).*
— H_2O_2 (not less than 3%).

ACTION AND USES: Nontoxic and fairly efficient local antiseptic and detergent, useful mainly for the destruction of pus. Used on suppurating wounds and as a mouthwash and gargle; also as a bleaching agent. Should not be injected into a wound unless free drainage is present.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.). For external application should be diluted about 1:4.

Hyoscyaminae Hydrobromidum (Hyoscyamin. Hydrobr.), Hyoscyamine Hydrobromide, U. S. P. (Hyoscyamine Bromide).—Obtained from hyoscyamus and other plants of the nightshade family.

White, odorless crystals. Freely soluble in alcohol (1:2.5) and in chloroform (1:1.7); very soluble in water.

ACTION AND USES: Similar to those of atropine; is usually more sedative to the brain, but at times produces acute delirium. The drug is very uncertain.

DOSAGE: 0.3 mg. or $\frac{1}{200}$ grain (U. S. P.).

**Hyoscyamus (Hyosc.), Hyoscyamus, U. S. P. (Henbane, Hyoscyami Folium, P. I.).*—Leaves and tops, yielding not less than 0.065 per cent. of alkaloids.

ACTION AND USES: Similar to those of belladonna.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Extractum Hyoscyami (Ext. Hyosc.), Extract of Hyoscyamus, U. S. P.—One Gm. extract represents about 4 Gm. hyoscyamus. Yields about 0.25 per cent. of alkaloids.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Fluidextractum Hyoscyami (Fldext. Hyosc.), Fluidextract of Hyoscyamus, U. S. P. (Fluidextract of Henbane).—Hyoscyamus (100%) yielding about 0.065 per cent. of alkaloids, chiefly hyoscyamine. Absolute alcohol content about 60 per cent.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Oleum Hyoscyami Compositum (Ol. Hyosc. Co.), Compound Oil of Hyoscyamus, N. F. (Balsamum Tranquillans).—Infused oil of hyoscyamus (hyoscyamus 10%) flavored with oils of lavender, peppermint, rosemary and thyme.

**Tinctura Hyoscyami (Tr. Hyosc.), Tincture of Hyoscyamus, U. S. P. (Tincture of Henbane, Hyoscyami Tinctura, P. I.).*—Hyoscyamus (10%), yielding about 0.0065 per

cent. of hyoscyamus alkaloids; in diluted alcohol. Absolute alcohol content about 47 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

***Hypophysis Sicca (Hypophysis Sic.), Desiccated Hypophysis, U. S. P.** (Desiccated Pituitary Body).—The posterior lobe obtained from the pituitary body of cattle.

A yellowish or grayish powder with a characteristic odor. Partially soluble in water.

ACTION AND USES: Solutions are used, hypodermically, to strengthen uterine contractions in prolonged labor. They also stimulate peristalsis and have been used with some success after abdominal operations and as a vasomotor stimulant. Oral administration is ineffective.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

***Liquor Hypophysis (Liq. Hypophysis), Solution of Hypophysis, U. S. P.** (Solution of the Pituitary Body).—

Contains the water-soluble principles from the fresh posterior lobe of the pituitary body of cattle. Assayed by testing against a standard solution of beta-iminazolyethylamine hydrochloride by the method of the U. S. Hygienic Laboratory.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Ignatia (Ignat.), Ignatia, N. F. (Saint Ignatius Bean, Ignatia Amara).—Seeds of *Strychnos Ignatii*.

ACTION AND USES: Constituents and actions similar to those of nux vomica, over which ignatia has no advantage, being less reliable.

DOSAGE: 0.06 Gm. or 1 grain (N. F.).

Extractum Ignatiæ (Ext. Ignat.), Extract of Ignatia, N. F. (Powdered Extract of Ignatia).—A hydroalcoholic extract, yielding about 6 per cent. of alkaloids.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (N. F.).

Tinctura Ignatiæ (Tr. Ignat.), Tincture of Ignatia, N. F.—Ignatia (10%), yielding about 0.2 per cent. of ignatia alkaloids; in alcohol and water. Absolute alcohol content about 85 per cent.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

INFUSA, INFUSIONS.—Aqueous preparations made by pouring hot or cold water over a vegetable drug, allowing the mixture to stand for a definite period and then straining. For the official infusions, see under the names of the respective drugs.

Inula (Inula), Inula, N. F. (Elecampane).—Rhizome and roots.

ACTION AND USES: At one time used in bronchitis. No scientific indications for its use.

DOSAGE: 2 Gm. or 30 grains (N. F.).

INUNCTA, INUNCTIONS.—Semisolid ointment-like preparations used for external application. For the various inunctions see under the names of the principal ingredients.

***Iodoformum (Iodof.), Iodoform, U. S. P.**—Triiodomethane, CHI_3 .

Lemon-yellow powder or lustrous crystals, with a peculiar, penetrating odor and an unpleasant, slightly sweetish taste suggestive of iodine. Nearly insoluble in water, to which, however, it imparts its odor and taste; soluble in alcohol (1:60) and in glycerin (1:80).

ACTION AND USES: Used as dusting powder, promoting healing by granulation. Oily suspensions are also used in treatment of tuberculous fistulas.

Collodium Iodoformi (Collod. Iodof.), Iodoform Collodion, N. F.—Iodoform (5%) in flexible collodion.

USES: There appears to be no advantage in applying iodoform in collodion.

Glycerogelatinum Iodoformi (Glycerogel. Iodof.), Iodoform Glycerogelatin, N. F.—Iodoform (10%), water, glycerin and glycerinated gelatin.

Iodoformum Aromatisatum (Iodof. Arom.), Aromatized Iodoform, N. F.—Coumarin (4%) and iodoform.

Petroxolinum Iodoformi (Petrox. Iodof.), Iodoform Petroxolin, N. F. (Iodoform Petrox.).—An unstable preparation made from iodoform (3%), acetone, oleic acid, eucalyptol and liquid petroxolin.

Unguentum Iodoformi (Ung. Iodof.), Iodoform Ointment, U. S. P.—Iodoform (10%) in benzoinated lard.

***Iodum (Iodum), Iodine, U. S. P.—I.**

Heavy, bluish-black, brittle plates, with a metallic luster, a distinctive odor and a sharp, acrid taste. Very slightly soluble in water (1:2950) and soluble in alcohol (1:12.5). Freely soluble in solutions of alkali iodides.

ACTION AND USES: The tincture and compound solution are used externally as local irritant and antiseptic. The irritant action can be easily graduated by successive application. It is rarely used internally for the systemic effects of iodides.

DOSAGE: 0.005 Gm. or $\frac{1}{2}$ grain (U. S. P.), largely diluted.

Collodium Iodi (*Collod. Iodi*), *Iodine Collodion*, N. F.—Iodine (5%) in flexible collodion.

USES: There appears little advantage in applying iodine in collodion.

Liquor Iodi Compositus (*Liq. Iodi Co.*), *Compound Solution of Iodine*, U. S. P. (Lugol's Solution).—Iodine (5%) and potassium iodide (10%) in water.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Liquor Iodi Phenolatus (*Liq. Iod. Phenol.*), *Phenolated Solution of Iodine*, N. F. (*Liquor Iodi Carbolatus*, N. F. III, Carbolized Solution of Iodine, Boulton's Solution, French Mixture).—Compound solution of iodine (1.5%), liquefied phenol (0.6%), glycerin and water. The iodine combines with the phenol.

USES: Antiseptic.

Petroxolinum Iodi (*Petrox. Iod.*), *Iodine Petroxolin*, N. F. (Iodine Petrox 10%).—An unstable preparation, containing iodine, ammonium iodide and other iodine compounds in variable proportions, with a mixture of liquid petrolatum, alcohol, ammonium soap and oil of lavender; prepared from iodine (10%) and liquid petroxolin.

Petroxolinum Iodi Dilutum (*Petrox. Iod. Dil.*), *Diluted Iodine Petroxolin*, N. F. (Iodine Petrox 5%).—An unstable preparation containing iodine, ammonium iodide and other iodine compounds in variable proportions with a mixture of liquid petrolatum, alcohol, ammonium soap and oil of lavender; made from iodine (5%) and liquid petroxolin. Absolute alcohol content about 19 per cent.

Syrupus Iodotannicus (*Syr. Iodotan.*), *Syrup of Iodotannin*, N. F.—Made by combining iodine (0.27%) with tannic acid (0.54%) in syrup. Contains iodine probably in the form of an iodide.

USES: Similar to those of potassium iodide; without advantage over the latter.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Tinctura Iodi* (*Tr. Iodi*), *Tincture of Iodine*, U. S. P.—An alcoholic solution of iodine (about 7%) and potassium iodide (about 5%). Absolute alcohol content about 83 per cent.

DOSAGE: 0.1 Cc. or $1\frac{1}{2}$ minims (U. S. P.).

Tinctura Iodi Decolorata (*Tr. Iod. Decolor.*), *Decolorized Tincture of Iodine*, N. F.—A solution of sodium iodide

and ammonium iodide obtained by mixing iodine and sodium thiosulphate (each 8.3%), stronger ammonia water (6.5%) and alcohol. Absolute alcohol content about 75 per cent. Does not contain free iodine.

USES: For external use; of doubtful utility.

Tinctura Iodi Fortior (*Tr. Iod. Fort.*), *Stronger Tincture of Iodine*, N. F. (Churchill's Tincture of Iodine).—Iodine (16.5%) and potassium iodide (3.3%) in water and alcohol. Absolute alcohol content about 65 per cent.

Unguentum Iodi (*Ung. Iodi*), *Iodine Ointment*, U. S. P.—Iodine (4%) and potassium iodide (4%) in glycerin and benzoinated lard. Should be made as required.

**Ipecacuanha* (*Ipecac.*), *Ipecac*, U. S. P. (*Ipecacuanhae Radix*, P. I.).—A root, yielding not less than 1.75 per cent. of alkaloids.

ACTION AND USES: Used as irritant, nauseant, expectorant and emetic. Also specific against amebic dysentery, but in this use has been practically displaced by emetine.

DOSAGE: Emetic, 1 Gm. or 15 grains (U. S. P.). Expectorant, 0.06 Gm. or 1 grain.

**Fluidextractum Ipecacuanhae* (*Fldext. Ipecac.*), *Fluidextract of Ipecac*, U. S. P.—*Ipecac* (100%), yielding about 2 per cent. of alkaloids. Absolute alcohol content about 30 per cent.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.).

Pulvis Ipecacuanhae et Opii.—See under *Opium*.

**Syrupus Ipecacuanhae* (*Syr. Ipecac.*), *Syrup of Ipecac*, U. S. P.—Fluidextract of *ipeacac* (7%) and acetic acid (1%) in glycerin and syrup.

DOSAGE: Expectorant, 1 Cc. or 15 minims (U. S. P.); emetic, 15 Cc. or 4 fluidrachms (U. S. P.).

Syrupus Ipecacuanhae et Opii and *Tinctura Ipecacuanhae et Opii*.—See under *Opium*.

Vinum Ipecacuanhae (*Vin. Ipecac.*), *Wine of Ipecac*, N. F.—Fluidextract of *ipeacac* (10%, yielding about 0.2% *ipeacac* alkaloids) in alcohol and sherry wine. Absolute alcohol content about 27 per cent.

DOSAGE: 1 Cc. or 15 minims.

Iris (*Iris*), *Orris*, N. F. (*Orris Root*).—Rhizome.

ACTION AND USES: At one time thought to be a mild digestive stimulant, slightly cathartic and diuretic. Now employed as dusting powder, tooth powder and perfume.

Iris Versicolor (Iris Vers.), Blue Flag, N. F.—Rhizome and roots.

ACTION AND USES: Obsolete cathartic; without advantage over resin of podophyllum.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Iridis Versicoloris (Fldext. Iri. Ver.), Fluidextract of Blue Flag, N. F.—Blue flag (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Jalapa (Jalap.), Jalap, U. S. P.**—A root, yielding not less than 7 per cent. of resins.

ACTION AND USES: Drastic hydragogue cathartic, used especially in dropsies. Preferred to other resinous cathartics, as being rather less irritant.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Extractum Jalapae (Ext. Jalap.), Extract of Jalap, N. F.—An alcoholic extract.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Jalapae (Fldext. Jalap.), Fluidextract of Jalap, N. F.—Jalap (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Pulvis Jalapae Compositus (Pulv. Jalap. Co.), Compound Powder of Jalap, U. S. P.** (Pulvis Purgans).—Jalap (35%) and potassium bitartrate (65%).

USES: Useful hydragogue cathartic.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Resina Jalapae (Res. Jalap.), Resin of Jalap, U. S. P.—The resin from jalap.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Tinctura Jalapae (Tr. Jalap.), Tincture of Jalap, N. F.—Jalap (20%), in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Tinctura Jalapae Composita (Tr. Jalap. Co.), Compound Tincture of Jalap, N. F.—Jalap (12.5%) and resin of scammony (3%), in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Juglans (Juglan.), Juglans, N. F. (Butternut Bark, White Walnut Bark).—Inner bark of the roots.

ACTION AND USES: Obsolete cathartic, without advantage over cascara.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Juglandis (Fldext. Jugland.), *Fluidextract of Juglans*, N. F.—Juglans (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Juniperus (Junip.), Juniper Berries, N. F.—Ripe fruit.

ACTION AND USES: Irritant to urinary organs, owing to the chief constituent, a volatile oil. The spirit is sometimes used as an irritant diuretic.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Juniperi (Fldext. Junip.), *Fluidextract of Juniper Berries*, N. F.—Juniper (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Kaolinum (Kaolinum), Kaolin, N. F.—Purified native aluminum silicate.

White or nearly white powder or lumps with an earthy taste. Insoluble in water.

ACTION AND USES: Used in a poultice, cataplasma kaolini, similar to proprietary preparations. Internally, as absorbent in diarrhea and dysentery.

DOSAGE: Tablespoonful dry or in tea several times daily.

Cataplasma Kaolini (Catapl. Kaolin.), *Cataplasma of Kaolin*, N. F.—A mixture of kaolin and glycerin with boric acid aromatized with thymol, methyl salicylate and oil of peppermint.

USES: Not superior to flaxseed or bread-and-milk poultices.

Kava (Kava), Kava, N. F. (Methysticum, Kava Kava).—Rhizome and roots of *Piper methysticum*.

ACTION AND USES: Mild irritant, without advantage over cubeb. Often inert. Owes its properties to an oleoresin.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Kavae (Fldext. Kav.), *Fluidextract of Kava*, N. F.—Kava (100%). Absolute alcohol content about 50 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Kino (Kino), Kino, U. S. P.—A dried plant juice.

Dark brown fragments or brick-red powder, odorless and having an astringent taste. Partly soluble in water.

ACTION AND USES: Used as intestinal and pharyngeal astringent; has no special advantage over other vegetable astringents, such as krameria.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Pulvis Kino et Opii Compositus (*Pulv. Kino et Opii Co.*), *Compound Powder of Kino and Opium*, N. F. (*Pulvis Kino Compositus*, N. F. III).—Kino (75%), and powdered opium (5%) with Saigon cinnamon.

USES: Antidiarrheic.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Tinctura Kino (*Tr. Kino*), *Tincture of Kino*, U. S. P.—Kino (10%) in diluted alcohol. Absolute alcohol content about 49 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Tinctura Kino et Opii Composita (*Tr. Kino et Opii Co.*), *Compound Tincture of Kino and Opium*, N. F. (*Tinctura Kino Composita*, N. F. III, *Compound Tincture of Kino*).—Kino (2%), opium (1%), spirit of camphor (6.5%), oil of clove, cochineal and aromatic spirit of ammonia in diluted alcohol. Absolute alcohol content about 50 per cent.

USES: Antidiarrheic; undesirably complex.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Kola (Kola), Kola, N. F. (*Cola, Kola Nuts, Cola Nuts*).—The peeled seeds.

ACTION AND USES: Possesses stimulant actions of its caffeine and theobromin. Without advantage over coffee.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Kolae (*Fldext. Kol.*), *Fluidextract of Kola*, N. F. (*Fluidextractum Sterculiae*, N. F. III).—Kola (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Krameria (Kramer.), Krameria, N. F. (*Rhatany*).—Root.

ACTION AND USES: Powerful astringent; like other tannin-containing drugs, is rapidly passing out of use.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Extractum Krameriae (*Ext. Kramer.*), *Extract of Krameria*, N. F. (*Powdered Extract of Krameria*).—An aqueous extract diluted with starch, 1 Gm. representing 4 Gm. krameria.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Fluidextractum Krameriae (*Fldext. Kramer.*), *Fluidextract of Krameria*, N. F.—*Krameria* (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Fluidglyceratum Krameriae (*Fldglycer. Kramer.*), *Fluidglycerate of Krameria*, N. F.—*Krameria* (100%) in glycerin and water.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Syrupus Krameriae (*Syr. Kramer.*), *Syrup of Krameria*, N. F.—*Fluidextract of krameria* (45%) in syrup. Absolute alcohol content about 18 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Tinctura Krameriae (*Tr. Kramer.*), *Tincture of Krameria*, N. F.—*Krameria* (20%) in diluted alcohol. Absolute alcohol content about 45 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Lac Vaccinum (*Lac Vaccinum*), *Cow's Milk*, N. F.

Lac Fermentatum (*Lac Ferment.*), *Fermented Milk*, N. F. (*Kumyss*).—A fermentation product obtained by the action of yeast on cow's milk to which sugar (sucrose) has been added. The following is an unofficial formula:

Take a quart of skim milk, one-fifth of a cake of yeast and two tablespoonfuls of sugar. Heat the milk. Dissolve the yeast in a little water and mix it with the sugar and lukewarm milk. Pour the mixture into strong bottles, stopper them tightly with new corks, and tie down the corks with strong twine. Shake the bottles well and place in a refrigerator to ferment slowly. After three days lay the bottles on their sides, turning them occasionally. Five days are required to complete the fermentation.

DOSAGE: 250 Cc. or 8 fluidounces (N. F.).

Lactucarium (*Lactucar.*), *Lactucarium*, U. S. P.—The dried juice of a species of wild lettuce.

ACTION AND USES: Inefficient and obsolete hypnotic and respiratory sedative. The proprietary preparation to which its popularity was due was found to contain morphine. *Lactucarium* is now seldom used.

DOSAGE: 1 Gm. or 15 grains (U. S. P.). Not given as such.

Syrupus Lactucarii (*Syr. Lactucar.*), *Syrup of Lactucarium*, U. S. P.—*Tincture of lactucarium* (10%) and citric acid, flavored with orange flower water, in glycerin and syrup.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.). Absolute alcohol content about 5 per cent.

Tinctura Lactucarii (*Tr. Lactucar.*), *Tincture of Lactucarium*, U. S. P.—Lactucarium (50%) in glycerin, alcohol and water. Absolute alcohol content about 46 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Lappa (Lappa), Lappa, N. F. (Burdock Root).

ACTION AND USES: Obsolete "alterative," used in nostrums.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Lappae (*Fldext. Lapp.*), *Fluidextract of Lappa*, N. F.—Lappa (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Leptandra (Leptand.), Leptandra, N. F. (Culver's Root).—Rhizome and roots.

ACTION AND USES: Unreliable cathartic; without advantage over resin of podophyllum.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Extractum Leptandrae (*Ext. Leptand.*), *Extract of Leptandra*, N. F. (Powdered Extract of Leptandra).—A hydroalcoholic extract, diluted with starch. One Gm. represents 4 Gm. leptandra.

DOSAGE: 0.25 Gm. or 4 grains (N. F.).

Fluidextractum Leptandrae (*Fldext. Leptand.*), *Fluidextract of Leptandra*, N. F.—Leptandra (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Limonis Cortex (Limon. Cort.), Lemon Peel, U. S. P.

ACTION AND USES: Used for the preparation of lemon flavors.

Tinctura Limonis Corticis (*Tr. Limon. Cort.*), *Tincture of Lemon Peel*, U. S. P.—Fresh lemon peel (50%) treated with alcohol. Absolute alcohol content about 83 per cent.

NIMENTA, LINIMENTS.—Liquid preparations intended for external application. For the various official liniments, under the names of the respective chief ingredients.

(*Linum*), *Linseed*, U. S. P. (Flaxseed).—Used in *ulcent* infusions, also when crushed or ground in

poultices. If employed where the skin is broken the entire poultice should be sterilized by boiling before it is applied, in order to avoid infection.

LIQUORES, SOLUTIONS.—Aqueous liquid preparations in which one or more substances are completely dissolved.

For the various official solutions, see under the names of the respective principal ingredients. For *Liquor Acidi Arsenosi*, see under *Arseni Trioxidum*; for *Liquor Antisepticus*, under *Acidum Boricum*; for *Liquor Antisepticus Alkalinus*, under *Potassii Bicarbonas*; for *Liquor Arseni et Hydrargyri Iodidi*, under *Arseni Iodidum*; for *Liquor Arsenicalis Clemens*, under *Potassii Arsenas*; for *Liquor Calcis Sulphuratae*, under *Calcii Sulphidum Crudum*; for *Liquor Ferri et Ammonii Acetas*, under *Ammonii Acetas*; for *Liquor Zinci et Alumini Compositus* and *Liquor Zinci et Ferri Compositus*, under *Zinci Sulphas*.

Lithii Bromidum (Lith. Brom.), Lithium Bromide, U. S. P.—
 LiBr (not less than 85%).

White, granular, odorless, very deliquescent salt with a sharp, slightly bitter taste. Very soluble in water (1:0.6) and soluble in alcohol and in ether.

ACTION AND USES: Used like other bromides, over which it has no advantage. For the supposed urate-solvent properties of lithium, see *Lithii Carbonas*.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Lithii Bromidi (Elix. Lith. Brom.), Elixir of Lithium Bromide, N. F.—Lithium bromide (8.5%) in syrup, water and aromatic elixir. Absolute alcohol content about 7 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Lithii Carbonas (Lith. Carb.), Lithium Carbonate, U. S. P.—
 Li_2CO_3 .

Light white, odorless powder with an alkaline taste. Soluble in water (1:78); almost insoluble in alcohol.

ACTION AND USES: Used in the same way as sodium carbonate; without advantage over the latter. Lithium was formerly given to promote the elimination of urates because it was supposed to have special urate-solvent properties, but under the conditions existing in the body these properties do not come into play.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Lithii Citras (Lith. Cit.), Lithium Citrate, U. S. P.

White, odorless powder or granules, with a cooling, faintly alkaline taste. Freely soluble in water (1:1.4), very slightly soluble in alcohol.

ACTION AND USES: Used like citrates of sodium or potassium, over which it has no advantage. For the supposed urate-solvent properties of lithium, see **Lithii Carbonas**.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Elixir Lithii Citratis (Elix. Lith. Cit.), Elixir of Lithium Citrate, N. F.—Lithium citrate (8.5%), in aromatic elixir. Absolute alcohol content about 21 per cent.

USES: An objectionable alcoholic preparation of lithium.

DOSAGE: 8 Cc. or 2 fluidrachms (U. S. P.).

Sal Lithii Citratis Effervescens (Sal. Lith. Cit. Eff.), Effervescent Salt of Lithium Citrate, N. F.—Lithium citrate (5%), sodium bicarbonate (57%), tartaric acid (30%) and citric acid, the three last-named yielding sodium tartrate and sodium citrate.

USES: Superfluous, and in view of its small lithium content, misnamed.

DOSAGE: 8 Gm. or 2 drachms (N. F.).

Lithii Salicylas (Lith. Salicyl.), Lithium Salicylate, N. F.

White or grayish white odorless powder, with a sweetish taste; deliquescent in a moist atmosphere. Very soluble in water or alcohol.

ACTION AND USES: As a salicylate, inferior to sodium salicylate. Adequate doses would produce the toxic effects of lithium. For the supposed urate-solvent effects of lithium salts, see **Lithii Carbonas**.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Elixir Lithii Salicylatis (Elix. Lith. Salicyl.), Elixir of Lithium Salicylate, N. F.—Lithium salicylate (8.5%) in aromatic elixir. Absolute alcohol content about 21 per cent.

USES: An objectionable alcoholic salicylic preparation.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Lobelia (Lobel.), Lobelia, U. S. P. (Indian Tobacco).

Leaves and flowering tops.

ACTION AND USES: Expectorant, nauseant and emetic, resembling nicotine. Has fallen into disuse because of dangerous effects.

DOSAGE: 0.15 Gm. or 2½ grains (U. S. P.).

Fluidextractum Lobeliae (Fldext. Lobel.), Fluidextract of Lobelia, U. S. P.—Lobelia (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 0.15 Cc. or 2½ minims (U. S. P.).

Tinctura Lobeliae (Tr. Lobel.), *Tincture of Lobelia*, U. S. P. (Lobeliae Tinctura, P. I.).—Lobelia (10%) in diluted alcohol. Absolute alcohol content about 46 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

LOTIONES, LOTIONS (Washes). — Liquid mixtures, usually of solid medicinal substances suspended in water for external use.

For the various official lotions see under the names of the respective principal ingredients. For Lotio Ammoniacalis Camphorata, see under *Ammonia*; for Lotio Flava, under *Hydrargyri Oxidum Flavum*; for Lotio Nigra, under *Hydrargyri Chloridum Mite*; for Lotio Plumbi et Opii, under *Plumbi Acetas*.

Lupulinum (Lupul.), *Lupulin*, N. F.—Glandular powder separated from hops.

ACTION AND USES: Bitter tonic without material advantage over gentian. Reputed hypnotic and anaphrodisiac effects probably attributable to the alcohol in which it is commonly dissolved.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Fluidextractum Lupulini (Fldext. Lupul.), *Fluidextract of Lupulin*, N. F.—Lupulin (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

Oleoresina Lupulini (Oleores. Lupul.), *Oleoresin of Lupulin*, N. F.—The ether extract of lupulin.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Lycopodium (Lycopod.), *Lycopodium*, U. S. P.—The spores of a moss.

Very inflammable.

ACTION AND USES: Used as an inert dusting powder for the skin, and as diluent for insufflations of boric acid, tannin, etc., for the throat, nose and ear.

Macis (Macis), *Mace*, N. F.—The arilode (inner covering) removed from nutmeg (myristica).

Yellowish or brownish, fragrant, branched and lobed bands, or orange-buff or orange-brown powder, with a warm, aromatic taste.

ACTION AND USES: Aromatic flavor and carminative without advantage over cardamom.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

MAGMATA, MAGMAS.—Suspensions of insoluble or nearly insoluble substances.

***Magnesii Carbonas** (Mag. Carb.), Magnesium Carbonate, U. S. P.—A mixture of magnesium carbonate and magnesium hydroxide.

Light, white, friable masses or bulky white powder, odorless and with a slight earthy taste. Practically insoluble in water; insoluble in alcohol.

ACTION AND USES: Used internally against gastric hyperacidity and as a mild laxative; externally as dusting powder. See also *Magma Magnesiae*, under *Magnesii Hydroxidum*.

DOSAGE: 3 Gm. or 45 grains (U. S. P.).

Mistura Magnesiae, Asafoetidae et Opii (Mist. Mag. Asafoet. et Opii), Mixture of Magnesia, Asafoetida and Opium, N. F. (*Mistura Magnesiae et Asafoetidae*, N. F. III, Dewees' Carminative).—Magnesium carbonate (5%), tincture of asafoetida (7.5%), tincture of opium (1%), and sugar in water. Absolute alcohol content about 7 per cent.

USES: Complex and irrational mixture intended for use in diarrhea.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Magnesii Chloridum (Magnes. Chlorid.), Magnesium Chloride, N. F.— $MgCl_2 + 6H_2O$.

Colorless transparent crystals or white translucent pieces; deliquescent in moist air. Very soluble in water (1:0.6) and soluble in alcohol.

ACTION AND USES: Similar to but without advantage over magnesium sulphate.

DOSAGE: 15 Gm. or 4 drachms (N. F.).

***Magnesii Citras**, Magnesium Citrate.

Liquor Magnesii Citratis (Liq. Mag. Cit.), Solution of Magnesium Citrate, U. S. P.—Magnesium citrate corresponding to not less than 1.5 per cent. of magnesium oxide.

USES: Efficient and not distasteful mild saline laxative.

DOSAGE: 350 Cc. or 12 fluidounces (1 bottle) (U. S. P.).

Magnesii Hydroxidum, Magnesium Hydroxide.

Magma Magnesiae (Magma Mag.), Magnesia Magma, U. S. P. (Milk of Magnesia).—A suspension of magnesium hydroxide in water [equivalent to about 7% $Mg(OH)_2$] forming a thick, white liquid.

USES: Widely used as mild alkaline laxative and tooth wash.

DOSAGE: 10 Cc. or $2\frac{1}{2}$ fluidrachms (U. S. P.).

***Magnesii Oxidum** (Mag. Oxid.), Magnesium Oxide, U. S. P. (Magnesia, Calcined Magnesia, Light Magnesia).— MgO (not less than 96%).

A white, bulky, odorless powder with an earthy but not saline taste. Almost insoluble in water; insoluble in alcohol.

ACTION AND USES: Useful antacid and laxative.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Magnesii Oxidum Ponderosum (Mag. Oxid. Pond.), Heavy Magnesium Oxide, U. S. P. (Heavy Magnesia).— MgO (96%).

White, dense powder.

ACTION AND USES: Used like magnesium oxide.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

***Magnesii Sulphas** (Mag. Sulph.), Magnesium Sulphate, U. S. P. (Epsom Salt).— $MgSO_4 + 7H_2O$.

Small, colorless, odorless crystals, with a cooling, saline, bitter taste. Freely soluble in water (1:1); almost insoluble in alcohol.

ACTION AND USES: One of the most active of the saline cathartics.

DOSAGE: 15 Gm. or 4 drachms (U. S. P.), in solution. A solution containing 1 Gm. of salt in each Cc. may be given, 5 Cc. every hour until a laxative effect is secured. This should usually be diluted with water.

Liquor Magnesii Sulphatis Effervescens (Liq. Mag. Sulph. Eff.), Effervescent Solution of Magnesium Sulphate, N. F.—Magnesium sulphate (about 7%) with syrup of citric acid, potassium citrate and water.

DOSAGE: The contents of a bottle, about 350 Cc. or nearly 12 fluidounces (N. F.).

Maltum (Maltum), Malt, U. S. P.—Malted barley, capable of converting not less than 5 times its weight of starch into sugars.

Yellowish grains with a characteristic odor and a sweet taste.

ACTION AND USES: Used to digest starch by its diastase.

Extractum Malti (Ext. Malt.), Extract of Malt, U. S. P.—Syrupy liquid.

DOSAGE: 15 Gm. or 4 drachms (U. S. P.).

Malvae Folia (Malv. Fol.), Mallow Leaves, N. F.

ACTION AND USES: Demulcent "herb tea," without definite indications for its use.

Mangani Citras Solubilis (Mangan. Cit. Sol.), Soluble Manganese Citrate, N. F. (Manganese and Sodium Citrate).

Yellowish or pinkish white powder or translucent scales, odorless and having a slightly bitter, astringent taste. Freely soluble in water (1:4) and nearly insoluble in alcohol.

ACTION AND USES: Manganese salts have been tried as hematinic tonics, but are of very doubtful value. The choice among the various salts is a matter of convenience. The soluble citrate is typical of the soluble manganese salts, the dioxide of the insoluble.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Mangani Dioxidum Praecipitatum (Mangan. Diox. Praec.), Precipitated Manganese Dioxide, U. S. P.—Chiefly MnO_2 .

Heavy, odorless, tasteless black powder. Insoluble in water or alcohol.

ACTION AND USES: See *Mangani Citras Solubilis*.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Mangani Glycerophosphas Solubilis (Mangan. Glycerophos. Sol.), Soluble Manganese Glycerophosphate, N. F. (Soluble Manganous Glycerinophosphate).—A mixture of manganese glycérophosphate and citric acid.

Yellowish or pinkish white, odorless powder, with an acid taste. Freely soluble in water (1:4); alcohol dissolves the citric acid and leaves a residue which is nearly insoluble in water.

ACTION AND USES: As a manganese compound it has no material advantage over other manganese salts. For a discussion of glycerophosphates, see *Sodii Glycerophosphas*.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Mangani Hypophosphis (Mangan. Hypophos.), Manganese Hypophosphite, N. F.— $Mn(H_2PO_2)_2 + H_2O$.

Pink, odorless, nearly tasteless powder. Freely soluble in water and insoluble in alcohol.

ACTION AND USES: As a manganese compound, has no material advantage over other manganese salts. For a discussion of hypophosphites, see *Calcii Hypophosphis*.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Mangani Sulphas (Mangan. Sulph.), Manganese Sulphate, N. F.— MnSO_4 (about 65%).

Pale rose-colored, translucent, odorless crystals, with a slightly bitter taste, slightly efflorescent in moist air. Freely soluble in water (1:1) and insoluble in alcohol.

ACTION AND USES: Without material advantage over other manganese salts.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Manna (Manna), Manna, U. S. P.—A dried sap.

ACTION AND USES: Mild laxative; antiquated.

DOSAGE: 15 Gm. or 4 drachms (U. S. P.) as infusion.

Syrupus Mannae (Syr. Mann.), *Syrup of Manna*, N. F.—Manna (12.5%) and alcohol in syrup. Absolute alcohol content about 6 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

MASSAE, MASSES.—Soft solid preparations of such consistency that they can readily be made into pills. For the various masses, see under the names of the respective drugs.

Mastiche (Mastic), Mastic, N. F.—A resin.

Completely soluble in ether and almost completely soluble in alcohol.

ACTION AND USES: Carminative, without special advantage over cardamom or fennel.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Matico (Matic.), Matico, N. F. (Matico Leaves).

ACTION AND USES: Obsolete genito-urinary "stimulant," without advantage over buchu, etc.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Matico (Fldext. Matic.), *Fluidextract of Matico*, N. F.—Matico (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Matricaria (Matricar.), Matricaria, U. S. P. (German Chamomile, Wild Chamomile).—Flower heads.

ACTION AND USES: Popularly used as a bitter and aromatic "tea" and poultice, any effects being due mostly to the hot water.

DOSAGE: 15 Gm. or 4 drachms (U. S. P.).

Mel (Mel), Honey, U. S. P.

ACTION AND USES: Demulcent.

Mel Depuratum (Mel Depurat.), Clarified Honey, U. S. P.

Mel Rosae (Mel Rosae), Honey of Rose, U. S. P.—Fluid-extract of rose and clarified honey. Absolute alcohol content about 5 per cent.

USES: Flavoring vehicle; rarely used.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Melilotus (Melilot.), Melilot, N. F. (Yellow Sweet Clover).—
Leaves and flowering tops.

ACTION AND USES: Formerly used as a stimulant and antispasmodic and in the treatment of neuralgia especially associated with debility. Now seldom used; no scientific indications for its use.

MELLITA, HONEYS.—Sweet liquids having honey as a base. For the various honeys, see under the names of the respective principal ingredients. For *Mel Rosae*, see under *Mel*; for *Mel Rosae et Sodii Boratis*, see under *Sodii Boras*.***Mentha Piperita (Menth. Pip.), Peppermint, U. S. P.**—
Leaves and flowering tops.

ACTION AND USES: Carminative and flavor, used chiefly in the form of the oil and spirit, which see under *Oleum Menthae Piperitae*.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Mentha Viridis (Menth. Vir.), Spearmint, U. S. P.—Leaves and flowering tops.

ACTION AND USES: Same as those of peppermint. See *Oleum Menthae Viridis*.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

***Menthol (Menthol), Menthol, U. S. P.**—Obtained from oil of peppermint or other mint oils.

Colorless crystals, with a strong peppermint odor and an aromatic taste. Slightly soluble in water; very soluble in alcohol, chloroform or petroleum benzin.

ACTION AND USES: Used locally as counterirritant and analgesic; sometimes internally as carminative.

DOSAGE: 0.06 Gm. or 1 grain.

Inunctum Mentholis (Inunct. Menthol.), Menthol Inunction, N. F.—Menthol (5%) and hydrous wool fat.

Inunctum Mentholis Compositum (*Inunct. Menthol. Co.*), *Compound Menthol Inunction*, N. F.—Menthol (5%), methyl salicylate (10%) and hydrous wool fat.

Menthol Camphoratum (*Menthol Camph.*), *Camphorated Menthol*, N. F. (Campho-Menthol, N. F. III, Camphor and Menthol).—Camphor (47.5%), menthol (47.5%) and alcohol.

Nebula Mentholis (*Nebul. Menthol.*), *Menthol Spray*, N. F.—Menthol (2%) in light liquid petrolatum.

Nebula Mentholis Composita (*Nebul. Menthol. Co.*), *Compound Menthol Spray*, N. F.—Menthol (1%), camphor (1%), methyl salicylate, eucalyptol and oil of cinnamon in light liquid petrolatum.

Petroxolinum Mentholis (*Petrox. Menthol.*), *Menthol Petroxolin*, N. F. (Menthol Petrox).—Menthol (17%) in liquid petroxolin.

Menyanthes (*Menyanth.*), *Menyanthes*, N. F. (Buckbean, Marsh Trefoil).—Leaves of *Menyanthes trifoliata*.

ACTION AND USES: Obsolete bitter; no special indications for its use.

DOSAGE: 1 Gm. or 15 grains (N. F.).

**Methylis Salicylas* (*Methyl. Salicyl.*), *Methyl Salicylate*, U. S. P. (*Oleum Gaultheriae*, U. S. P. VIII, Oil of Wintergreen, *Oleum Betulae*, U. S. P. VIII, Oil of Sweet Birch, Oil of Teaberry).—Produced synthetically or obtained from wintergreen (oil of wintergreen) or from birch (oil of sweet birch).

Colorless, yellowish or reddish liquid, with a wintergreen odor and taste. Sparingly soluble in water and miscible with alcohol and glacial acetic acid.

ACTION AND USES: Chiefly used as flavor. In sufficient amounts, antirheumatic and antipyretic.

DOSAGE: 0.75 Cc. or 12 minims (U. S. P.). May be applied externally either pure or diluted (10%) in ointment. For absorption through the skin is best diluted with a fatty oil.

Petroxolinum Methylis Salicylatis (*Petrox. Methyl. Salic. Methyl Salicylate Petroxolin*, N. F. (Methyl Salicy Petrox).—Methyl salicylate (20%), in liquid petroxolin.

Methylthioninae Chloridum (*Methylthionin. Chlor.*), *Methylthionine Chloride*, U. S. P. (*Methylthioninae Hydrochloridum*, U. S. P. VIII, Methylene Blue).

Dark green powder or crystals with a bronze-like luster. Freely soluble in water and alcohol, forming a deep blue solution.

ACTION AND USES: Sometimes used as urinary antiseptic; also in the treatment of malaria and some neuralgic conditions. In the latter use, is of doubtful value.

DOSAGE: 0.15 Gm. or $2\frac{1}{2}$ grains (U. S. P.).

Mezereum (Mezereum), Mezereum, U. S. P. (Mezereon).—A bark.

ACTION AND USES: Antiquated antisyphilitic and vesicant. *Fluidextractum Mezerei (Fldext. Mezer.)*, *Fluidextract of Mezereum, N. F.*—Mezereum (100%). Absolute alcohol content about 70 per cent.

MISTURAE, MIXTURES.—Liquid preparations containing insoluble or partly soluble medicinal substances suspended in them.

For the various official mixtures, see under the names of the respective principal ingredients. For *Mistura Adstringens*, see under *Cupri Sulphas*; for *Mistura Camphorae Acida*, *Mistura Carminativa* and *Mistura Glycyrrhizae Composita*, under *Opium*; for *Mistura Ferri Composita*, under *Ferri Carbonas*; for *Mistura Camphorae Aromatica* and *Mistura Oleo-Balsamica*, under *Oleum Lavandulae*; for *Mistura Pectoralis Stokes*, under *Ammonii Carbonas*.

Morphina (Morphina), Morphine, U. S. P.—An alkaloid obtained from opium.

Colorless or white, odorless crystals or powder. Very slightly soluble in water (1:3,340) and soluble in alcohol (1:210).

ACTION AND USES: Central analgesic, hypnotic and respiratory sedative and depressant; also constipative. Is only slightly depressant to the circulation. Eliminated mainly through the bowels. Should not be used to control cough if this is associated with marked expectoration. Its habit-forming power should be constantly remembered.

DOSAGE: 0.008 Gm. or $\frac{1}{8}$ grain (U. S. P.).

***Morphinae Hydrochloridum (Morph. Hydrochl.), Morphine Hydrochloride, U. S. P. (Morphine Chloride).**

White, odorless needles, masses or powder. Soluble in water (1:17.5) and in alcohol (1:52).

ACTION AND USES: See *Morphina*.

DOSAGE: 0.008 Gm. or $\frac{1}{8}$ grain (U. S. P.).

Syrupus Morphinae et Acaciae (Syr. Morph. et Acac.), *Syrup of Morphine and Acacia, N. F.* (*Syrupus Pectoralis, N. F. III*, Jackson's Pectoral Syrup).—Morphine

hydrochloride (0.055%) flavored with oil of sassafras in syrup of acacia.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Morphinae Sulphas (Morph. Sulph.), Morphine Sulphate, U. S. P.—**

White, odorless crystals or masses. Soluble in water (1:15.2) and slightly soluble in alcohol (1:565).

ACTION AND USES: See *Morphina*.

DOSAGE: 0.008 Gm. or $\frac{1}{8}$ grain (U. S. P.).

Syrupus Pini Strobi Compositus cum Morphina (Syr. Pin. Strob. c. Morph.), *Compound Syrup of White Pine with Morphine, N. F.* (Syrupus Pini Strobi Compositus, N. F. III).—Morphine sulphate (0.04%) in compound syrup of white pine. Absolute alcohol content about 20 per cent.

Uses: Objectionable expectorant mixture containing morphine.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Moschus (Mosch.), Musk, U. S. P. (Tonquin Musk, Deer Musk).—A dried animal secretion.

ACTION AND USES: Antiquated analeptic.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Tinctura Moschi (Tr. Mosch.), *Tincture of Musk, U. S. P.*—Musk (5%) in diluted alcohol. Absolute alcohol content about 48 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

MUCILAGINES, MUCILAGES.—Aqueous solutions of gums or the mucilaginous principles of vegetable substances. For the various mucilages, see under the names of the respective drugs.

MULLAE, MULLS.—Ointments of high fusing points spread on soft muslin or "mull." For the various mulls, see under the names of the respective active constituents.

Myrica (Myrica), Bayberry Bark, N. F.—Bark of the roo

ACTION AND USES: Formerly recommended as digest nutritive and nervous stimulant. No scientific indication for its use.

DOSAGE: 0.5 Gm. or 8 grains (N. F.).

Pulvis Myricae Compositus (Pulv. Myric. Co.), *Compound Powder of Bayberry, N. F.* (Composition Pow.

Bayberry bark (60%), Jamaica ginger (30%), capsicum and clove.

USES: An obsolete carminative.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Myristica (Myrist.), Myristica, U. S. P. (Nutmeg).

ACTION AND USES: Aromatic and carminative.

***Myrrha (Myrrh.), Myrrh, U. S. P. (Gum Myrrh).—A gum resin.**

ACTION AND USES: Protective and local stimulant to the mucous membranes. Internally, carminative.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

***Tinctura Myrrhae (Tr. Myrrh.), Tincture of Myrrh, U. S. P.—Myrrh (20%) in alcohol. Absolute alcohol content about 87 per cent.**

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

NEBULAE, SPRAYS.—Spray solutions, intended for the application of medicaments in solution to the throat and nose by means of suitable atomizers. For Nebula Aromatica, see under *Petrolatum Liquidum*; for the various other sprays, see under *Eucalyptol*, *Menthol* and *Thymol*.

Nitrogenii Monoxidum (Nitrogen. Monox.), Nitrogen Monoxide, U. S. P. (Nitrous Oxide, "Laughing Gas").—N₂O.

A colorless gas with a slight characteristic odor and a sweetish taste.

ACTION AND USES: For inhalation anesthesia.

***Nux Vomica (Nux Vom.), Nux Vomica, U. S. P. (Strychni Semen, P. I.).—Seeds yielding not less than 2.5 per cent. of alkaloids, chiefly strychnine and brucine.**

ACTION AND USES: Used as bitter stomachic and tonic, depending on its strychnine.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

***Extractum Nucis Vomicae (Ext. Nuc. Vom.), Extract of Nux Vomica, U. S. P. (Powdered Extract of Nux Vomica, Nucis Vomicae Extractum P. I.).—Yields about 16 per cent. of alkaloids.**

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (U. S. P.).

Fluidextractum Nucis Vomicae (Fldext. Nuc. Vom.), Fluid-extract of Nux Vomica, U. S. P.—Nux vomica (100%) yielding about 2.5 per cent. of alkaloids. Absolute alcohol content about 60 per cent.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.).

**Tinctura Nucis Vomicae (Tr. Nuc. Vom.), Tincture of Nux Vomica, U. S. P. (Strychni Tinctura, P. I.).—Nux vomica (10%) yielding about 0.25 per cent. of alkaloids. Absolute alcohol content about 71 per cent.*

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

OLEA INFUSA (OL. INF.), INFUSED OILS, N. F.—Liquid preparation made by extracting drugs by fixed oils.

OLEATA, OLEATES.—Oleates are combinations of alkaloids or metallic oxids with oleic acid or with mixtures of oleic acid with a fixed oil. They have no real advantage over simpler ointments. For the official oleates see under the names of the respective drugs.

OLEORESINAE, OLEORESINS.—Oleoresins are thick liquid preparations made by extracting the drug with ether, alcohol or acetone. For the official oleoresins see under the names of the respective drugs.

OLEOSACCHARA (OLEOSACCHARA), OIL-SUGARS, N. F. (Elaeosacchara).—Mixtures of volatile oil (2%) with sugar in fine powder.

Oleum Aethereum (Ol. Aether.), Ethereal Oil, N. F.—A volatile liquid consisting of equal volumes of heavy oil of wine (chiefly ethyl esters of sulphuric acid) and ether. An indefinite and obsolete ingredient of the medieval "Hoffmann's anodyne."

Oleum Amygdalae Amarae (Ol. Amygd. Amar.), Oil of Bitter Almond, U. S. P. (Bitter Almond Oil).—A volatile oil yielding benzaldehyde (not less than 85%) and hydrocyanic acid (about 3%).

Slightly soluble in water; miscible with alcohol or ether.

ACTION AND USES: Flavoring agent, especially for emulsions. Also used as a "sedative," but probably without value.

DOSAGE: 0.03 Cc. or ½ minim (U. S. P.).

Aqua Amygdalae Amarae (Aq. Amygd. Amar.), Bitter Almond Water, U. S. P.—A solution of oil of bitter almond in water.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Elixir Amygdalae Compositum (Elix. Amygd. Co.), Compound Elixir of Almond, N. F.—Oil of bitter almond, vanillin, orange flower water, syrup, water and alcohol. Absolute alcohol content about 5 per cent.

Spiritus Amygdalae Amarae (Sp. Amygd. Amar.), *Spirit of Bitter Almond*, U. S. P.—Oil of bitter almond, alcohol and water. Absolute alcohol content about 76 per cent.

Oleum Amygdalae Expressum (Ol. Amygd. Exp.), *Expressed Oil of Almond*, U. S. P. (Oil of Sweet Almond).—A fixed oil; similar to olive oil.

ACTION AND USES: Emollient, nutritive and laxative.

Oleum Anisi (Ol. Anisi), *Oil of Anise*, *Oil of Star Anise*, U. S. P. (Anise Oil).—A volatile oil from anise and from star anise.

Very slightly soluble in water; very soluble in alcohol.

ACTION AND USES: Aromatic carminative and flavor.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.), on sugar.

Aqua Anisi (Aq. Anisi), *Anise Water*, U. S. P.—A solution of oil of anise in water.

Spiritus Anisi (Sp. Anisi), *Spirit of Anise*, U. S. P.—Oil of anise (10%) in alcohol. Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Oleum Aurantii (Ol. Aurant.), *Oil of Orange*, U. S. P. (*Oleum Aurantii Corticis*, U. S. P. VIII, *Orange Oil*, *Oil of Sweet Orange*).—A volatile oil.

Nearly insoluble in water; soluble in alcohol and ether.

ACTION AND USES: Aromatic flavor.

Spiritus Aurantii Compositus (Sp. Aurant. Co.), *Compound Spirit of Orange*, U. S. P.—Oils of orange, lemon, coriander and anise, in alcohol. Absolute alcohol content about 69 per cent.

Oleum Aurantii Amari (Ol. Aurant. Amar.), *Oil of Bitter Orange*, N. F.—A volatile oil.

Practically insoluble in water; soluble in alcohol (1:4).

ACTION AND USES: Flavoring agent.

Elixir Aurantii Amari (Elix. Aurant. Amar.), *Elixir of Bitter Orange*, N. F. (Replacing *Elixir Curassao*, N. F. III).—Oil of bitter orange, tincture of bitter orange peel, stronger orange flower water, in syrup, water and alcohol. Absolute alcohol content about 28 per cent.

Oleum Aurantii Florum (Ol. Aurant. Flor.), *Oil of Orange Flowers*, N. F. (Oil of Neroli).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Flavoring agent.

Oleum Bergamottae (Ol. Bergam.), Oil of Bergamot, N. F.—
A volatile oil.

Practically insoluble in water; freely soluble in alcohol.

ACTION AND USES: Flavoring agent.

Spiritus Odoratus (Sp. Odorat.), *Perfumed Spirit*, N. F.—
Oils of bergamot, lemon, rosemary, lavender and orange
flowers, acetic ether, water and alcohol. Absolute alcohol
content about 80 per cent.

Oleum Betulae Empyreumaticum Rectificatum (Ol. Bet.
Empyr. Rect.), Rectified Oil of Birch Tar, N. F. (Recti-
fied Empyroligneous Oil of Birch, *Oleum Rusci Rectifi-*
catum).—An oil obtained by the dry distillation of the
bark and wood of white birch.

Insoluble in water; soluble in dehydrated alcohol, ether and
chloroform.

ACTION AND USES: Similar to tar; employed locally as
a stimulant and antiseptic in the treatment of psoriasis
and eczema.

DOSAGE: 2 to 10 per cent. ointment.

Oleum Cadinum (Ol. Cadin.), Oil of Cade, U. S. P. (Cade
Oil, Oil of Juniper Tar, *Oleum Juniperi Empyreumaticum*).—An empyreumatic oil obtained by the dry distilla-
tion of juniper wood.

Nearly insoluble in water; only partially soluble in alcohol.

ACTION AND USES: Epidermal stimulant in chronic
inflammatory skin diseases, acting similarly to tar.

DOSAGE: 1 to 10 per cent. ointment.

Petroxolinum Cadini (*Petrox. Cadin.*), *Cade Petroxolin*,
N. F. (Cade Petrox).—Oil of cade (25%) in liquid
petroxolin.

Oleum Cajuputi (Ol. Cajup.), Oil of Cajuput, U. S. P.
(Cajuput Oil, Oil of Cajuput).—A volatile oil.

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Counterirritant.

Oleum Cardamomi (Ol. Cardam.), Oil of Cardamom, N. F.—
A volatile oil.

Nearly insoluble in water; very soluble in alcohol.

ACTION AND USES: Aromatic flavoring agent.

Elixir Cardamomi Compositum (Elix. Card. Co.), Compound Elixir of Cardamom, N. F.—Compound spirit of cardamom, alcohol, syrup and water. Absolute alcohol content about 9 per cent.

Spiritus Cardamomi Compositus (Sp. Card. Co.), Compound Spirit of Cardamom, N. F.—Oil of cardamom, anethol, oils of orange, cinnamon, caraway and clove, in alcohol. Absolute alcohol content about 72 per cent.

USES: Flavoring agent.

Oleum Cari (Ol. Cari), Oil of Caraway, U. S. P. (Caraway Oil).—A volatile oil.

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Aromatic carminative.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

***Oleum Caryophylli** (Ol. Caryoph.), Oil of Clove, U. S. P. (Clove Oil, Oil of Cloves).—A volatile oil.

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Externally, rubefacient and counter-irritant; internally, carminative; to some extent a local anesthetic.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

***Oleum Cassiae** (Ol. Cass.), Oil of Cinnamon, U. S. P. (Oleum Cinnamomi, U. S. P. VIII, Cassia Oil).—A volatile oil.

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Aromatic flavor.

***Aqua Cinnamomi** (Aq. Cinnam.), Cinnamon Water, U. S. P.—A saturated solution of oil of cinnamon in water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Spiritus Cinnamomi (Sp. Cinnam.), Spirit of Cinnamon, U. S. P.—Oil of cinnamon (10%) in alcohol. Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

***Oleum Chenopodii** (Ol. Chenopod.), Oil of Chenopodium, U. S. P. (Oil of American Wormseed).—A volatile oil.

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Anthelmintic, especially for round-worms and hookworms.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.). Subminimal doses repeated at intervals of several days become toxic.

Oleum Coriandri (Ol. Coriand.), Oil of Coriander, U. S. P. (Coriander Oil).

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Carminative aromatic.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Oleum Cubebae (Ol. Cubeb.), Oil of Cubeb, U. S. P. (Cubeb Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol (1:18).

ACTION AND USES: Used internally as a stimulating antiseptic, expectorant and against subacute gonorrhea. Of doubtful value.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

***Oleum Eucalypti** (Ol. Eucalypt.), Oil of Eucalyptus, U. S. P. (Eucalyptus Oil).—A volatile oil.

Nearly insoluble in water; freely soluble in alcohol.

ACTION AND USES: Largely used especially for inhalation in the form of vapor or oily sprays (about 10%) as antiseptic and deodorant, for subacute and chronic inflammations of the respiratory mucosae.

Oleum Foeniculi (Ol. Foenic.), Oil of Fennel, U. S. P. (Fennel Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatic carminative frequently employed with purgative medicines to prevent griping.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Aqua Foeniculi (Aq. Foenic.), Fennel Water, U. S. P.—A solution of oil of fennel in water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Oleum Gossypii Seminis (Ol. Gossyp. Sem.), Cottonseed Oil, U. S. P.—A fixed oil.

ACTION AND USES: Frequently employed in place of olive oil, in preparations for external use.

Oleum Juniperi (Ol. Junip.), Oil of Juniper, U. S. P. (Juniper Oil, Oil of Juniper Berries).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Irritant diuretic.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.); not given as such.

Spiritus Juniperi (*Sp. Junip.*), *Spirit of Juniper*, U. S. P.—Oil of juniper (5%) in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Spiritus Juniperi Compositus (*Sp. Junip. Co.*), *Compound Spirit of Juniper*, U. S. P.—Oil of juniper (0.4%) and oils of caraway and fennel in alcohol and water. Absolute alcohol content about 66 per cent.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Oleum Lavandulae (*Ol. Lavand.*), *Oil of Lavender*, U. S. P. (*Oleum Lavandulae Florum* U. S. P. VIII).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatic and flavoring agent.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Acetum Aromaticum (*Acet. Arom.*), *Aromatic Vinegar*, N. F.—Acetic acid, alcohol and water, aromatized with oils of lavender, rosemary, juniper, peppermint, cinnamon, lemon and clove.

USES: Refreshing inhalant and counterirritant.

Mistura Camphorae Aromatica (*Mist. Camph. Arom.*), *Aromatic Camphor Mixture*, N. F. (Parrish's Camphor Mixture).—Camphor water (about 70%) in compound tincture of lavender and sugar. Absolute alcohol content about 17.5 per cent.

USES: Aromatic carminative.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Mistura Oleo-Balsamica (*Mist. Ol.-Balsam.*), *Oleo-Balsamic Mixture*, N. F.—Oil of lavender, eugenol, oils of thyme, cinnamon, lemon and nutmeg with balsam of Peru in alcohol. Absolute alcohol content about 90 per cent.

ACTION AND USES: Local stimulant.

Spiritus Lavandulae (*Sp. Lavand.*), *Spirit of Lavender*, U. S. P.—Oil of lavender (5%) in alcohol. Absolute alcohol content about 90 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Lavandulae Composita (*Tr. Lavand. Co.*), *Compound Tincture of Lavender*, U. S. P. (Compound Spirit of Lavender).—Oil of lavender, oil of rosemary, Saigon cinnamon, clove, nutmeg, and red saunders in alcohol and water. Absolute alcohol content about 70 per cent.

USES: Aromatic flavor and rubefacient.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Oleum Limonis (Ol. Limon.), Oil of Lemon, U. S. P. (Lemon Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Flavoring agent.

***Oleum Lini (Ol. Lini), Linseed Oil, U. S. P. (Oil of Flaxseed, Raw Linseed Oil).**—A fixed oil.

ACTION AND USES: Internally, laxative; its disagreeable taste and odor have made it unpopular. Externally, protective emollient, especially as Linimentum Calcis (which see under Calx) in burns.

DOSAGE: 30 Cc. or 1 fluidounce (U. S. P.).

***Oleum Menthae Piperitae (Ol. Menth. Pip.), Oil of Peppermint, U. S. P. (Peppermint Oil).**—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatic carminative and flavoring agent.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

***Aqua Menthae Piperitae (Aq. Menth. Pip.), Peppermint Water, U. S. P.**—A saturated solution of oil of peppermint in distilled water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

***Spiritus Menthae Piperitae (Sp. Menth. Pip.), Spirit of Peppermint, U. S. P. (Essence of Peppermint).**—Oil of peppermint (10%), colored with peppermint in alcohol. Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Trochisci Menthae Piperitae (Troch. Menth. Pip.), Troches of Peppermint, N. F.—Oil of peppermint, with sugar and mucilage of tragacanth.

DOSAGE: 1 troche (N. F.).

Oleum Menthae Viridis (Ol. Menth. Vir.), Oil of Spearmint, U. S. P. (Spearmint Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatic carminative and flavoring agent.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Aqua Menthae Viridis (Aq. Menth. Vir.), Spearmint Water, U. S. P.—A solution of oil of spearmint in water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Spiritus Menthae Viridis (Sp. Menth. Vir.), Spirit of Spearmint, U. S. P.—Oil of spearmint (10%), colored with

spearmint, in alcohol. Absolute alcohol content about 85 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

***Oleum Morrhuæ** (Ol. Morr.), Cod Liver Oil, U. S. P. (Oleum Jecoris Aselli).—A fixed oil.

ACTION AND USES: Generally considered to be a food rather than a medicine. Its value depends probably entirely on its nutritive property. Preferably administered as such or in the form of recently prepared emulsions.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Emulsum Olei Morrhuæ (Emuls. Ol. Morr.), *Emulsion of Cod Liver Oil*, U. S. P.—Cod liver oil (50%) with acacia, syrup, methyl salicylate and water. Other flavors may be substituted for the methyl salicylate.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Emulsum Olei Morrhuæ cum Calcii Lactophosphate (Emul. Ol. Morr. c. Calc. Lactophos.), *Emulsion of Cod Liver Oil with Calcium Lactophosphate*, N. F.—Cod liver oil (50%), calcium lactophosphate (5%), lactic acid, acacia, syrup of Tolu and water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Emulsum Olei Morrhuæ cum Calcii Phosphate (Emul. Ol. Morr. c. Calc. Phos.), *Emulsion of Cod Liver Oil with Calcium Phosphate*, N. F. (Emulsion of Cod Liver Oil with Phosphate of Lime).—Cod liver oil (50%), precipitated calcium phosphate (3.5%), syrup of Tolu, acacia and water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Emulsum Olei Morrhuæ cum Hypophosphitibus (Emul. Ol. Morr. c. Hypophos.), *Emulsion of Cod Liver Oil with Hypophosphites*, N. F.—Cod liver oil (50%), calcium hypophosphite (1%), potassium hypophosphite and sodium hypophosphite (each 0.5%), acacia, syrup and water.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Emulsum Olei Morrhuæ cum Malto (Emul. Ol. Morr. c. Malt.), *Emulsion of Cod Liver Oil with Malt*, N. F.—Cod liver oil (30%), extract of malt (about 55%), tragacanth and water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Emulsum Olei Morrhuæ cum Pruno Virginiana (Emul. Ol. Morr. c. Prun. Virg.), *Emulsion of Cod Liver Oil with Wild Cherry*, N. F.—Cod liver oil (50%), fluidextract of wild cherry (6.5%), acacia, syrup of Tolu and water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Emulsum Olei Morrhuae cum Vitello (*Emul. Ol. Morrh. c. Vitel.*), *Emulsion of Cod Liver Oil with Egg*, N. F.—Cod liver oil (50%), glycerite of yolk of egg, syrup of Tolu and water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Oleum Myrciae (*Ol. Myrc.*), *Oil of Myrcia*, N. F. (*Oil of Bay*).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatizing agent used in bay-rum. *Spiritus Myrciae Compositus* (*Sp. Myrciae Co.*), *Compound Spirit of Myrcia*, N. F.—Oils of myrcia, orange, pimenta, alcohol and water. Absolute alcohol content about 58 per cent.

USES: Perfume and stimulant to the skin.

Oleum Myristicae (*Ol. Myrist.*), *Oil of Myristica*, U. S. P. (*Myristica Oil*, *Oil of Nutmeg*).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatic flavor and carminative.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Oleum Olivae (*Ol. Oliv.*), *Olive Oil*, U. S. P.—A fixed oil.

ACTION AND USES: Emollient, laxative and nutrient.

DOSAGE: 30 Cc. or 1 fluidounce (U. S. P.).

Oleum Picis Liquidae Rectificatum (*Ol. Pic. Liq. Rect.*), *Rectified Oil of Tar*, U. S. P. (*Oleum Picis Liquidae*, U. S. P. VIII, *Rectified Tar Oil*).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Externally antiseptic, irritant and parasiticide; used in skin diseases.

DOSAGE: Internally, 0.2 Cc. or 3 minims (U. S. P.).

Mistura Olei Picis (*Mist. Ol. Pic.*), *Mixture of Oil of Tar*, N. F. (*Mistura Picis Liquidae*, *Tar Mixture*).—Rectified oil of tar (3.5%) and chloroform (1%), with pure extract of glycyrrhiza, sugar and oil of peppermint in alcohol and water. Absolute alcohol content about 16 per cent.

USES: Complex, disagreeable expectorant.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Petroxolinum Picis (*Petrox. Pic.*), *Tar Petroxolin*, N. F. (*Tar Petrox.*).—Rectified oil of tar (25%) in liquid petroxolin.

Unguentum Picis Compositum (Ung. Pic. Co.), Compound Tar Ointment, N. F.—Rectified oil of tar (4%), zinc oxide (3%), and tincture of benzoin in yellow wax, lard and cottonseed oil.

Oleum Pimentae (Ol. Piment.), Oil of Pimenta, U. S. P. (Pimento Oil, Oil of Allspice).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Flavoring agent and carminative.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Oleum Pini Pumilionis (Ol. Pin. Puml.), Oil of Dwarf Pine Needles, U. S. P. (Dwarf Pine Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: More aromatic than oil of turpentine, and used as an inhalation in catarrh of the respiratory passages.

***Oleum Ricini** (Ol. Ricin.), Castor Oil, U. S. P.—A fixed oil.

Soluble in equal parts of alcohol.

ACTION AND USES: Effective and prompt cathartic, but likely to be followed by constipation and therefore not indicated in chronic constipation. Useful as an initial purge in acute diarrheas.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Emulsum Olei Ricini (Emul. Ol. Ricin.), Emulsion of Castor Oil, N. F.—Castor oil (35%), with acacia, flavored with tincture of vanilla, in syrup and water.

USES: Renders castor oil somewhat less disagreeable.

DOSAGE: 45 Cc. or 1½ fluidounces (N. F.).

Oleum Ricini Aromaticum (Ol. Ricin. Arom.), Aromatic Castor Oil, N. F.—Castor oil flavored with benzosulphinide, oil of cinnamon, oil of clove, vanillin and coumarin.

USES: A partly successful attempt to render castor oil less disagreeable.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Oleum Rosmarini (Ol. Rosmar.), Oil of Rosemary, U. S. P. (Rosemary Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Aromatic flavor and carminative.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

***Oleum Santali** (Ol. Santal.), Oil of Santal, U. S. P. (Santalwood Oil, Oil of Sandalwood).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Urinary disinfectant and stimulant. Used in subacute stages of cystitis and gonorrhea.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.), preferably in capsules.

Oleum Sassafras (Ol. Sassaf.), Oil of Sassafras, U. S. P. (Sassafras Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Flavor; used externally as rubefacient counterirritant.

Oleum Sesami (Ol. Sesam.), Sesame Oil, U. S. P. (Teel Oil, Benne Oil).—A fixed oil.

ACTION AND USES: Emollient, used as substitute for olive oil.

***Oleum Sinapis Volatile (Ol. Sinap. Vol.), Volatile Oil of Mustard, U. S. P. (Mustard Oil).—Produced synthetically or from black mustard seed.**

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Powerful irritant and rapid vesicant; employed as counterirritant dissolved in alcohol or fixed oils.

Linimentum Sinapis Compositum (Lin. Sinap. Co.), Compound Liniment of Mustard, N. F.—Volatile oil of mustard (3%), fluidextract of mezereum (20%) and camphor (6%) in castor oil and alcohol.

USES: Complex and antiquated counterirritant.

Spiritus Sinapis (Sp. Sinap.), Spirit of Mustard, N. F.—Volatile oil of mustard (2%) in alcohol.

USES: Rubefacient.

***Oleum Terebinthinae (Ol. Tereb.), Oil of Turpentine, U. S. P. (Turpentine Oil, "Spirits of Turpentine").—A volatile oil obtained from the oleoresin of pine wood (turpentine).**

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Applied externally, rubefacient and counterirritant. For internal use, see *Oleum Terebinthinae Rectificatum*, below.

Oleum Terebinthinae Rectificatum (Ol. Tereb. Rect.), Rectified Oil of Turpentine, U. S. P. (Rectified Turpentine Oil).—Redistilled oil of turpentine.

USES: Antiseptic, anthelmintic, diuretic and carminative. Used to relieve tympanites.

DOSAGE: 0.3 Cc. or 5 minims (U. S. P.).

Emulsum Olei Terebinthinae (*Emuls. Ol. Tereb.*), *Emulsion of Oil of Turpentine*, U. S. P.—Rectified oil of turpentine (15%) with expressed oil of almond, syrup, acacia and water.

DOSAGE: 2 Cc. or $\frac{1}{2}$ fluidrachm (U. S. P.).

Linimentum Terebinthinae (*Lin. Terebinth.*), *Turpentine Liniment*, U. S. P. (Kentish's Ointment).—Rosin cerate (65%) in oil of turpentine.

Linimentum Terebinthinae Aceticum (*Lin. Terebinth. Acet.*), *Acetic Turpentine Liniment*, N. F. (*Linimentum Album*, Stokes' Liniment, St. John Long's Liniment).—Oil of turpentine (40%) and acetic acid (8%) with oil of lemon, fresh egg and rose water.

**Oleum Theobromatis* (*Ol. Theobrom.*), *Oil of Theobroma*, U. S. P. (Butter of Cacao, Cacao Butter).—A solid fixed oil.

Melts at body temperature.

ACTION AND USES: Used principally in suppositories, etc.; sometimes in emollient ointments and as a lubricant in massage.

Oleum Thymi (*Ol. Thymi*), *Oil of Thyme*, U. S. P. (Thyme Oil).—A volatile oil.

Nearly insoluble in water; soluble in alcohol.

ACTION AND USES: Rubefacient and counterirritant; usually applied as a liniment, diluted with a fixed oil or the well-known soap liniment. Internally, carminative but now seldom used.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

**Oleum Tiglii* (*Ol. Tiglii*), *Croton Oil*, U. S. P.—A fixed oil.

ACTION AND USES: Very powerful cathartic and gastric and intestinal irritant, frequently causing violent vomiting and purging, sometimes followed by collapse. Externally, rubefacient and vesicant.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.). Preferably administered in drop doses on sugar or in a pill with bread or other absorbent.

Collodium Tiglii (*Collod. Tiglii*), *Croton Oil Collodion*, N. F.—Croton oil (10%) and flexible collodion.

USES: There is no advantage in applying croton oil with collodion.

Linimentum Tiglii (Lin. Tiglii), *Liniment of Croton Oil*, N. F. (*Linimentum Crotonis*).—Croton oil (13%), oil of cajuput, and alcohol.

Linimentum Tiglii Compositum (Lin. Tiglii Co.), *Compound Croton Oil Liniment*, N. F.—Croton oil, oil of sassafras and oil of turpentine (each 20%) and olive oil.

***Opium** (Opium), **Opium**, U. S. P.—The dried juice of the opium poppy. Yields not less than 9.5 per cent. of anhydrous morphine.

ACTION AND USES: Like those of morphine, but opium is absorbed more slowly, and is generally preferred in the treatment of the gastro-intestinal tract, especially for the purpose of checking diarrhea. It is ineffective for external or local application, because the opium alkaloids are not absorbed to any appreciable extent through the unbroken skin.

The routine use of complicated mixtures containing opiates deserves condemnation, since opium should not be prescribed unless it is adapted to the conditions of the individual patient.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Acetum Opii (Acet. Opii), *Vinegar of Opium*, N. F.—Opium (10%) in diluted acetic acid, flavored with nutmeg and sugar.

USES: An obsolete and superfluous preparation.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

**Extractum Opii* (Ext. Opii), *Extract of Opium*, U. S. P. (Powdered Extract of Opium, Opii Extractum, P. I.).—One Gm. extract represents about 2 Gm. opium. Yields about 20 per cent. of anhydrous morphine.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

Linimentum Opii Compositum (Lin. Opii Co.), *Compound Liniment of Opium*, N. F. (Canada Liniment).—Tincture of opium (10%), camphor (1.75%), oil of peppermint (2.5%) in alcohol, fresh egg albumen, ammonia water and oil of turpentine. Absolute alcohol content about 28 per cent.

USES: Irrational opium preparation.

Mistura Camphorae Acida (Mist. Camph. Acid.), *Acid Camphor Mixture*, N. F. (*Mistura Antidysenterica*, Hope's Mixture).—Tincture of opium (1.2%) and nitric acid (1.75%) in camphor water.

USES: Against diarrhea.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Mistura Carminativa (*Mist. Carminat.*), *Carminative Mixture*, N. F. (Dalby's Carminative).—Tincture of opium (2.5%), magnesium carbonate (6.5%), potassium carbonate (0.3%), with oils of caraway, fennel and peppermint in syrup and water.

USES: Against colic.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

Mistura Glycyrrhizae Composita (*Mist. Glycyrrh. Co.*), *Compound Mixture of Glycyrrhiza*, U. S. P. (Brown Mixture).—Camphorated tincture of opium (12%), antimony and potassium tartrate (0.024%), with pure extract of glycyrrhiza, syrup, acacia and spirit of nitrous ether in water. Absolute alcohol content about 8 per cent.

USES: A misnamed but popular expectorant, efficient mainly through its opium and antimony.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Mistura Opii et Chloroformi Composita (*Mist. Opii et Chlorof. Co.*), *Compound Mixture of Opium and Chloroform*, N. F. (Squibb's Diarrhea Mixture).—Tincture of opium (20%), spirit of camphor (20%), tincture of capsicum (10%), chloroform (8%) and alcohol. Absolute alcohol content about 75 per cent.

USES: Against diarrhea.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Mistura Opii et Rhei Composita (*Mist. Opii et Rhei Co.*), *Compound Mixture of Opium and Rhubarb*, N. F. (Sun Cholera Mixture).—Tincture of opium (20%), tincture of rhubarb (10%), tincture of capsicum (10%), spirit of camphor (20%), spirit of peppermint and alcohol. Absolute alcohol content about 78 per cent.

USES: Against diarrhea.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Mistura Opii et Sassafras (*Mist. Opii et Sassaf.*), *Mixture of Opium and Sassafras*, N. F. (*Mistura Sassafras et Opii*, N. F. III, *Mistura Opii Alkalina*, Godfrey's Cordial).—Tincture of opium (3.5%) and oil of sassafras (0.1%) with alcohol and potassium carbonate in syrup and water. Absolute alcohol content about 6.5 per cent.

USES: Sedative.

DOSAGE: Infants, 0.3 Cc. or 5 minims (N. F.).

**Opii Pulvis* (*Opii Pulv.*), *Powdered Opium*, U. S. P. (*Opii Pulvis*, P. I.).—Contains about 10.25 per cent. of anhydrous morphine.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Opium Deodoratum (*Opium Deod.*), *Deodorized Opium*, U. S. P.—Opium extracted with benzin. Yields about 10.25 per cent of anhydrous morphine.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Opium Granulatum (*Opium Gran.*), *Granulated Opium*, U. S. P.—Yields about 10.25 per cent. of anhydrous morphine.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Pilulae Opii, Digitalis et Quininae (*Pil. Opii Digit. et Quin.*), *Pills of Opium, Digitalis and Quinine*, N. F. (Niemeyer Pills for Phthisis).—Each pill contains powdered opium (0.01 Gm. or $\frac{1}{6}$ grain), digitalis and quinine sulphate (each 0.065 Gm. or 1 grain) with clarified honey.

DOSAGE: 1 pill (N. F.).

Pilulae Opii et Camphorae (*Pil. Opii et Camph.*), *Pills of Opium and Camphor*, N. F.—Each pill contains powdered opium (0.065 Gm. or 1 grain) and camphor (0.13 Gm. or 2 grains) with clarified honey.

DOSAGE: 1 pill (N. F.).

Pilulae Opii et Plumbi (*Pil. Opii et Plumb.*), *Pills of Opium and Lead*, N. F.—Each pill contains powdered opium and lead acetate (each 0.065 Gm. or 1 grain) with clarified honey.

USES: Constipative. The internal use of lead is rarely advisable.

DOSAGE: 1 pill (N. F.).

Pulvis Cretae et Opii Aromaticus (*Pulv. Cret. et Opii Arom.*), *Aromatic Powder of Chalk and Opium*, N. F.—Powdered opium (2.5%) and aromatic powder of chalk.

DOSAGE: 1 Gm. or 15 grains (N. F.).

**Pulvis Ipecacuanhae et Opii* (*Pulv. Ipecac. et Opii*), *Powder of Ipecac and Opium*, U. S. P. (Compound Powder of Ipecac, Dover's Powder, Opii et Ipecacuanhae Pulvis Compositus, P. I.).—Powdered opium and ipecac (each 10%) with sugar of milk.

USES: Especially as diaphoretic in incipient colds.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Syrupus Ipecacuanhae et Opii (*Syr. Ipecac. et Opii*), *Syrup of Ipecac and Opium*, N. F. (Syrup of Dover's Powder).—Tincture of ipecac and opium (8.5%) flavored with spirit of cinnamon and cinnamon water in syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Tinctura Ipecacuanhae et Opium (Tr. Ipecac. et Opium), Tincture of Ipecac and Opium, N. F. (Tincture of Dover's Powder).—Tincture of deodorized opium (100%) and fluidextract of ipecac (10%) in diluted alcohol. Absolute alcohol content about 15 per cent.

DOSAGE: 0.5 Cc. or 8 minims (N. F.).

**Tinctura Opium (Tr. Opium), Tincture of Opium, U. S. P.* (Laudanum, *Opium Tinctura, P. I.*).—Granulated opium (10%) yielding about 1 per cent. of anhydrous morphine in diluted alcohol. Absolute alcohol content about 46 per cent.

DOSAGE: 0.5 Cc. or 8 minims.

**Tinctura Opium Camphorata (Tr. Opium Camph.), Camphorated Tincture of Opium, U. S. P.* (Paregoric, *Opium Tinctura Benzoici P. I.*).—Powdered opium and camphor (each 0.4%), benzoic acid, oil of anise and glycerin in diluted alcohol. Absolute alcohol content about 46 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Tinctura Opium Crocata (Tr. Opium Crocat.), Tincture of Opium with Saffron, N. F. (Sydenham's Laudanum, *Tinctura Opium Crocata, P. I.*).—Granulated opium (10%) with saffron, Saigon cinnamon and clove in diluted alcohol. Absolute alcohol content about 45 per cent.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

**Tinctura Opium Deodorati (Tr. Opium Deod.), Tincture of Deodorized Opium, U. S. P.*—Opium (10%, a purified or deodorized aqueous extract yielding about 1% anhydrous morphine), in alcohol and water. Absolute alcohol content about 19 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

Tinctura Pectoralis (Tr. Pectoral.), Pectoral Tincture, N. F. (Guttae Pectorales, Pectoral Drops, Bateman's Pectoral Drops).—Opium (0.4%), gambir (about 0.3%), camphor (0.4%), with oil of anise and caramel in diluted alcohol. Absolute alcohol content about 50 per cent.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

Ovi Albumen Recens (Ovi Album.), Fresh Egg Albumen, N. F.

ACTION AND USES: Used in pharmacy.

Ovi Vitellum Recens (Ovi Vitel.), Fresh Egg Yolk, N. F.

ACTION AND USES: Emulsifying agent

Glyceritum Vitelli (*Glycer. Vitell.*), *Glycerite of Egg Yolk*
N. F. (Glyconin).—Fresh yolk of egg and glycerin.

Ovum Gallinaceum (**Ovum Gallin.**), **Fresh Egg**, N. F.

ACTION AND USES: Used in pharmaceutical preparations.

***Oxygenium** (**Oxygen.**), **Oxygen**, U. S. P.

ACTION AND USES: Inhaled for relief of asphyxia, carbon monoxid poisoning, etc.

***Pancreatinum** (**Pancreat.**), **Pancreatin**, U. S. P.—Contains enzymes from the pancreas of the hog or beef, principally amylpsin, trypsin and steapsin. Converts not less than 25 per cent. of its own weight of starch into sugars.

A cream-colored powder with a faint odor.

ACTION AND USES: Chiefly for the predigestion of protein and starchy foods. Incompatible with acids and is therefore destroyed by the action of normal gastric juice. Occasionally used to assist gastric digestion in achlorhydria but the administration of dilute hydrochloric acid seems preferable. When the pancreatic juice is absent pancreatin may be given in salol-coated pills.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Liquor Pancreatini (*Liq. Pancreat.*), *Solution of Pancreatin*, N. F. (*Liquor Pancreaticus*, N. F. III. *Pancreatic Solution*).—Pancreatin (1.75%), sodium bicarbonate (5%), with compound spirit of cardamom, sodium chloride and chloroform in alcohol, glycerin and water. Absolute alcohol content about 6 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Pulvis Pancreatini Compositus (*Pulv. Pancreat. Co.*), *Compound Pancreatin Powder*, N. F. (*Pulvis Pancreaticus Compositus*, N. F. III, *Peptonizing Powder*).—Pancreatin (20%) and sodium bicarbonate.

USES: Predigestion of milk.

Papaveris Fructus (**Papav. Fruct.**), **Poppy Capsules**, N. F.—Fully grown unripe poppy heads.

ACTION AND USES: Weak and unreliable opiate.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Syrupus Papaveris (*Syr. Papaver.*), *Syrup of Poppy*, N. F.—Poppy capsules (10%) in syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Paracoto (**Paracot.**), **Paracoto**, N. F.—Bark of an unidentified tree.

ACTION AND USES: At one time used in diarrhea. The identity of the drug is uncertain and there is no scientific evidence for its value; hence it is now little used.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Fluidextractum Paracoto (Fldext. Paracoto), Fluidextract of Paracoto, N. F. (Fluidextractum Coto, N. F. III).—Paracoto (100%). Absolute alcohol content about 75 per cent.

DOSAGE: 0.3 Cc. or 5 minims (N. F.).

Tinctura Paracoto (Tr. Paracoto), Tincture of Paracoto, N. F. (Tinctura Coto, N. F. III).—Paracoto (12.5%) in alcohol. Absolute alcohol content about 90 per cent.

USES: Antidiarrheic, of doubtful efficacy.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Paraffinum (Paraff.), Paraffin, U. S. P.**—A purified mixture of solid hydrocarbons, usually obtained from petroleum.

A white, waxy, odorless and tasteless solid, greasy to the touch, melting between 50 and 57 C.

ACTION AND USES: Used in surgery for prosthetic purposes and in pharmacy for raising the melting point of ointments and similar products.

Paraformaldehydum (Paraform.), Paraformaldehyde, U. S. P. (Paraform, Trioxymethylene).—A polymeric form of formaldehyde.

White friable masses or powder with a slight odor of formaldehyde. Soluble in cold water, insoluble in alcohol or ether.

***ACTION AND USES:** Employed in place of formaldehyde to disinfect rooms by vaporization in presence of moisture (to insure formation of formaldehyde). As it is active only if decomposed there may be some doubt as to its efficiency. Internally, it has been used as intestinal antiseptic, but is probably useless and may be harmful.

***Paraldehydum (Paraldehyd.), Paraldehyde, U. S. P.**—A polymer of acetaldehyde.

A colorless liquid, having a strong, characteristic odor and an extremely unpleasant taste, and producing first a burning and then a cooling sensation in the mouth. Soluble in water (1:8), and miscible with alcohol or chloroform.

ACTION AND USES: Prompt and fairly active hypnotic and sedative devoid of serious side-effects.

DOSAGE: 2 Cc. or 30 minims (U. S. P.). Best administered with cracked ice or ice-cold liquids.

Pareira (Pareira), Pareira, N. F. (Pareira Brava).—Roots.

ACTION AND USES: Obsolete bitter tonic.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Pareirae (*Fldext. Pareir.*), *Fluidextract of Pareira*, N. F.—Pareira (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Passiflora (Passiflor.), Passion Flower, N. F. (Passion Vine).—Dried herbage.

ACTION AND USES: Exploited by manufacturers of proprietary remedies for the treatment of insomnia, but probably inert.

DOSAGE: 0.2 Gm. or 3 grains (N. F.).

Tinctura Passiflorae (*Tr. Passiflor.*), *Tincture of Passion Flower*, N. F.—Passion flower (20%), in diluted alcohol. Absolute alcohol content about 45 per cent.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

PASTAE DERMATOLOGICAE, DERMATOLOGIC

PASTES.—Dermatologic pastes are medicaments for external use. They are ointment-like mixtures containing starch or dextrin with glycerin, soft soap, petrolatum or lard mixed with some active medicinal agent. For the official dermatologic pastes see under the names of the principal constituents.

**Pelletierinae Tannas* (Pellet. Tann.), *Pelletierine Tannate*, U. S. P.—A mixture of alkaloids from pomegranate.

A light yellow, odorless powder with an astringent taste. Slightly soluble in water (1:240) and soluble in alcohol (1:16).

ACTION AND USES: Anthelmintic and teniafuge.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.). May be administered suspended in water. Should be given on an empty stomach followed after one or two hours by brisk purging. Not more than 0.3 Gm. or 5 grains should be given.

Pepo (Pepo), *Pepo*, U. S. P. (Pumpkin Seed).

ACTION AND USES: Domestic anthelmintic; probably inefficient.

DOSAGE: 30 Gm. or 1 ounce (U. S. P.).

**Pepsinum* (Pepsin.), *Pepsin*, U. S. P.—Contains a proteolytic ferment or enzyme from the stomach of the hog.

White or yellowish scales, grains, or powder, having a slightly acid or saline taste and no offensive odor. Soluble in water (1:50).

nearly insoluble in alcohol. Digests 3000 times its own weight of coagulated egg albumen.

ACTION AND USES: Used to assist in the gastric digestion of proteins. Usually superfluous, since gastric juice generally contains sufficient pepsin to perform gastric digestion. The alcohol of the elixirs and wines may be distinctly harmful in gastric disorders, while the other ingredients of the many complex mixtures need be prescribed only when direct indications for them exist. Many of these mixtures are irrational.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Elixir Cinchonae Alkaloidorum, Ferri et Pepsini (*Elix. Cinchon. Ferr. et Pepsin.*), *Elixir of Cinchona Alkaloids, Iron and Pepsin*, N. F. (*Elixir Cinchonae, Ferri et Pepsini*, N. F. III, *Elixir of Calisaya, Alkaloidal with Iron and Pepsin*).—Glycerite of pepsin (20%) and elixir of cinchona alkaloids and iron. Absolute alcohol content about 15 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Pepsini (*Elix. Pepsin.*), *Elixir of Pepsin*, N. F.—Glycerite of pepsin (20%), glycerin (10%), hydrochloric acid (0.4%) and aromatic elixir. Absolute alcohol content about 15 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Pepsini, Bismuthi et Strychninae.—See under *Strychnina*.

Elixir Pepsini et Bismuthi (*Elix. Pepsin. et Bism.*), *Elixir of Pepsin and Bismuth*, N. F.—Pepsin (0.85%), glycerin (12.5%), glycerite of bismuth (12.5%) in water and aromatic elixir. Absolute alcohol content about 10 per cent.

USES: This preparation lacks the protective action of the basic bismuth salts.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Pepsini et Ferri (*Elix. Pepsin. et Ferr.*), *Elixir of Pepsin and Iron*, N. F.—Tincture of ferric citro-chloride (7.5%) and elixir of pepsin. Absolute alcohol content about 15 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Pepsini et Rennini Compositum (*Elix. Pepsin. et Rennin. Co.*), *Compound Elixir of Pepsin and Rennin*, N. F. (*Essentia Pepsini*, N. F. III).—Pepsin (2.25%), rennin (1.65%), lactic acid, tincture of sweet orange peel, glycerin, alcohol, oil of myristica and water. Absolute alcohol content about 19 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Glyceritum Pepsini (*Glycer. Pepsin.*), *Glycerite of Pepsin*, N. F.—Pepsin (8.5%), hydrochloric acid (1%) in glycerin and water.

DOSAGE: 3 Cc. or 45 minims (N. F.).

Liquor Pepsini (*Liq. Pepsin.*), *Solution of Pepsin*, N. F.—Glycerite of pepsin (5%) and hydrochloric acid (1%) in glycerin and water.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Liquor Pepsini Antisepticus (*Liq. Pepsin. Antisept.*), *Antiseptic Solution of Pepsin*, N. F.—Pepsin (5%), menthol, eucalyptol, methyl salicylate, alcohol, glycerin and diluted hydrochloric acid in water. Absolute alcohol content about 1 per cent.

USES: This solution appears misnamed, since its administration cannot render the stomach antiseptic.

Liquor Pepsini Aromaticus (*Liq. Pepsin. Arom.*), *Aromatic Solution of Pepsin*, N. F.—Pepsin (1.75%), oils of cinnamon, pimenta and clove, alcohol, hydrochloric acid and glycerin in water. Absolute alcohol content about 3 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Pepsinum Saccharatum (*Pepsin. Sacchar.*), *Saccharated Pepsin*, N. F.—Pepsin (10%) with sugar of milk.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Succus Citri et Pepsinum (*Suc. Cit. et Pepsin.*), *Lime Juice and Pepsin*, N. F. (*Succus Limettae cum Pepsino*, N. F. III).—Glycerite of pepsin (40%) and lime juice.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Vinum Pepsini (*Vin. Pepsin.*), *Wine of Pepsin*, N. F. (*Pepsin Wine*).—Glycerite of pepsin (20%) in alcohol and sherry wine. Absolute alcohol content about 23 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Persio (**Persio**), **Cudbear**, N. F. (Red Indigo).—A purplish-red powder prepared from lichens.

ACTION AND USES: Coloring agent.

Tinctura Persionis (*Tr. Persion.*), *Tincture of Cudbear*, N. F.—Cudbear in alcohol and water.

USES: To impart a reddish color.

Tinctura Persionis Composita (*Tr. Persion. Co.*), *Compound Tincture of Cudbear*, N. F.—Cudbear and caramel in alcohol and water.

USES: To impart a reddish-brown color.

- ***Petrolatum** (*Petrolat.*), **Petrolatum**, U. S. P. (*Petrolatum Ointment*, *Petroleum Jelly*).—A purified mixture of semi-solid hydrocarbons from crude petroleum.

An unctuous, nearly odorless and nearly tasteless semisolid. Insoluble in water and almost insoluble in alcohol.

ACTION AND USES: Protective to the skin and basis for ointments, etc.

Emulsum Petrolati (*Emul. Petrolat.*), *Emulsion of Petrolatum*, N. F. (*Emulsum Petrolei*, N. F. III).—Petrolatum (22.5%), expressed oil of almond and acacia in syrup and water flavored with tincture of lemon peel.

USES: Emollient laxative.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Petrolatum Album (*Petrolat. Alb.*), **White Petrolatum**, U. S. P. (*White Petroleum Jelly*).—Petrolatum decolorized or nearly so.

ACTION AND USES: Same as those of petrolatum.

- ***Petrolatum Liquidum** (*Petrolat. Liq.*), **Liquid Petrolatum**, U. S. P. (*Liquid Paraffin*, *Mineral Oil*).—A mixture of liquid hydrocarbons. Official as Heavy Liquid Petrolatum and Light Liquid Petrolatum.

A colorless, transparent, oily, nearly odorless and nearly tasteless liquid. Insoluble in water or alcohol.

ACTION AND USES: Vehicle for medicinal agents to be applied externally or to the mucous membranes of the nose and throat. Also given internally largely for its mechanical action in constipation. Is not absorbed by the intestine and has no nutritive properties.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Nebula Aromatica (*Nebul. Arôm.*), *Aromatic Oil Spray*, N. F. (*Aromatol*).—Light liquid petrolatum aromatized with phenol, menthol, thymol, camphor, benzoic acid, eucalyptol, oil of cinnamon, oil of clove and methyl salicylate.

USES: An aromatic spray.

Petroxolinum Liquidum (*Petrox. Liq.*), *Liquid Petroxolin*, N. F. (*Petrolatum Saponatum Liquidum*, N. F. III, *Liquid Petrox.*).—Light liquid petrolatum and ammonium oleate, scented with oil of lavender and diluted with water and alcohol.

USES: A soap-emulsion of petrolatum intended as vehicle for oily applications.

Petroxolinum Spissum (*Petrox. Spiss.*), *Solid Petroxolin*, N. F. (*Petrolatum Saponatum Spissum*, N. F. III, *Solid Petrox.*).—An ointment composed of ammonium oleate, white wax, light liquid petrolatum and alcohol scented with oil of lavender.

Petroselini Radix (*Petrosel. Rad.*), *Parsley Root*, N. F.

ACTION AND USES: Obsolete carminative.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Petroselini Radicis (*Fldext. Petrosel. Rad.*), *Fluidextract of Parsley Root*, N. F.—Parsley root (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Petroselinum (*Petrosel.*), *Parsley Fruit*, U. S. P. (*Parsley Seed*).

ACTION AND USES: See the oleoresin below.

Oleoresina Petroselini (*Oleores. Petrosel.*), *Oleoresin of Parsley Fruit*, U. S. P. (*Liquid Apiol*).

USES: Advertised as an emmenagogue, but of doubtful value.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

PETROXOLINA, PETROXOLINS.—The petroxolins are solutions or suspensions of active medicaments in a mixture of ammonia soap and petrolatum or liquid petrolatum. Their advantage over other forms of external applications is doubtful. For official petroxolins see under the names of the principal ingredients.

***Phenol** (*Phenol*), *Phenol*, U. S. P. (*Carbolic Acid*).—Hydroxybenzene obtained from coal-tar or made synthetically.

Colorless crystals or white crystalline masses, sometimes becoming reddish, with a characteristic odor. Soluble in water (1:15), and very soluble in alcohol, glycerin or the fixed or volatile oils.

ACTION AND USES: Standard antiseptic and germicide; also local anesthetic and caustic.

Aqua Phenolata (*Aq. Phenol.*), *Phenolated Water*, N. F. (*Carbolic Acid Water*).—Phenol (2%) in water.

Glyceritum Phenolis (*Glycer. Phenol.*), *Glycerite of Phenol*, U. S. P. (*Glycerite of Carbolic Acid*).—Liquefied phenol (20%) in glycerin.

DOSAGE: 0.3 Cc. or 5 minims (U. S. P.).

Oleum Phenolatum (*Ol. Phenol.*), *Phenolated Oil*, N. F. (*Oleum Carbolatum*, N. F. III, *Carbolized Oil*).—Phenol (5%) in olive oil.

USES: A mild local stimulant and anesthetic but practically not antiseptic.

Petroxolinum Phenolis (*Petrox. Phenol.*), *Phenol Petroxolin*, N. F. (Phenol Petrox).—Phenol (5%) in liquid petroxolin.

Petroxolinum Phenolis Camphoratum (*Petrox. Phenol. Camph.*), *Camphorated Phenol Petroxolin*, N. F. (Camphorated Phenol Petrox).—Phenol (12.5%) and camphor (37.5%) in liquid petroxolin.

**Phenol Liquefactum* (*Phenol Liq.*), *Liquefied Phenol*, U. S. P. (Liquefied Carbolic Acid).—Phenol liquefied by water (about 10%).

A colorless liquid, which may become reddish.

DOSAGE: 0.05 Cc. or 1 minim (U. S. P.).

Unguentum Phenolis (*Ung. Phenol.*), *Ointment of Phenol*, U. S. P. (Ointment of Carbolic Acid).—Liquefied phenol (2.25%) in ointment.

Phenol Iodatum (*Phenol Iodat.*), *Iodized Phenol*, N. F. (*Acidum Carbolicum Iodatum*, N. F. III).—Iodine (20%) and phenol (60%) in glycerin.

**Phenolphthaleinum* (*Phenolphthal.*), *Phenolphthalein*, U. S. P.

White or nearly white, odorless, tasteless powder. Almost insoluble in water; soluble in alcohol (1:13).

ACTION AND USES: Mild but efficient cathartic. It may cause some irritation to the rectum and lower bowel and very large doses have produced alarming secondary symptoms.

DOSAGE: 0.15 Gm. or 2½ grains (U. S. P.).

Trochisci Phenolphthaleini (*Troch. Phenolphthal.*), *Troches of Phenolphthalein*, N. F.—Each troche contains phenolphthalein (0.06 Gm. or 1 grain) with acacia, sugar, vanillin and carmine.

DOSAGE: 1 troche.

**Phenylis Salicylas* (*Phenyl. Salicyl.*), *Phenyl Salicylate*, U. S. P. (*Salol*).

White powder, with an aromatic odor and a slight, characteristic taste. Slightly soluble in water (1:6670); freely soluble in alcohol (1:6), and in fixed or volatile oils.

ACTION AND USES: Intestinal antiseptic, by the liberation of phenol and salicylic acid.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.) best in powder; may be inclosed dry in capsules or cachets.

***Phosphorus (Phosphor.), Phosphorus, U. S. P.**

Translucent, nearly colorless, waxy solid with a characteristic odor, becoming white, red or black on the surface on long keeping. Readily inflammable in air (preserved under water). Almost insoluble in water, to which phosphorus imparts its characteristic odor and taste; slightly soluble in dehydrated alcohol (1:400) and soluble in chloroform (1:17).

ACTION AND USES: In rickets for stimulation of bone formation. Its use as "nerve-food" is irrational.

DOSAGE: 0.5 mg. or $\frac{1}{420}$ grain (U. S. P.). May be administered in pills or as phosphorated oil (1:100). The preparations are unreliable unless they have been recently made, since phosphorus rapidly loses its efficiency through oxidation.

Elixir Phosphori (Elix. Phosphor.), Elixir of Phosphorus, N. F.—Phosphorus (0.025%) and chloroform in alcohol, glycerin and water flavored with compound spirit of orange and oil of anise. Absolute alcohol content about 32 per cent. Deteriorates rapidly.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Phosphori et Nucis Vomicae (Elix. Phosphor. et Nuc. Vom.), Elixir of Phosphorus and Nux Vomica, N. F.—Tincture of nux vomica (3.5%) and elixir of phosphorus. Absolute alcohol content about 32 per cent.

USES: An irrational "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Liquor Phosphori (Liq. Phosphor.), Solution of Phosphorus, N. F. (Thompson's Solution of Phosphorus).—Phosphorus (0.07%) in glycerin and dehydrated alcohol flavored with spirit of peppermint. Absolute alcohol content about 35 per cent.

DOSAGE: 0.6 Cc. or 10 minims (N. F.).

Oleum Phosphoratum (Ol. Phosphorat.), Phosphorated Oil, N. F.—Phosphorus (1%), dissolved in ether and added to expressed oil of almond.

DOSAGE: 0.05 Cc. or 1 minim (N. F.).

Pilulae Phosphori (Pil. Phosphor.), Pills of Phosphorus, U. S. P.—Each pill should contain phosphorus (0.6 mg. or $\frac{1}{400}$ grain) coated with tolu.

DOSAGE: 1 pill (U. S. P.).

Physostigma (Physostig.), Physostigma, U. S. P. (Calabar Bean, Ordeal Bean).—Yields not less than 0.15 per cent. of alkaloids (chiefly physostigmine).

ACTION AND USES: Antitetic, antineuralgic, peristaltic. See *Physostigminae Salicylas*, which has the same actions

and deserves preference because of its uniform composition.

DOSAGE: 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.). Not used as such.

Extractum Physostigmatis (Ext. Physostig.), *Extract of Physostigma*, U. S. P. (Powdered Extract of Physostigma).—One Gm. extract represents about 13 Gm. of physostigma. Yields about 2 per cent. of alkaloids.

DOSAGE: 0.008 Gm. or $\frac{1}{8}$ grain (U. S. P.).

Tinctura Physostigmatis (Tr. Physostig.), *Tincture of Physostigma*, U. S. P. (Tincture of Calabar Bean).—Physostigma (10%), yielding about 0.015 per cent. of alkaloids. Absolute alcohol content about 92 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

***Physostigminae Salicylas (Physostig. Salicyl.), Physostigmine Salicylate, U. S. P. (Eserine Salicylate).**

Colorless or nearly colorless, odorless crystals. Soluble in water (1:75) and in alcohol (1:16).

ACTION AND USES: Stimulant of parasympathetic nerve endings. Used especially as a powerful miotic, particularly in glaucoma, sometimes internally against ileus or intestinal paresis.

DOSAGE: Internally, 0.001 Gm. or $\frac{1}{60}$ grain (U. S. P.). Externally, solutions (should be freshly prepared) from 0.1 to 1 per cent.

Phytolacca (Phytolac.), Phytolacca, N. F. (Poke Root).

ACTION AND USES: Unreliable irritant, emetic, cathartic and "alterative."

DOSAGE: Emetic, 1 Gm. or 15 grains (N. F.); alterative, 0.1 Gm. or $1\frac{1}{2}$ grains (N. F.).

Fluidextractum Phytolaccae (Fldext. Phytolac.), *Fluid-extract of Phytolacca*, N. F.—Phytolacca (100%). Absolute alcohol content about 40 per cent.

DOSAGE: Emetic, 1 Cc. or 15 minims (N. F.); alterative, 0.1 Cc. or $1\frac{1}{2}$ minims (N. F.).

***Pilocarpinae Hydrochloridum (Pilocarpin. Hydrochl.), Pilocarpine Hydrochloride, U. S. P. (Pilocarpine Chloride).**

Colorless, odorless crystals with a faintly bitter taste. Very soluble in water (1:0.3) and freely soluble in alcohol (1:3).

ACTION AND USES: Used especially to increase sweat *other secretions*, also as miotic. Must be used with caution when the circulation is weak.

DOSAGE: By mouth 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.); hypodermic, 0.005 Gm. or $\frac{1}{12}$ grain (U. S. P.). Best administered by hypodermic injection.

***Pilocarpinae Nitras (Pilocarpin. Nit.), Pilocarpine Nitrate, U. S. P.**

Shining, odorless crystals. Freely soluble in water (1:4) and soluble in alcohol (1:75).

ACTION AND USES: Has no advantage over pilocarpine hydrochloride.

DOSAGE: By mouth, 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.); hypodermic, 0.005 Gm. or $\frac{1}{12}$ grain (U. S. P.).

Pilocarpus (Pilocarp.), Pilocarpus, U. S. P. (Jaborandi).—

Leaves, yielding 0.6 per cent. of alkaloids (chiefly pilocarpine).

ACTION AND USES: See Pilocarpine.

DOSAGE: 2 Gm. or 30 grains.

***Fluidextractum Pilocarpi (Fldext. Pilocarp.), Fluidextract of Pilocarpus, U. S. P.* (Fluidextract of Jaborandi).—**

Pilocarpus (100%), yielding about 0.6 per cent. of alkaloids. Absolute alcohol content about 55 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

PILULAE, PILLS.—Pills are dosage forms of medicinal substances intended for administration by the mouth. For the official pills see under the names of the principal ingredients. For Pilulae ad Prandium see under **Aloe**; for Pilulae Antidyspepticae, under **Strychnina**; for Pilulae Antimonii Compositae, under **Antimonium Sulphuratum**; for Pilulae Antiperiodicae and Pilulae Antiperiodicae sine Aloe, under **Quininae Sulphas**; for Pilulae Catharticae Compositae, Pilulae Catharticae Vegetabiles and Pilulae Laxativae Post Partum, under **Colocynthis**; for Pilulae Ferri, Quininae, Aloes et Nucis Vomicae, under **Ferri Sulphas**; for Pilulae Ferri, Quininae Strychninae et Arseni Fortiores and Pilulae Ferri, Quininae, Strychninae et Arseni Mites, under **Ferri Reductum**; for Pilulae Laxativae Compositae, under **Aloinum**.

Pimenta (Piment.), Pimenta, N. F. (Pimento, Allspice).—
Nearly ripe fruit.

ACTION AND USES: Aromatic; stomachic; has no advantage over cardamom.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Pimpinella (Pimpinell.), Pimpinella, N. F. (Pimpernel Root).
—Rhizome and roots.

ACTION AND USES: Formerly used as a diuretic, dia-phoretic and stomachic. Now obsolete; no definite indications for its use.

DOSAGE: 1 Gm. or 15 grains.

Tinctura Pimpinellae (Tr. Pimpinell.), Tincture of Pimpinella, N. F.—Pimpinella (20%) in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Pinus Alba (Pinus Alb.), White Pine Bark, N. F.—Inner bark.

ACTION AND USES: Used as "expectorant" in proprietary cough syrups. Of doubtful value.

DOSAGE: 2 Gm. or 30 grains.

Syrupus Pini Strobi Compositus (Syr. Pin. Strob. Co.), Compound Syrup of White Pine, N. F.—White pine bark and wild cherry (each 8.5%), aralia (1%), balsam poplar buds (1%), sanguinaria (0.8%), sassafras (0.7%) and chloroform (0.6%) in glycerin, alcohol and syrup, colored with cudbear and flavored with oil of sassafras. Absolute alcohol content about 12 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Pini Strobi Compositus cum Morphina.—See under Morphinae Sulphas.

Piper (Piper), Pepper, U. S. P. (Black Pepper).

ACTION AND USES: Sometimes used as carminative, generally in the form of the oleoresin, which is highly irritant.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Oleoresina Piperis (Oleores. Piper.), Oleoresin of Pepper, U. S. P.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

***Pix Liquida (Pix Liq.), Tar, U. S. P. (Pine Tar).**—Obtained by the destructive distillation of pine wood.

A semiliquid viscid blackish-brown substance, with an empyreu-matic odor and taste. Slightly soluble in water; miscible with alcohol.

ACTION AND USES: Externally, as antiseptic and stimulant in chronic diseases of the skin associated with thickening and scaling. Internally, as irritant expectorant in subacute bronchitis.

DOSAGE: Internally, 0.5 Gm. or 8 grains (U. S. P.) as syrup. Externally, in ointment, at first from 2 to 4 per cent., later in higher strength.

Glyceritum Picis Liquidæ (*Glycer. Pic. Liq.*), *Glycerite of Tar*, N. F.—Tar (6.3%) in glycerin, alcohol and water. Absolute alcohol content about 12 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Liquor Picis Alkalinus (*Liq. Pic. Alk.*), *Alkaline Solution of Tar*, N. F.—Tar (25%), in potassium hydroxide (12.5%), and water.

Syrupus Picis Liquidæ (*Syr. Pic. Liq.*), *Syrup of Tar*, U. S. P.—Tar (0.5%) in alcohol, mixed with syrup. Absolute alcohol content about 5 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

**Unguentum Picis Liquidæ* (*Ung. Pic. Liq.*), *Tar Ointment*, U. S. P.—Tar (50%), yellow wax and lard.

Vinum Picis (*Vin. Pic.*), *Wine of Tar*, N. F.—Tar (10%) in alcohol and sherry wine. Absolute alcohol content about 29 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Pix Lithanthracis (**Pix Lith.**), **Coal Tar**, N. F. (**Pix Carbonis**).—Obtained by the destructive distillation of coal.

Nearly black semisolid, with a characteristic naphthalene-like odor and a sharp, burning taste. Only slightly soluble in water; partially dissolved by alcohol.

ACTION AND USES: Antiseptic and irritant; used in skin diseases in the form of:

Liquor Picis Carbonis (*Liq. Pic. Carbon.*), *Coal Tar Solution*, N. F.—Coal tar (20%), quillaja and alcohol. Absolute alcohol content about 75 per cent.

***Plumbi Acetas** (**Plumb. Acet.**), **Lead Acetate**, U. S. P. (**Sugar of Lead**).— $\text{Pb}(\text{CH}_3\text{COO})_2 + 3\text{H}_2\text{O}$.

Colorless, shining transparent crystals or heavy white crystalline masses, efflorescent, with a faint vinegar odor and a sweetish, astringent, afterward metallic taste. Freely soluble in water (1:1.4) and soluble in alcohol (1:38).

ACTION AND USES: Seldom used internally for fear of lead poisoning. Externally, used as astringent lotion.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Lotio Plumbi et Opii (*Lot. Plumb. et Opii*), *Lotion of Lead and Opium*, N. F. (**Lead and Opium Wash**).—A mixture of lead acetate (1.75%) and tincture of opium (3.5%) in water.

USES: A popular but irrational application, astringent and protective, but not a true anesthetic. The opium alkaloids are not absorbed to any appreciable extent through the unbroken skin.

Plumbi Carbonas (Plumb. Carb.), Lead Carbonate, N. F. (White Lead).—A mixture of lead carbonate and hydroxide.

Heavy, white, odorless, tasteless powder. Insoluble in alcohol or water.

ACTION AND USES: External protective and astringent, without marked advantage over zinc oxide or the bismuth preparations, but involving some danger of lead poisoning.

Plumbi Iodidum (Plumb. Iod.), Lead Iodide, N. F.— PbI_2 .

Heavy, bright yellow, odorless, tasteless powder. Very slightly soluble in water (1:1,300), and in alcohol.

ACTION AND USES: Has been used externally in the form of an ointment for application to glandular enlargement and for chronic joint enlargement. Its use is irrational.

Unguentum Plumbi Iodidi (Ung. Plumb. Iod.), Ointment of Lead Iodide, N. F.—Lead iodide (10%), and benzoinated lard.

Plumbi Oxidum (Plumb. Oxid.), Lead Oxide, U. S. P. (Litharge).— PbO .

Heavy yellowish or reddish-yellow, odorless and tasteless powder or minute scales. Almost insoluble in water; insoluble in alcohol.

ACTION AND USES: Used to make pharmaceutical preparations.

Emplastrum Plumbi (Emp. Plumb.), Lead Plaster, U. S. P. (Diachylon Plaster).—Lead oleate obtained by heating a mixture of lead oxide, olive oil and lard.

USES: Used in plaster masses and as a basis for diachylon ointment.

Unguentum Diachylon (Ung. Diachyl.), Diachylon Ointment, U. S. P.—Lead plaster (50%), oil of lavender and white petrolatum.

Plumbi Oxidum Rubrum (Plumb. Oxid. Rub.), Red Oxide of Lead, N. F. (Red Lead).—Chiefly Pb_3O_4 .

Heavy, odorless, tasteless red powder. Almost insoluble in alcohol.

ACTION AND USES: Similar to those of Plum

Plumbi Subacetat, Lead Subacetate.

ACTION AND USES: Astringent; used externally in the form of the following preparations:

Ceratum Plumbi Subacetatis (*Cerat. Plumb. Subacet.*), *Cerate of Lead Subacetate*, N. F. (Goulard's Cerate).—A mixture of solution of lead subacetate (20%), camphor (2%), wool fat, white wax, and white petrolatum.

**Liquor Plumbi Subacetatis* (*Liq. Plumb. Subacet.*), *Solution of Lead Subacetate*, U. S. P. (Goulard's Extract).—Contains lead subacetate corresponding to about 18 per cent. of lead, Pb.

Liquor Plumbi Subacetatis Dilutus (*Liq. Plumb. Subacet. Dil.*), *Diluted Solution of Lead Subacetate*, U. S. P. (Lead Water).—Solution of lead subacetate (4%) with water.

***Podophyllum** (*Podoph.*), **Podophyllum**, U. S. P. (Mandrake, May Apple Rhizome).—Yields not less than 3 per cent. of resin.

ACTION AND USES: See the resin, below.

Extractum Podophyllum (*Ext. Podophyll.*), *Extract of Podophyllum*, N. F.—A hydro-alcoholic extract.

DOSAGE: 0.015 Gm. or $\frac{1}{4}$ grain (N. F.).

Fluidextractum Podophylli (*Fldext. Podophyll.*), *Fluid-extract of Podophyllum*, U. S. P.—Podophyllum (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

**Resina Podophylli* (*Res. Podoph.*), *Resin of Podophyllum*, U. S. P. (Podophyllin).—The resin from podophyllum.

USES: Slow but effective and rather irritant cathartic.

DOSAGE: 0.01 Gm. or $\frac{1}{6}$ grain (U. S. P.).

Populi Gemmae (*Pop. Gem.*), **Balsam Poplar Buds**, N. F. (Balm of Gilead Buds).

ACTION AND USES: Similar to but without advantage over cubeb or turpentine. Used almost exclusively in proprietary medicines and imitations.

Potassae Chlorinata, Chlorinated Potassa.

Liquor Potassae Chlorinatae (*Liq. Pot. Chlorinat.*), *Solution of Potassium Chlorate*, U. S. P. (Liquor Potassae Chlorate, carbonate (5.8%), chlor-

of Liquor Sodae
ata.

Potassa Sulphurata (Pot. Sulphurat.), Sulphurated Potassa, U. S. P. (Liver of Sulphur).—Chiefly potassium polysulphides and potassium thiosulphate.

Irregular pieces, at first liver-brown, later greenish-yellow to gray, with a strong hydrogen sulphide odor and a bitter, acrid, alkaline taste. Very soluble in water; alcohol dissolves only the sulphides. Incompatible with acids and the salts of the heavy metals.

ACTION AND USES: Employed as a parasiticide and to stimulate and soften the skin in chronic cutaneous diseases.

***Potassii Acetas (Pot. Acet.), Potassium Acetate, U. S. P.— KCH_3COO .**

White, odorless or nearly odorless powder or crystalline masses with a warming, saline taste. Freely soluble in water (1:0.5) and in alcohol (1:2.9).

ACTION AND USES: Systemic alkali and diuretic, practically equivalent to sodium acetate. It can easily be administered in water and milk, so that there appears to be little excuse for the alcoholic elixirs.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Potassii Acetatis (Elix. Pot. Acet.), Elixir of Potassium Acetate, N. F.—Potassium acetate (8.5%), and aromatic elixir. Absolute alcohol content about 20 per cent.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Elixir Potassii Acetatis et Juniperi (Elix. Pot. Acet. et Junip.), Elixir of Potassium Acetate and Juniper, N. F.—Potassium acetate (8.5%), fluidextract of juniper (12.5%) and aromatic elixir. Absolute alcohol content about 23 per cent.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Potassii Arsenas, Potassium Arsenate.

Liquor Arsenicalis Clemens (Liq. Ars. Clemens), Clemens' Solution of Arsenic, N. F. (Liquor Potassi Arsenatis et Bromidi, N. F. III, Solution of Potassium Arsenate and Bromides).—Contains potassium arsenate corresponding to about 1 per cent. of arsenic trioxide, As_2O_3 .

USES: A superfluous arsenic solution.

DOSAGE: 0.2 Cc. or 3 minims (N. F.).

Potassii Arsenis, Potassium Arsenite.

**Liquor Potassii Arsenitis (Liq. Pot. Arsen.), Solⁿ Potassium Arsenite, U. S. P.* (Fowler's Solution Arsenicalis, Liquor Arsenicalis Fowleri, P.).—*trioxide* (1%) and potassium bicarbonate

pound tincture of lavender and water. Contains potassium arsenite corresponding to 1 per cent. of As_2O_3 .

USES: The most frequently used inorganic arsenical preparation. It has an alkaline reaction and therefore tends to precipitate alkaloids.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

***Potassii Bicarbonas (Pot. Bicarb.), Potassium Bicarbonate, U. S. P.— KHCO_3 .**

Colorless, transparent crystals or white, granular powder, odorless and having a saline, slightly alkaline taste. Freely soluble in water (1:2.8) and almost insoluble in alcohol.

ACTION AND USES: Mild alkali; similar to sodium bicarbonate, but without advantage over the latter.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Liquor Antisepticus Alkalinus (Liq. Antisept. Alk.), Alkaline Antiseptic Solution, N. F.—Potassium bicarbonate (3.2%), sodium borate (3.2%) and sodium benzoate (0.8%) with thymol, eucalyptol, oil of peppermint, methyl salicylate and cudbear in alcohol, glycerin and water.

USES: Of slight value as antiseptic, but a pleasant cleansing gargle.

***Potassi Bitartras (Pot. Bitart.), Potassium Bitartrate, U. S. P. (Cream of Tartar).**

Colorless, slightly opaque crystals or white, somewhat gritty powder, odorless and having an acidulous taste. Only slightly soluble in water (1:155) and very slightly soluble in alcohol (1:8,820).

ACTION AND USES: Diuretic and aperient. Frequently administered in combination with jalap as a hydragogue cathartic. See *Pulvis Jalapae Compositus* under Jalapa.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

***Potassii Bromidum (Pot. Brom.), Potassium Bromide, U. S. P.— KBr .**

White crystals or granular powder, odorless and having a strongly saline taste. Freely soluble in water (1:1.5); slightly soluble in alcohol (1:250).

ACTION AND USES: Nerve sedative; diminishes reflex excitability and depresses the motor area of the cortex. It is a general sedative, but more likely to disturb

the (U. S. P.); preferably in aqueous solutions; increased administration should be

interrupted occasionally to prevent bromism and accumulation in the system.

Elixir Potassii Bromidi (*Elix. Pot. Brom.*), *Elixir of Potassium Bromide*, N. F.—Potassium bromide (17.5%) in syrup, water and aromatic elixir; may be colored with compound tincture of cudbear. Absolute alcohol content about 7.5 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Sal Potassii Bromidi Effervescens (*Sal Pot. Brom. Eff.*), *Effervescent Salt of Potassium Bromide*, N. F. (*Pulvis Potassii Bromidi Effervescens*, N. F. III).—Potassium bromide (16.6%), sodium bicarbonate (53%), tartaric acid (20.4%) and citric acid, the last-named three yielding sodium citrate and sodium carbonate.

DOSAGE: 6 Gm. or 1½ drachms.

Sal Potassii Bromidi Effervescens Compositus (*Sal Pot. Brom. Eff. Co.*), *Compound Effervescent Salt of Potassium Bromide*, N. F. (*Pulvis Potassii Bromidi Effervescens cum Caffeina*, N. F. III, *Effervescent Potassium Bromide with Caffeine*).—Caffeine (0.8%) and potassium bromide (8.3%) in a mixture of sodium bicarbonate (58.7%), lithium carbonate (4.2%), tartaric acid (18%) and citric acid. (Yielding citrate and tartrate of lithium and sodium.)

USES: "Headache" mixture; generally undesirable.

DOSAGE: 6 Gm. or 1½ drachms (N. F.).

Syrupus Bromidorum (*Syr. Bromidor.*), *Syrup of the Bromides*, N. F.—Potassium bromide and sodium bromide (each 8%), ammonium bromide (5%), calcium bromide (2.5%), lithium bromide (0.8%) flavored with tincture of vanilla and colored with compound tincture of cudbear in compound syrup of sarsaparilla and syrup.

USES: It is very improbable that this complex mixture has any advantage over a simple bromide.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Potassii Carbonas* (*Pot. Carb.*), *Potassium Carbonate*, U. S. P.— K_2CO_3 . (A hydrated form is sold as salt of tartar).

White, odorless, deliquescent granular powder with a strongly alkaline taste. Very soluble in water (1:0.9), but insoluble in alcohol.

ACTION AND USES: Strongly alkaline and caustic; very dilute solutions sometimes used as antacid.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

***Potassii Chloras (Pot. Chloras), Potassium Chlorate, U. S. P.—KClO₃.**

Colorless, lustrous crystals or white, granular powder, odorless and having a saline taste. Soluble in water (1:11.5) and almost insoluble in alcohol.

ACTION AND USES: Used as mouthwash and gargles in stomatitis, etc. Large doses are actively poisonous, causing disintegration of the blood corpuscles. Has been used in the form of gargle and troches for sore throat (swallowing the latter may be dangerous).

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Trochisci Potassii Chloratis (Troch. Pot. Chlorat.), Troches of Potassium Chlorate, U. S. P.—Each troche contains potassium chlorate (0.15 Gm. or 2½ grains) with sugar and tragacanth.

Potassii Chloridum (Pot. Chlorid.), Potassium Chloride, N. F.—KCl.

Colorless crystals or white powder, odorless and having a saline taste. Freely soluble in water (1:2.8) and insoluble in alcohol.

ACTION AND USES: Has been recommended for use as table salt in place of sodium chloride, in cases in which the absorption of salt was thought to be undesirable. Its value is not established.

***Potassii Citras (Pot. Cit.), Potassium Citrate, U. S. P.**

Transparent crystals or white, granular powder; odorless and having a cooling saline taste. Very soluble in water (1:0.6) and almost insoluble in alcohol.

ACTION AND USES: Systemic alkali and diuretic like potassium acetate, but more laxative.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

**Potassii Citras Effervescens (Pot. Cit. Eff.), Effervescent Potassium Citrate, U. S. P.*—An effervescent mixture representing potassium citrate (20%), sodium citrate (27.55%) and sodium tartrate (38.7%).

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Liquor Potassii Citratis (Liq. Pot. Cit.), Solution of Potassium Citrate, U. S. P.—Potassium citrate (8%) in water.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

***Potassii et Sodii Tartras (Pot. et Sod. Tart.), Potassium and Sodium Tartrate, U. S. P. (Rochelle Salt).**

Transparent crystals or white powder, odorless and having a cooling, saline taste. Very soluble in water (1:0.9).

ACTION AND USES: Saline cathartic, of relatively unobjectionable taste.

DOSAGE: 10 Gm. or 2½ drachms (U. S. P.) in water.

**Pulvis Effervescens Compositus* (*Pulv. Eff. Co.*), *Compound Effervescent Powder*, U. S. P. (Seidlitz Powder).—The blue paper contains sodium bicarbonate (2.5 Gm.) and potassium and sodium tartrate (7.5 Gm.). The white paper contains tartaric acid (2.16 Gm.).

DOSAGE: 1 set of two powders (U. S. P.).

**Potassii Hydroxidum* (*Pot. Hydrox.*), *Potassium Hydroxide*, U. S. P. (Caustic Potash, Potassium Hydrate).—KOH (not less than 85%).

Dry white or nearly white, odorless, deliquescent, hard and brittle flakes, masses or sticks. Very soluble in water (1:0.9) and freely soluble in alcohol (1:3).

ACTION AND USES: Internally, an antacid, but objectionable for it is more irritant than the bicarbonate and without advantage over the latter. Externally, caustic and solvent, but too diffident.

**Liquor Potassii Hydroxidi* (*Liq. Pot. Hydrox.*), *Solution of Potassium Hydroxide*, U. S. P. (*Liquor Potassae*, *Solution of Potassa*).—KOH (not less than 4.5%).

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Potassa cum Calce (*Pot. c. Calc.*), *Potassa with Lime*, N. F.—Potassium hydroxide (50%) and calcium oxide.

USES: Caustic.

Potassii Hypophosphis (*Pot. Hypophos.*), *Potassium Hypophosphite*, U. S. P.— KH_2PO_2 . (Caution should be observed in handling this product to avoid explosion.)

White, opaque deliquescent plates, crystalline masses or granular powder, odorless and having a pungent, saline taste. Very soluble in water (1:0.6) and freely soluble in alcohol (1:9).

ACTION AND USES: Introduced in common with other hypophosphites as a cure for tuberculosis; since used as "nerve-foods." There is no proof that the hypophosphites exert any physiologic effect, influence any pathologic process, or have any food value.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

**Potassii Iodidum* (*Pot. Iod.*), *Potassium Iodide*, U. S. P.—KI.

Transparent, translucent or opaque crystals or white granular powder, with a pungent, saline, afterward bitter taste. Very soluble in water (1:0.7) and soluble in alcohol (1:22). Incompatible with

mineral acids and oxidizing agents and should not be prescribed in solution with alkaloids or alkaloid-containing drugs.

ACTION AND USES: Saline expectorant, especially in asthma and chronic bronchitis; diuretic; antisyphilitic; "alterative" in rheumatic conditions, sclerosis, etc.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.); best administered in simple solution and well diluted; in lues the dose should be gradually increased to the limit of tolerance.

Unguentum Potassii Iodidi (Ung. Pot. Iod.), Ointment of Potassium Iodide, N. F.—Potassium iodide (10%), sodium thiosulphate (1%), water and benzoated lard.

USES: An irrational method of administering potassium iodide, as none of the drug is absorbed through the skin.

Potassi Nitras (Pot. Nitras), Potassium Nitrate, U. S. P. (Saltpetre).— KNO_3 .

Transparent crystals or white powder, odorless and having a cooling saline taste. Freely soluble in water (1:2.8) and slightly soluble in alcohol (1:620).

ACTION AND USES: Diuretic, irritant to kidneys and intestines; has largely gone out of use.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Charta Potassii Nitratis (Chart. Pot. Nit.), Potassium Nitrate Paper, N. F.—Paper impregnated with potassium nitrate.

USES: The smoke is inhaled in asthma, but is not very effective.

***Potassii Permanganas (Pot. Permang.), Potassium Permanganate, U. S. P.**— KMnO_4 .

Slender, dark purple odorless crystals, with a taste (in solution) at first sweet, afterward disagreeable and astringent. Soluble in water (1:13.5), forming violet-red to rose-colored solutions which stain the fingers and clothes. Decomposed by alcohol and glycerin. Should not be brought into contact with organic or other readily oxidizable substances.

ACTION AND USES: Effective deodorant, disinfectant and astringent. Internally as emmenagogue, probably useless.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.), usually in pills. For application to the skin, 1:500 solution.

Potassii Sulphas (Pot. Sulph.), Potassium Sulphate, N. F.— K_2SO_4 .

Colorless crystals or white powder, odorless and having a saline, somewhat bitter taste. Freely soluble in water (1:10) and insoluble in alcohol.

ACTION AND USES: Saline purgative closely resembling sodium sulphate in its action, but without advantage over the latter.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Prunum (Prunum), Prune, N. F.

ACTION AND USES: One of the ingredients in confection of senna but seldom used in medicine apart from this.

***Prunus Virginiana (Prun. Virg.), Wild Cherry, U. S. P. (Wild Black Cherry Bark).**

ACTION AND USES: Sedative expectorant, mainly used as flavor.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Fluidextractum Pruni Virginianae (Fldext. Prun. Virg.), Fluidextract of Wild Cherry, N. F.—Wild cherry (100%). Absolute alcohol content about 20 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Infusum Pruni Virginianae (Inf. Prun. Virg.), Infusion of Wild Cherry, N. F.—Wild cherry (4%), in glycerin and water.

DOSAGE: 60 Cc. or 2 fluidounces (N. F.).

***Syrupus Pruni Virginianae (Syr. Prun. Virg.), Syrup of Wild Cherry, U. S. P.**—Wild cherry bark (15%), in glycerin and syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Vinum Pruni Virginianae (Vin. Prun. Virg.), Wine of Wild Cherry, N. F.—Wild cherry (25%) in water, alcohol, sugar and sherry wine. Absolute alcohol content about 23 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Vinum Pruni Virginianae Ferratum (Vin. Prun. Virg. Ferr.), Ferrated Wine of Wild Cherry, N. F.—Tincture of ferric citrochloride (8%) in wine of wild cherry. Absolute alcohol content about 21 per cent.

USES: Superfluous "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Pulsatilla (Pulsatil.), Pulsatilla, N. F. (Pasque Flower, Meadow Anemone).—The plant.

ACTION AND USES: Has been recommended as an alterative, sedative and antispasmodic. Has been used in syphilis with undoubted harmful results because of its inefficiency. Its use is fortunately decreasing.

DOSAGE: 0.3 Gm. or 5 grains (N. F.).

Tinctura Pulsatillae (Tr. Pulsatil.), *Tincture of Pulsatilla*, N. F.—Pulsatilla (10%) in alcohol and water. Absolute alcohol content about 65 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

PULVERES, POWDERS.—The official powders are mixtures of several substances in powdered form. For the preparations now official see under the names of the principal constituents. For Pulvis Antimonialis, see under *Antimonii Oxidum*; for Pulvis Antisepticus, see under *Zinci Sulphas*; for Pulvis Aromaticus and Pulvis Cretae Aromaticus, under *Cinnamomum Saigoncum*; for Pulvis Aromaticus Rubefaciens, under *Caryophyllus*; Pulvis Cretae Aromaticus et Opii, under *Opium*; for Pulvis Effervescens Compositus, under *Potassii et Sodii Tartras*; for Pulvis Glycyrrhizae Compositus, under *Senna*; for Pulvis Ipecacuanhae et Opii, under *Opium*, and for Pulvis Talci Compositus, under *Acidum Salicylicum*.

Pumex (Pumex), Pumice, N. F.—Consists chiefly of complex silicates of volcanic origin.

ACTION AND USES: Used as an abrasive material, and in pharmacy as an absorbent.

Pyrethrum (Pyreth.), Pyrethrum, U. S. P. (Pellitory Root).

ACTION AND USES: Obsolete, irritant sialagogue and local irritant.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Tinctura Pyrethri (Tr. Pyreth.), *Tincture of Pyrethrum*, U. S. P. (Tincture of Pellitory).—Pyrethrum (20%) in alcohol. Absolute alcohol content about 90 per cent.

USES: External.

Pyrogallol (Pyrogall.), Pyrogallol, U. S. P. (Pyrogallic Acid).—Trihydroxybenzene.

Light, white or nearly white, odorless, bitter laminae, fine needles or powder. Freely soluble in water (1:1.7) and in alcohol (1:1.3).

ACTION AND USES: Irritant antiseptic in chronic skin diseases. Internally highly toxic.

DOSAGE: 5 per cent. ointment.

Pyroxylinum (Pyroxylin), Pyroxylin, U. S. P. (Soluble Gun Cotton).—Chiefly cellulose tetranitrate.

A yellowish-white matted mass of filaments, resembling raw cotton in appearance; very inflammable. Slowly but completely soluble in a mixture of three volumes of ether and one volume of alcohol. Also soluble in acetone and in glacial acetic acid. Practically insoluble in alcohol or water.

ACTION AND USES: Basis for collodions.

**Collodium* (*Collod.*), *Collodium*, U. S. P.—Pyroxylin (4%) in a mixture of ether and alcohol.

USES: Used to form a protective film and as a vehicle for external applications.

**Collodium Flexile* (*Collod. Flex.*), *Flexible Collodium*, U. S. P.—A mixture of collodion (95%) with camphor (2%) and castor oil (3%).

USES: More pliable than collodion and does not contract as much in drying.

Quassia (*Quass.*), *Quassia*, U. S. P. (Bitter Wood).

ACTION AND USES: Simple bitter stomachic (more disagreeable than gentian); an infusion is also used as enema in the treatment of pinworms.

DOSEAGE: 0.5 Gm. or 8 grains (U. S. P.).

Extractum Quassiae (*Ext. Quas.*), *Extract of Quassia*, N. F. (Powdered Extract of Quassia).—An aqueous extract diluted with starch, 1 Gm. representing 10 Gm of quassia.

DOSEAGE: 0.06 Gm. or 1 grain (N. F.).

Fluidextractum Quassiae (*Flidext. Quass.*), *Fluidextract of Quassia*, N. F.—Quassia (100%). Absolute alcohol content about 25 per cent.

DOSEAGE: 0.5 Cc. or 8 minims (N. F.).

Tinctura Quassiae (*Tr. Quas.*), *Tincture of Quassia*, U. S. P.—Quassia (20%) in alcohol and water. Absolute alcohol content about 30 per cent.

DOSEAGE: 2 Cc. or 30 minims (U. S. P.).

Quercus (*Querc.*), *Quercus*, N. F. (White Oak Bark).

ACTION AND USES: Tannin-bearing astringent, without advantage over krameria, etc.

DOSEAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Quercus (*Flidext. Querc.*), *Fluidextract of Quercus*, N. F.—Quercus (100%). Absolute alcohol content about 40 per cent.

DOSEAGE: 1 Cc. or 15 minims (N. F.).

Quillaja (*Quillaja*), *Quillaja*, N. F. (Soap-tree Bark).

ACTION AND USES: Contains saponin and has been recommended as an expectorant. Similar to, but without advantage over senega. Has been used as a detergent and as an emulsifying agent; in the latter use is objectionable.

Tinctura Quillajae (Tr. *Quillaj.*), *Tincture of Quillaja*, N. F.—Quillaja (20%) in alcohol and water. Absolute alcohol content about 30 per cent.

Quinidina (Quinid.), Quinidine, N. F.—An alkaloid isomeric with quinine.

White, odorless crystals or powder, with an intense and persistent bitter taste.

ACTION AND USES: At one time recommended as a cheap substitute for quinine. Inferior and obsolete.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (N. F.); anti-malarial, 1 Gm. or 15 grains (N. F.) daily.

Syrupus Quinidinae (Syr. *Quinid.*), *Syrup of Quinidine*, N. F. (Bitterless Syrup of Quinidine).—Quinidine (3.3%) and oil of orange in syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Quinina (Quin.), Quinine, U. S. P.**—Obtained from cinchona bark.

White, odorless, intensely bitter powder. Very slightly soluble in water (1:1560), and very soluble in alcohol (1:0.8).

ACTION AND USES: Bitter tonic, analgesic and antipyretic; specific against malaria. Contraindicated in ear diseases. Idiosyncrasies are common.

The official quinine salts are more numerous than is necessary. Those that are starred would meet every indication.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.); anti-malarial, at least 1 Gm. or 15 grains daily (U. S. P.).

Oleatum Quininae (Oleat. *Quin.*), *Oleate of Quinine*, N. F.—Quinine (25%) in oleic acid.

***Quininae Bisulphas (Quin. Bisulph.), Quinine Bisulphate, U. S. P.**—Quinine acid sulphate.

Transparent or whitish, odorless, very bitter crystals. Freely soluble in water (1:9) and soluble in alcohol (1:23).

ACTION AND USES: Similar to, and without advantage over, *Quininae Dihydrochloridum*.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.); antimalarial, at least 1 Gm. or 15 grains daily (U. S. P.).

Tinctura Antiperiodica (Tr. *Antiperiod.*), *Antiperiodic Tincture*, N. F. (Warburg's Tincture).—Quinine bisulphate (2%), extract of aloes (1.75%), rhubarb (0.8%), angelica fruit, elecampane, saffron, fennel, gentian, zedoary, cubeb, myrrh, camphor, agaric, pepper, Saigon cinnamon and

jamaica ginger in alcohol and water. Absolute alcohol content about 55 per cent.

USES: Antiquated, complex and irrational quinine mixture.

DOSAGE: From 4 to 15 Cc. or 1 to 4 fluidrachms (N. F.).

Tinctura Antiperiodica sine Aloe (*Tr. Antiperiod. s. Aloe*), *Antiperiodic Tincture without Aloes*, N. F. (Warburg's Tincture without Aloes).—The same as the preceding (*Tinctura Antiperiodica*) with the omission of extract of aloes.

USES AND DOSAGE: See *Tinctura Antiperiodica*.

Quininae Dihydrochloridum (Quin. Dihydrochl.), Quinine Dihydrochloride, U. S. P.

White, odorless, very bitter powder. Very soluble in water (1:0.6), and soluble in alcohol (1:12).

ACTION AND USES: Used where concentrated solutions of quinine are wanted, as for subcutaneous injections, etc.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.); antimalarial, at least 1 Gm. or 15 grains daily (U. S. P.).

***Quininae et Ureae Hydrochloridum** (Quin. et Urea. Hydrochl.), Quinine and Urea Hydrochloride, U. S. P. (Quinine and Urea Chloride).—A double salt of quinine and urea.

Colorless, translucent crystals or white, granular powder, odorless and having a very bitter taste. Very soluble in water (1:0.9) and freely soluble in alcohol (1:2.4).

ACTION AND USES: Much less irritating than ordinary quinine salts and therefore suited for hypodermic injections, in malaria or as local anesthetic.

DOSAGE: Hypodermic, one dose daily, 1 Gm. or 15 grains (U. S. P.). For local anesthesia (hypodermic) solution of from 0.25 to 1 per cent.; (applied to mucous membranes), solutions of from 10 to 20 per cent.

Quininae Glycerophosphas (Quin. Glycerophos.), Quinine Glycerophosphate, N. F. (Quinine Glycerinophosphate).

Fine, white, odorless crystals or powder, with a very bitter taste.

ACTION AND USES: Expensive and superfluous quinine salt.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (N. F.); antimalarial, 1 Gm. or 15 grains (N. F.).

Quininae Hydrobromidum (Quin. Hydrobr.), Quinine Hydrobromide, U. S. P. (Quinine Bromide).

Light, white, odorless, very bitter silky needles. Soluble in water (1:40), and very soluble in alcohol (1:0.9).

ACTION AND USES: Superfluous; see *Quininae Hydrochloridum*.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.); antimalarial, at least 1 Gm. or 15 grains, daily (U. S. P.).

***Quininae Hydrochloridum (Quin. Hydrochl.), Quinine Hydrochloride, U. S. P. (Quinine Chloride).**

White, odorless, very bitter, silky needles. Soluble in water (1:18) and very soluble in alcohol (1:0.8).

ACTION AND USES: Has the action of other quinine salts; frequently preferred to the sulphate, because neutral and more soluble.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.); antimalarial, at least 1 Gm. or 15 grains daily (U. S. P.).

Syrupus Phosphatum cum Quinina et Strychnina (Syr. Phosphat. c. Quin. et Strych.), *Syrup of Phosphates with Quinine and Strychnine, N. F.* (Syrupus Hydrochlorophosphatum, N. F. III, Compound Syrup of Hydrochlorophosphates).—Quinine hydrochloride (0.44%), strychnine nitrate (0.014%), compound solution of phosphates (50%) in glycerin and syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Quininae Hypophosphis (Quin. Hypophos.), Quinine Hypophosphite, N. F.

Fine, white, odorless crystals, with a very bitter taste.

ACTION AND USES: Superfluous quinine salt.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (N. F.); antimalarial, 1 Gm. or 15 grains daily (N. F.).

Quininae Salicylas (Quin. Salicyl.), Quinine Salicylate, U. S. P.

Colorless (often becoming pinkish), odorless, bitter needles. Soluble in alcohol (1:14) and in glycerin (1:13). Only slightly soluble in water.

ACTION AND USES: Superfluous; see *Quininae Hydrochloridum*. The quantity of salicylate in a dose is us too small to be effective.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. antimalarial, at least 1 Gm. or 15 grains daily (U. S.

***Quininae Sulphas (Quin. Sulph.), Quinine Sulphate, 1**

White, odorless, very bitter, efflorescent crystals. Slightly soluble in water (1:725) and in alcohol (1:107).

ACTION AND USES: The most commonly used quinine salt but inferior to the hydrochloride, the dihydrochloride and quinine and urea hydrochloride where a soluble salt is desired, and to the alkaloid or the tannate where an insoluble one is needed.

DOSAGE: Tonic, 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.); antimalarial, at least 1 Gm. or 15 grains daily (U. S. P.).

Elixir Cinchonae Alkaloidorum (Elix. Cinchon. Alk.), *Elixir of Cinchona Alkaloids*, N. F. (Elixir Cinchonae, N. F. III, Elixir Calisaya, Alkaloidal).—Quinine sulphate (0.2%), cinchonidine sulphate (0.1%), cinchonine sulphate (0.1%), compound tincture of cudbear and aromatic elixir. Absolute alcohol content about 21 per cent.

USES: Imitation of cinchona, minus the tannin. No advantage over quinine.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Cinchonae Alkaloidorum et Hypophosphitum (Elix. Cinchon. et Hypophos.), *Elixir of Cinchona Alkaloids and Hypophosphites*, N. F. (Elixir Cinchonae et Hypophosphitum, N. F. III, Elixir of Calisaya, Alkaloidal, with Hypophosphites).—Calcium hypophosphite and sodium hypophosphite (each 1.75%), elixir of cinchona alkaloids (about 85%), hypophosphorous acid and water. Absolute alcohol content about 18 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Elixir Cinchonae Alkaloidorum, Ferri et Calcii Lactophosphatis.—See under *Calcii Lactophosphas*.

Pilulae Antiperiodicae (Pil. Antiperiod.), *Antiperiodic Pills*, N. F. (Warburg's Pills).—Each pill contains quinine sulphate (0.09 Gm. or $1\frac{1}{2}$ grains), extract of aloes (0.065 Gm. or 1 grain) and rhubarb (0.032 Gm. or $\frac{1}{2}$ grain) with angelica fruit, elecampane, saffron, fennel, zedoary, cubeb, myrrh, agaric, camphor and extract of gentian.

DOSAGE: 1 pill (N. F.).

Pilulae Antiperiodicae sine Aloe (Pil. Antiperiod. s. Aloe), *Antiperiodic Pills without Aloes*, N. F. (Warburg's Pills without Aloes).—Each pill contains quinine sulphate (0.09 Gm. or $1\frac{1}{2}$ grains), rhubarb (0.032 Gm. or $\frac{1}{2}$ grain) with angelica fruit, elecampane, saffron, fennel, zedoary, cubeb, myrrh, agaric, camphor and extract of gentian.

DOSAGE: 1 pill (N. F.).

***Quininae Tannas (Quin. Tann.), Quinine Tannate, U. S. P.**

Pale yellow, or yellowish-white, odorless and tasteless, or nearly tasteless powder. Only very slightly soluble in water. Somewhat more soluble in alcohol.

ACTION AND USES: Used where comparatively tasteless preparations are wanted.

DOSAGE: 0.2 Gm. or 3 grains (U. S. P.); sometimes administered with chocolate or sugar.

Trochisci Quininae Tannatis (Troch. Quin. Tan.), Troches of Quinine Tannate, N. F.—Each troche contains quinine tannate (0.06 Gm. or 1 grain) with tragacanth, oil of theobroma, cocoa, sugar and sodium benzosulphide.

DOSAGE: 1 troche (N. F.).

Quininae Valeras (Quin. Valer.), Quinine Valerate, N. F.

White, lustrous crystals, with a valerian-like odor and an intensely bitter taste.

ACTION AND USES: Those of quinine and valerates, but disagreeable and superfluous.

DOSAGE: Tonic, 0.1 Gm. or 1½ grains (N. F.); anti-malarial, 1 Gm. or 15 grains daily (N. F.).

Elixir Quininae Valeratis et Strychninae (Elix. Quin. Valer. et Strych.), Elixir of Quinine Valerate and Strychnine, N. F. (*Elixir Quininae Valerianatis et Strychninae, N. F. III.*)—Quinine valerate (1.75%), strychnine sulphate (0.0175%), compound tincture of cudbear, distilled water and aromatic elixir. Absolute alcohol content about 22 per cent.

USES: Irrational "tonic" mixture.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Renninum (Rennin.), Rennin, N. F.—Milk-curdling enzyme from the stomach of the calf.

Grayish-white or yellowish-white powder or yellowish grains or scales, with a characteristic, slightly saline taste and a peculiar odor. Slowly soluble in water and in diluted alcohol.

ACTION AND USES: For making wheys and junkets.

Resina (Resin.), Rosin, U. S. P. (Colophony).—The residue left after distilling the volatile oil from turpentine (the oleoresin).

ACTION AND USES: Used in the preparation of ointments and plasters. Rubefacient.

Ceratum Resinae (Cerat. Res.), Rosin Cerate, U. S. P. (Basilicon Ointment).—Rosin, yellow wax and lard.

Ceratum Resinae Compositum (*Cerat. Resin. Co.*), *Compound Rosin Cerate*, N. F. (Deshler's Salve).—Rosin, yellow wax, prepared suet, turpentine and linseed oil.

Emplastrum Elasticum (*Emp. Elast.*), *Rubber Plaster*, U. S. P. (Rubber Adhesive Plaster).—A mixture of rubber, resins, and waxes, with a filler of an absorbent powder.

Emplastrum Resinae (*Emp. Res.*), *Rosin Plaster*, U. S. P. (Rosin Adhesive Plaster, Adhesive Plaster).—Rosin, lead plaster and yellow wax.

RESINAE, RESINS.—Resins are the residues obtained by distilling the volatile oil from oleoresins or the precipitates obtained from solutions of resins in alcohol by the addition of water. For the official resins see under the names of the principal constituents.

***Resorcinol (Resorcin.), Resorcinol, U. S. P. (Resorcin).**—Metadihydroxybenzene.

Colorless or nearly colorless crystals or powder with a faint, peculiar odor and a taste at first sweetish, afterward bitter. Very soluble in water (1:0.9) and in alcohol (1:0.9).

ACTION AND USES: Irritant, antiseptic; used internally in gastric fermentation; externally in skin diseases.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Pasta Resorcinolis Fortis (*Past. Resorcin. Fort.*), *Strong Resorcinol Paste*, N. F. (Lassar's Stronger Resorcinol Paste).—Resorcinol (20%), zinc oxide (20%), starch and light liquid petrolatum.

Pasta Resorcinolis Mitis (*Past. Resorcin. Mit.*), *Mild Resorcinol Paste*, N. F. (Lassar's Mild Resorcinol Paste).—Resorcinol (10%), zinc oxide (25%), starch and liquid petrolatum.

Unguentum Resorcinolis Compositus (*Ung. Resorcin. Co.*), *Compound Resorcinol Ointment*, N. F.—Resorcinol, zinc oxide, bismuth subnitrate and rectified oil of birch tar (each 6%), in yellow wax, petrolatum, anhydrous wool fat and glycerin.

USES: Complex antiseptic ointment.

Rhamnus Cathartica (*Rhamn. Cath.*), *Rhamnus Cathartica*, N. F. (Buckthorn Berries, *Baccae Spinae Cervinae*).—Ripe fruit.

ACTION AND USES: Superfluous hydragogue cathartic.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Rhamni Catharticae (*Fldext. Rham. Cathart.*), *Fluidextract of Rhamnus Cathartica*, N. F.—*Rhamnus cathartica* (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Syrupus Rhamni Catharticae (*Syr. Rham. Cathart.*), *Syrup of Rhamnus Cathartica*, N. F. (*Syrup of Buckthorn Berries*, *Syrupus Spinae Cervinae*).—*Fluidextract of rhamnus cathartica* (20%), oil of fennel and oil of cinnamon, in syrup. Absolute alcohol content about 8 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

***Rheum (Rheum), Rhubarb, U. S. P.**—Rhizome and root.

ACTION AND USES: Efficient and pleasant laxative, its action resembling cascara. It is, however, also somewhat astringent. Used especially in the form of aromatic tincture and aromatic syrup.

The official preparations of rhubarb are needlessly numerous, those that are starred being quite sufficient. Many of the official mixtures and pills are unnecessarily complex.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

***Extractum Rhei** (*Ext. Rhei*), *Extract of Rhubarb*, U. S. P. (*Powdered Extract of Rhubarb*).—One Gm. extract represents 2 Gm. rhubarb.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Fluidextractum Rhei (*Fldext. Rhei*), *Fluidextract of Rhubarb*, U. S. P.—*Rhubarb* (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Fluidglyceratum Rhei (*Fldglycer. Rhei*), *Fluidglycerate of Rhubarb*, N. F.—*Rhubarb* (100%) in water and glycerin.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Mistura Rhei Alkalina (*Mist. Rhei Alk.*), *Alkaline Mixture of Rhubarb*, N. F. (*Syrupus Rhei et Potassii Compositus*, N. F. III, *Neutralizing Cordial*).—*Fluidextract of rhubarb* (1.6%), *fluidextract of hydrastis* (0.8%), and *potassium carbonate* (1.6%) with tincture of cinnamon, spirit of peppermint in syrup and diluted alcohol. Absolute alcohol content about 37 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Mistura Rhei Composita (*Mist. Rhei Co.*), *Compound Mixture of Rhubarb*, N. F. (*Mistura Rhei et Sodae*, U. S. P. VIII, *Mixture of Rhubarb and Soda*).—*Fluidextract of*

rhubarb (1.5%), fluidextract of ipecac (0.3%), sodium bicarbonate (3.5%), glycerin and spirit of peppermint in water. Absolute alcohol content about 4 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm.

Pilulae Rhei (Pil. Rhei), Pills of Rhubarb, N. F.—Each pill contains rhubarb (0.2 Gm. or 3 grains) with soap.

DOSAGE: 1 pill (N. F.).

Pilulae Rhei Compositae (Pil. Rhei Co.), Compound Pills of Rhubarb, U. S. P.—Each pill contains rhubarb (0.13 Gm. or 2 grains) and aloes (0.1 Gm. or 1½ grains) with myrrh and oil of peppermint.

DOSAGE: 2 pills (U. S. P.).

Pulvis Rhei Compositus (Pulv. Rhei Co.), Compound Powder of Rhubarb, U. S. P. (Gregory's Powder).—Rhubarb (25%), Jamaica ginger (10%) and magnesium oxide.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Pulvis Rhei et Magnesiae Anisatus (Pulv. Rhei et Magnes. Anis.), Anisated Powder of Rhubarb and Magnesia, N. F. (Compound Anise Powder).—Rhubarb (35%), anethol and heavy magnesium oxide.

USES: Mild cathartic.

DOSAGE: For infants, 0.3 Gm. or 5 grains (N. F.).

Syrupus Rhei (Syr. Rhei), Syrup of Rhubarb, U. S. P.—Fluidextract of rhubarb (10%), spirit of cinnamon and potassium carbonate in water and syrup.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

**Syrupus Rhei Aromaticus (Syr. Rhei Arom.), Aromatic Syrup of Rhubarb, U. S. P. (Spiced Syrup of Rhubarb).*—Aromatic tincture of rhubarb (15%) and potassium carbonate in syrup.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Tinctura Rhei (Tr. Rhei), Tincture of Rhubarb, U. S. P.—Rhubarb (20%) and cardamom seed, in glycerin, alcohol and water. Absolute alcohol content about 46 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Tinctura Rhei Aquosa (Tr. Rhei Ag.), Aqueous Tincture of Rhubarb, N. F.—Rhubarb (10%) and potassium carbonate in cinnamon water, alcohol and water. Absolute alcohol content about 10 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Tinctura Rhei Aromatica (Tr. Rhei Arom.), Aromatic Tincture of Rhubarb, U. S. P.*—Rhubarb (20%), Saigon

cinnamon, clove and nutmeg in glycerin, alcohol and water. Absolute alcohol content about 46 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Rhei Dulcis (*Tr. Rhei Dulc.*), *Sweet Tincture of Rhubarb*, N. F.—Rhubarb (10%), glycyrrhiza, anise and cardamom seed in glycerin, diluted alcohol and water. Absolute alcohol content about 45 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Rhei et Gentianae (*Tr. Rhei et Gent.*), *Tincture of Rhubarb and Gentian*, N. F.—Rhubarb (7%) and gentian (1.75%) in diluted alcohol. Absolute alcohol content about 45 per cent.

USES: Stomachic and laxative.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Vinum Rhei Compositum (*Vin. Rhei Co.*), *Compound Wine of Rhubarb*, N. F. (*Tinctura Rhei Vinosa*, N. F. III).—Fluidextract of rhubarb (8%), fluidextract of bitter orange peel, tincture of cardamom in sugar and sherry wine. Absolute alcohol content about 25 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Rhus Glabra (*Rhus Glab.*), *Rhus Glabra*, N. F. (Sumac Berries).

ACTION AND USES: Tannin-bearing astringent; without advantage over krameria, etc.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Rhois Glabrae (*Fldext. Rhois Glab.*), *Fluidextract of Rhus Glabra*, N. F.—Rhus glabra (100 %). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

**Rosa*, *Rose* (*Cabbage Rose*, *Rosa centifolia*).

ACTION AND USES: The preparations are used for flavoring.

**Aqua Rosae* (*Aq. Ros.*), *Rose Water*, U. S. P.—A mixture of stronger rose water with distilled water.

Aqua Rosae Fortior (*Aq. Ros. Fort.*), *Stronger Rose Water*, U. S. P.—Prepared by distilling fresh cabbage roses with water.

Confectio Rosae (*Confect. Ros.*), *Confection of Rose*, N. F.—Red rose, sugar, clarified honey and stronger rose water.

Unguentum Aquae Rosae (*Ung. Aq. Ros.*), *Ointment of Rose Water*, U. S. P. (Cold Cream).—Sodium borate (0.5%), spermaceti, white wax, expressed oil of almond, and stronger rose water.

USES: Emollient.

Rosa Gallica (*Rosa Gall.*), Red Rose, U. S. P.—Petals.

ACTION AND USES: Mildly astringent but used chiefly as a flavoring agent in the confection of rose; antiquated.

Fluidextractum Rosae (*Fldext. Rosae*), *Fluidextract of Rose*, U. S. P.—Red rose (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Infusum Rosae Compositum (*Inf. Ros. Co.*), *Compound Infusion of Rose*, N. F.—Red rose (1.3%) with diluted sulphuric acid and sugar in water.

DOSAGE: 100 Cc. or 3 fluidounces (N. F.).

Syrupus Rosae (*Syr. Ros.*), *Syrup of Rose*, N. F.—Fluid-extract of rose and diluted sulphuric acid in syrup.

Rubi Fructus (*Rubi Fructus*), Blackberries, N. F.

ACTION AND USES: Flavor.

Cordiale Rubi Fructus (*Cord. Rubi Fruct.*), *Blackberry Cordial*, N. F.—Blackberry syrup (about 75%) and diluted alcohol (about 25%), with cinnamon, cloves and nutmeg. Absolute alcohol content about 12 per cent.

USES: Used as carminative in summer diarrhea. Not very efficient.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Syrupus Rubi Fructi (*Syr. Rubi Fruct.*), *Syrup of Blackberry Fruit*, N. F.—Juice of ripe blackberries in sugar.

USES: Flavoring vehicle.

Rubi Idaei Fructus (*Rubi Idaei Fructus*), Raspberries, N. F.

ACTION AND USES: Flavoring.

Syrupus Rubi Idaei (*Syr. Rubi Id.*), *Syrup of Raspberry*, N. F.—Juice of ripe raspberries in sugar.

Rubus (*Rubus*), Rubus, N. F. (Blackberry Bark).—Bark of the rhizome.

ACTION AND USES: Tannin-bearing astringent without advantage over krameria, etc.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Elixir Rubi Compositum (*Elix. Rubi Co.*), *Compound Elixir of Blackberry*, N. F.—Rubus (1.6%), nutgall (1.6%),

Saigon cinnamon, clove, mace, ginger, syrup of blackberry and diluted alcohol. Absolute alcohol content about 10 per cent.

USES: Used in "summer complaint."

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Fluidextractum Rubi (*Fldext. Rubi*), *Fluidextract of Rubus*, N. F.—*Rubus* (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Syrupus Rubi (*Syr. Rubi*), *Syrup of Rubus*, N. F.—*Fluidextract of rubus* (25%) in syrup.

USES: Intestinal astringent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Rumex (*Rumex*), *Rumex*, N. F. (Yellow Dock, Curled Dock).—Roots.

ACTION AND USES: Obsolete, unreliable and generally worthless "alterative and tonic"; mildly laxative.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Rumicis (*Fldext. Rumic.*), *Fluidextract of Rumex*, N. F.—*Rumex* (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Sabal (*Sabal*), *Sabal*, U. S. P. (Saw Palmetto Berries).—The partially dried ripe fruit.

ACTION AND USES: Widely exploited as a stimulant of the mucous membrane of the genito-urinary tract. Probably inferior to santal oil.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Fluidextractum Sabal (*Fldext. Sabal*), *Fluidextract of Sabal*, U. S. P. (*Fluidextract of Saw Palmetto*).—*Sabal* (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Tinctura Sabal et Santali (*Tr. Sabal et Santal.*), *Tincture of Saw Palmetto and Santal*, N. F.—*Sabal* (20%) and sandalwood (6.5%) in alcohol and water. Absolute alcohol content about 75 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Saccharum* (*Sacch.*), *Sugar*, U. S. P. (Sucrose).—Obtained from the sugar-cane, the sugar-beet, etc.

White, odorless crystals or powder with a sweet taste.

ACTION AND USES: Sweetening agent in official syrups and other preparations.

**Syrupus (Syrupus), Syrup, U. S. P.* (Sirup, Simple Syrup).—Sugar (85%) in water.

**Saccharum Lactis (Sacch. Lact.), Sugar of Milk, U. S. P.* (Milk Sugar, Lactose).—Obtained from the whey of cow's milk.

White, hard, odorless crystalline masses or powder, with a faintly sweet taste.

ACTION AND USES: Less sweet than cane sugar and less liable to ferment; slightly laxative and diuretic; frequently used to modify cow's milk in the feeding of infants.

SALES EFFERVESCENTES, GRANULAR EFFERVESCENT SALTS.—Effervescent salts are mixtures of active constituents with sodium bicarbonate, citric acid and tartaric acid. They are usually directed to be produced and dispensed in coarse granules. Their advantage over other methods of administering active ingredients is rather doubtful. For effervescent salts included in the Pharmacopeia and in the National Formulary, see under the names of the active ingredients. For *Sal Carolinum Factitium* and *Sal Carolinum Factitium Effervescens*, see under *Sodii Sulphas*; for *Sal Kissingense Factitium* and *Sal Kissingense Factitium Effervescens*, see under *Sodii Chloridum*; for *Sal Vichyanum Factitium*, *Sal Vichyanum Factitium Effervescens* and *Sal Vichyanum Factitium Effervescens cum Lithio*, see under *Sodii Bicarbonas*.

Salicinum (Salicin), Salicin, U. S. P.—A glucoside obtained from certain species of willows and poplars.

Colorless crystals or white powder, odorless and having a very bitter taste.

ACTION AND USES: Partially decomposed in the stomach and intestines and finally oxidized in the body into salicylic acid. Less irritant to the mucous membranes than the salicylates but also less certain in its action, and has been practically superseded by the latter.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Sambucus (Sambuc.), Sambucus, N. F. (Elder Flowers).

ACTION AND USES: Obsolete and worthless remedy in dropsy. Large doses are said to act like a hydragogue cathartic.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Sanguinaria (Sanguin.), Sanguinaria, U. S. P. (Blood Root).
—Rhizome and roots.

ACTION AND USES: Irritant and narcotic; unreliable; expectorant in small doses and nauseant in large doses, which may produce serious secondary symptoms.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Fluidextractum Sanguinariae (Fldext. Sanguin.), Fluid-extract of Sanguinaria, N. F.—Sanguinaria (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 0.1 Cc. or 1½ minims (N. F.).

Syrupus Sanguinariae (Syr. Sanguinar.), Syrup of Sanguinaria, N. F. (Syrup of Bloodroot).—Sanguinaria (22.5%) and acetic acid in syrup.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Sanguinariae (Tr. Sanguin.), Tincture of Sanguinaria, U. S. P. (Tincture of Bloodroot).—Sanguinaria (10%) and hydrochloric acid (1%) in alcohol and water. Absolute alcohol content about 56 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Santalum Album (Santal. Alb.), Sandal Wood, N. F. (White Sandal Wood).—Heartwood.

ACTION AND USES: Similar to those of *Oleum Santali*, which see.

Santalum Rubrum (Santal. Rub.), Red Saunders, U. S. P.—Heartwood.

ACTION AND USES: Used as a coloring agent.

***Santoninum (Santonin.), Santonin, U. S. P.**—Anhydride or lactone of santoninic acid, obtained from Levant wormseed.

Colorless crystals or powder, becoming yellow on exposure to light, odorless and at first nearly tasteless, afterward becoming bitter.

ACTION AND USES: Used for its poisonous action on intestinal parasites, especially ascaris. When absorbed in sufficient amounts it produces yellow vision and epileptiform convulsions.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.) in powder or capsules.

Trochisci Santonini (Troch. Santonin.), Troches of Santonin, N. F.—Each troche contains santonin (0.03 Gm. or ½ grain), with tragacanth, sugar, cocoa and vanillin.

USES: Anthelmintic. Objectionable since their resemblance to candy may lead to serious poisoning.

DOSAGE: 1 troche (N. F.).

Trochisci Santonini Compositi (*Troch. Santonin. Co.*), *Compound Troches of Santonin*, N. F. (Troches of Santonin and Calomel).—Each troche contains santonin and mild mercurous chloride (each 0.03 Gm. or $\frac{1}{2}$ grain) with tragacanth, sugar, cocoa and vanillin.

USES: Anthelmintic; objectionable like *Trochisci Santonini*, which see above.

DOSAGE: 1 troche (N. F.).

***Sapo (Sapo), Soap, U. S. P.** (White Castile Soap).—Soap prepared from olive oil and sodium hydroxide.

A white or whitish solid or powder, having a faint, peculiar odor and an alkaline taste.

ACTION AND USES: Used chiefly as a detergent and in solution as a vehicle for liniments. Also an ingredient of pills containing resinous drugs like aloes and scammony.

Emplastrum Saponis (*Emp. Sapon.*), *Soap Plaster*, N. F.—Made from soap (10%) and lead plaster (90%).

Linimentum Saponato-Camphoratum (*Lin. Sapon.-Camph.*), *Camphorated Soap Liniment*, N. F. (Opodeldoc, Solid Opodeldoc).—Camphor (2.5%), ammonia water, sodium stearate, oil of thyme, oil of rosemary, alcohol and water. Absolute alcohol content about 75 per cent.

***Linimentum Saponis** (*Lin. Sapon.*), *Soap Liniment, U. S. P.* (Liquid Opodeldoc).—Soap (6%), camphor (4.5%) in oil of rosemary, alcohol and water. Absolute alcohol content about 65 per cent.

USES: Mild rubefacient and vehicle for more active liniments.

***Sapo Mollis (Sapo Moll.)**, *Soft Soap, U. S. P.*—Prepared from cottonseed oil and potassium hydroxide.

A soft, unctuous, yellowish-white to brownish-yellow mass, having a slight, characteristic odor and an alkaline taste.

ACTION AND USES: Dissolved in diluted alcohol, is employed in the preparation of various liniments, and as a detergent.

Linimentum Saponis Mollis (*Lin. Sapon. Moll.*), *Liniment of Soft Soap, U. S. P.* (Tincture of Green Soap).—Soft soap (65%) and oil of lavender in alcohol. Absolute alcohol content about 30 per cent.

Linimentum Saponis Mollis Compositum (*Lin. Sapon. Mol. Co.*), *Compound Liniment of Soft Soap, N. F.* (*Tinctura Saponis Viridis Composita*, N. F. III).—Soft soap (15%), oil of cade (2%) and alcohol. Absolute alcohol content about 75 per cent.

Sarsaparilla (Sarsap.), Sarsaparilla, U. S. P. (Sarsaparilla Root).

ACTION AND USES: Formerly used in chronic rheumatism, skin diseases and syphilis; inefficient and therefore harmful; fortunately passing out of use.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Decoctum Sarsaparillae Compositum (Decoc. Sarsap. Co.), Compound Decoction of Sarsaparilla, N. F.—Sarsaparilla (10%), guaiac wood (2%), mezereum (1%), sassafras, glycyrrhiza and water.

USES: Antiquated and useless "alterative."

DOSAGE: 120 Cc. or 4 fluidounces (N. F.).

Fluidextractum Sarsaparillae (Fldext. Sarsap.), Fluidextract of Sarsaparilla, U. S. P.—Sarsaparilla (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Fluidextractum Sarsaparillae Compositum (Fldext. Sarsap. Co.), Compound Fluidextract of Sarsaparilla, U. S. P.—Sarsaparilla (75%), mezereum (3%), glycyrrhiza and sassafras. Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Syrupus Sarsaparillae Compositus (Syr. Sarsap. Co.), Compound Syrup of Sarsaparilla, U. S. P.—Fluidextract of sarsaparilla (20%), fluidextract of senna (1.5%), fluidextract of glycyrrhiza, oil of sassafras, oil of anise, methyl salicylate and alcohol in syrup. Absolute alcohol content about 10 per cent.

USES: Irrational mixture; no indications for its use.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

Sassafras (Sassaf.), Sassafras, U. S. P.—Bark of the root.

ACTION AND USES: Mild aromatic and carminative containing a volatile oil. No advantage over other aromatics, such as peppermint, ginger, etc.

DOSAGE: 10 Gm. or 2½ drachms (U. S. P.).

Sassafras Medulla (Sassaf. Med.), Sassafras Pith, N. F.

ACTION AND USES: Demulcent.

Mucilago Sassafras Medullae (Mucil. Sassaf. Medul.), Mucilage of Sassafras Pith, N. F.—Sassafras (0.2%) in water.

DOSAGE: 15 Cc. or 4 fluidrachms (N. F.).

Scammoniae Radix (Scam. Rad.), Scammony Root, U. S. P.

ACTION AND USES: Used principally for preparing resin, in which its activity as a cathartic resides.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Resina Scammoniae (Res. Scamm.), Resin of Scammony, U. S. P.—The resin from scammony root.

DOSAGE: 0.2 Gm. or 3 grains (U. S. P.).

**Scilla* (Scill.), Squill, U. S. P.—Fleshy inner scales of a bulb. The official drug is to be assayed biologically.

ACTION AND USES: Qualitatively like those of digitalis. Has been largely used as a nauseant and expectorant. Has no advantage over digitalis, and should pass out of use. If used, it should be standardized.

DOSAGE: 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.) of a standardized preparation.

Acetum Scillae (Acet. Scill.), Vinegar of Squill, U. S. P.—Squill (10%) in diluted acetic acid.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Fluidextractum Scillae (Fldext. Scill.), Fluidextract of Squill, U. S. P.—Squill (100%). Assayed biologically by producing death in the frog.

DOSAGE: 0.1 Cc. or $1\frac{1}{2}$ minims (U. S. P.). Physicians who prescribe it should specify the biologically assayed preparation.

Oxymel Scillae (Oxymel Scill.), Oxymel of Squill, N. F.—Vinegar of squill (50%) and clarified honey.

USES: Antiquated and irrational preparation of squill

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

**Syrupus Scillae* (Syr. Scill.), Syrup of Squill, U. S. P.—Vinegar of squill (45%), in sugar and water.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Syrupus Scillae Compositus (Syr. Scill. Co.), Compound Syrup of Squill, U. S. P. (Hive Syrup).—Fluidextract of squill (8%), fluidextract of senega (8%) and antimony and potassium tartrate (0.2%), in water and syrup.

USES: Complex and irrational expectorant.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

**Tinctura Scillae* (Tr. Scill.), Tincture of Squill, U. S. P.—Squill (10%) in alcohol and water. Biologically assayed by producing death in frogs. Absolute alcohol content about 69 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.). Physicians who prescribe it should specify the biologically assayed preparation.

Scoparius (Scopar.), Scoparius, N. F. (Broom Tops).

ACTION AND USES: Formerly used as diuretic, laxative and cathartic. See *Sparteinae Sulphas*.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Scoparii (Fldext. Scopar.), Fluidextract of Scoparius, N. F.—Scoparius (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

***Scopolaminae Hydrobromidum (Scopolamin. Hydrobrom.), Scopolamine Hydrobromide, U. S. P. (Hyosine Hydrobromide, Scopolamine Bromide).**—The hydrobromide of levorotatory scopolamine, also known as hyosine.

Colorless, odorless, transparent crystals.

ACTION AND USES: Closely resembling those of atropine in its influence on the nerve endings but differing from the latter in having a sedative instead of a stimulating effect on the brain. Used as a somnifacient in motor excitement and mania; as a preliminary to anesthesia and in "twilight sleep," and locally as a mydriatic. Uncertain in its action, at times producing acute delirium. The tendency to depress the respiratory center must be remembered.

DOSAGE: 0.3 Mg. or $\frac{1}{200}$ grain (U. S. P.).

Scutellaria (Scutell.), Scutellaria, N. F. (Skullcap).—Plant.

ACTION AND USES: Claimed, though without reliable evidence, to be a tonic, nervine and antispasmodic, particularly useful in chorea and convulsions.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Scutellariae (Fldext. Scutellar.), Fluidextract of Scutellaria, N. F.—Scutellaria (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Senecio (Senecio), Senecio, N. F. (Life Root).—Plant deprived of its root.

ACTION AND USES: Has been recommended, probably without warrant, as a tonic, diuretic, expectorant and diaphoretic.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Senecionis (*Fldext. Senecion.*), *Fluidextract of Senecio*, N. F.—Senecio (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Senega (*Seneg.*), **Senega**, U. S. P. (*Seneca Snakeroot*, *Senega Snakeroot*).

ACTION AND USES: Employed as nauseant expectorant (by virtue of the irritant saponin).

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Fluidextractum Senegae (*Fldext. Seneg.*), *Fluidextract of Senega*, U. S. P.—Senega (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Syrupus Senegae (*Syr. Seneg.*), *Syrup of Senega*, U. S. P.—Fluidextract of senega (20%) in syrup. Absolute alcohol content about 10 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

***Senna** (*Senn.*), **Senna**, U. S. P. (*Alexandria Senna* or *India Senna*).

ACTION AND USES: Efficient cathartic of anthraquinone series.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Confectio Sennae (*Confect. Senn.*), *Confection of Senna*, N. F.—Senna (10%), cassia fistula (16%), tamarind (10%), prune, fig, sugar, oil of coriander and water.

USES: A pleasant but needlessly complex laxative.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

***Fluidextractum Sennae** (*Fldext. Senn.*), *Fluidextract of Senna*, U. S. P.—Senna (100%). Absolute alcohol content about 25 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Infusum Sennae Compositum (*Inf. Senn. Co.*), *Compound Infusion of Senna*, U. S. P.—Senna (6%), magnesium sulphate (12%), manna (12%) and fennel in water.

DOSAGE: 120 Cc. or 4 fluidounces (U. S. P.).

***Pulvis Glycyrrhizae Compositus** (*Pulv. Glycyrrh. Co.*), *Compound Powder of Glycyrrhiza*, U. S. P. (*Compound Licorice Powder*).—Senna (18%), washed sulphur (8%) with glycyrrhiza, oil of fennel and sugar.

USES: A misnamed senna preparation.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Species Laxativae (Spec. Lax.), Laxative Species, N. F. (Species Laxantes, N. F. III, St. Germain Tea).—Senna (40%), potassium bitartrate (10%), elder flowers, fennel and anise.

DOSAGE: 1.3 Gm. or 20 grains (N. F.).

Syrupus Ficorum Compositus (Syr. Ficor. Co.), Compound Syrup of Figs, N. F.—Fluidextract of senna (20%), aromatic fluidglycerate of cascara sagrada (10%), with figs, oil of fennel and spirit of peppermint in syrup. Absolute alcohol content about 5 per cent.

USES: Complex and misnamed cathartic, imitating a widely advertised "patent medicine."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

**Syrupus Sennae (Syr. Senn.), Syrup of Senna, U. S. P.*—Fluidextract of senna (25%) and oil of coriander in syrup. Absolute alcohol content about 5 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

Syrupus Sennae Aromaticus (Syr. Senn. Arom.), Aromatic Syrup of Senna, N. F.—Fluidextract of senna (12.5%), jalap (5%), rhubarb (1.75%), Saigon cinnamon, clove, nutmeg, oil of lemon and sugar in diluted alcohol. Absolute alcohol content about 30 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Syrupus Sennae Compositus (Syr. Senn. Co.), Compound Syrup of Senna, N. F.—Fluidextract of senna (13.5%), fluidextract of rhubarb (3.5%), fluidextract of frangula (3.5%), methyl salicylate (0.4%) and alcohol in syrup. Absolute alcohol content about 13 per cent.

USES: Complex and irrational cathartic.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Serpentaria (Serpent.), Serpentina, U. S. P. (Texas Snake-root, Virginia Snakeroot).—Rhizome and root.

ACTION AND USES: Bitter with no advantage over gentian; now less frequently used than formerly.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Fluidextractum Serpentinae (Fldext. Serpentar.), Fluidextract of Serpentina, N. F.—Serpentina (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 1 Cc. or 15 minims.

Tinctura Serpentinae (Tr. Serpent.), Tincture of Serpentina, N. F.—Serpentina (20%) in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

- ***Serum Antidiphthericum** (Ser. Antidiph.), **Antidiphtheric Serum**, U. S. P. (Diphtheria Antitoxin).—Serum from a horse properly immunized against diphtheria toxin; has a potency of not less than 250 antitoxic units per Cc.

ACTION AND USES: Curative and prophylactic agent in diphtheria. The possibility of serum disease must be remembered. Caution should be exercised when antidiphtheric serum is administered to persons who have had asthma, and, above all, "horse asthma." In such cases a few drops only of the serum may be injected, followed in fifteen or thirty minutes by the full dose.

DOSAGE: Curative, 10,000 units (U. S. P.); protective, 1,000 units (U. S. P.); both hypodermic.

- ***Serum Antidiphthericum Purificatum** (Ser. Antidiph. Purif.), **Purified Antidiphtheric Serum**, U. S. P. (Antidiphtheric Globulins, Concentrated Diphtheria Antitoxin, Diphtheric Antitoxin Globulins, Refined and Concentrated Diphtheria Antitoxin).—Certain antitoxic proteins from the blood serum of a horse properly immunized against diphtheria toxin, dissolved in physiologic solution of sodium chloride.

USES: Less apt to produce serum disease.

DOSAGE: Curative, 10,000 units (U. S. P.); protective, 1,000 units (U. S. P.); both hypodermic.

- ***Serum Antidiphthericum Siccum** (Ser. Antidiph. Sicc.), **Dried Antidiphtheric Serum**, U. S. P. (Dried Diphtheria Antitoxin).

DOSAGE: Curative, 10,000 units (U. S. P.); protective, 1,000 units (U. S. P.); both hypodermic.

- ***Serum Antitetanicum** (Ser. Antitetan.), **Antitetanic Serum**, U. S. P. (Tetanus Antitoxin).—Serum from a horse properly immunized against tetanus toxin, dissolved in physiologic solution of sodium chloride.

ACTION AND USES: Prophylactic agent in tetanus; also used for curative purposes.

DOSAGE: Curative, 10,000 units (U. S. P.); protective, 1,500 units (U. S. P.); both hypodermic.

- ***Serum Antitetanicum Purificatum** (Ser. Antitetan. Purif.), **Purified Antitetanic Serum**, U. S. P. (Antitetanic Globulins, Concentrated Tetanus Antitoxin, Refined and Concentrated Tetanus Antitoxin, Tetanus Antitoxic Globulins).—Certain antitoxic proteins from the serum of a horse properly immunized against tetanus toxin.

DOSAGE: Curative, 10,000 units (U. S. P.); protective, 1,500 units (U. S. P.); both hypodermic.

**Serum Antitetanicum Siccum* (Ser. Antiletan. Sicc.), Dried Antitetanic Serum, U. S. P. (Dried Tetanus Antitoxin).

USES: May be used as a dusting powder in suspicious wounds.

DOSAGE: Curative, 10,000 units (U. S. P.); protective, 1,500 units (U. S. P.); both hypodermic.

Sevum Praeparatum (Sev. Praep.), Prepared Suet, U. S. P. (Mutton Suet).

A white, solid, almost odorless fat, having a bland taste when fresh, but becoming rancid on prolonged exposure to air.

ACTION AND USES: Used in the preparation of ointments and cerates.

Sevum Benzoinatum (Sev. Benz.), Benzoinated Suet, N. F. —Benzoin (3%) in prepared suet.

USES: Ointment base; superfluous.

Sinapis Alba (Sinap. Alb.), White Mustard, U. S. P. (Yellow Mustard).—Ripe seeds.

ACTION AND USES: Develops irritant volatile oil on contact with water; frequently mixed with black mustard in powder for use as ground mustard.

DOSAGE: Emetic, 10 Gm. or $2\frac{1}{2}$ drachms (U. S. P.).

**Sinapis Nigra* (Sinap. Nig.), Black Mustard, U. S. P. (Brown Mustard).—Ripe seeds.

ACTION AND USES: See *Oleum Sinapis Volatile*.

DOSAGE: Emetic, 10 Gm. or $2\frac{1}{2}$ drachms (U. S. P.).

**Emplastrum Sinapis* (Emp. Sinap.), Mustard Plaster, U. S. P. (Charta Sinapis, U. S. P. VIII, Mustard Paper).

—A mixture of black mustard, deprived of its fixed oil, and a solution of rubber, spread on paper, cotton cloth or other fabric.

USES: Counterirritant.

Soda Chlorinata, Chlorinated Soda.

ACTION AND USES: Hypochlorites in acid, alkaline and neutral solutions have been found useful in the treatment of infected wounds. The alkaline solution is represented by the *Liquor Sodae Chlorinatae* below, and by *Potassae Chlorinatae*, which see under *P*.

**Liquor Sodae Chlorinatae* (Liq. Sod. C of Chlorinated Soda, U. S. P. (Laba

Chlorine compounds of sodium containing not less than 2.5 per cent. available Cl.

Soda cum Calce (Sod. c. Calc.), Soda with Lime, N. F. (London Paste).—Sodium hydroxide and calcium oxide (each 50%).

ACTION AND USES: Caustic.

Sodii Acetas (Sod. Acet.), Sodium Acetate, U. S. P.— $\text{NaC}_2\text{H}_3\text{O}_2 + 3\text{H}_2\text{O}$.

Colorless crystals or granular powder, odorless or nearly odorless, and having a cooling, saline taste. Very soluble in water (1:0.8) and soluble in alcohol (1:19).

ACTION AND USES: Practically identical with those of *Potassii Acetas*, which see.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Sodii Arsenas (Sod. Arsen.), Sodium Arsenate, U. S. P. (Arsenas Sodii, P. I.).— $\text{Na}_2\text{HAsO}_4 + 7\text{H}_2\text{O}$.

Colorless, odorless crystals, efflorescent in dry air, and somewhat deliquescent in moist air. Freely soluble in water (1:1.5) and slightly soluble in alcohol.

ACTION AND USES: Has the same general action and uses as arsenic trioxide, but is less active.

DOSAGE: 0.005 Gm. or $\frac{1}{12}$ grain (U. S. P.). Usually given in the form of solution or pill.

Sodii Arsenas Exsiccatus (Sod. Arsen. Exsic.), Exsiccated Sodium Arsenate, U. S. P.— Na_2HAsO_4 .

White, odorless, amorphous powder. Freely soluble in water (1:3.1), but only slightly soluble in alcohol.

DOSAGE: 0.003 Gm. or $\frac{1}{20}$ grain (U. S. P.).

Liquor Sodii Arsenatis (Liq. Sod. Arsen.), Solution of Sodium Arsenate, U. S. P.—Exsiccated sodium arsenate (about 1%) in water.

USES: No advantage over other arsenicals.

DOSAGE: 0.2 Cc. or 3 minims (U. S. P.).

Liquor Sodii Arsenatis, Pearson (Liq. Sod. Arsen. Pearson), *Pearson's Solution of Sodium Arsenate*, N. F. (Liquor Arsenicalis Pearsonii).—Exsiccated sodium arsenate (0.1%) in water. Should not be confounded with *Liquor Sodii Arsenatis*, U. S. P., which is ten times stronger.

USES: Superfluous arsenic preparation.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Sodii Benzoas** (Sod. Benz.), Sodium Benzoate, U. S. P.— $\text{Na}(\text{C}_6\text{H}_5.\text{COO})$.

White, odorless, sweetish powder. Freely soluble in water (1:1.8), and soluble in alcohol (1:61). Incompatible with mineral acids and ferric salts.

ACTION AND USES: Has the action of benzoic acid, but is less irritating. Mild antiseptic, practically non-toxic.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Sodii Benzosulphinidum (Sod. Benzosulphin.), Sodium Benzosulphinide, U. S. P. (Sodium - Saccharin, Soluble Saccharin).

Colorless crystals or white powder, odorless or nearly odorless, and having an intensely sweet taste. Freely soluble in water (1:1.2) and soluble in alcohol (1:50).

ACTION AND USES: Those of benzosulphinide (saccharin), but with the advantage of ready solubility.

DOSAGE: 0.2 Gm. or 3 grains (U. S. P.).

***Sodii Bicarbonas (Sod. Bicarb.), Sodium Bicarbonate, U. S. P.** (Baking Soda).— NaHCO_3 .

A white, odorless powder, having a cooling, mildly alkaline taste. Decomposed by acids and converted by boiling into the normal carbonate. Soluble in water (1:10) and insoluble in alcohol.

ACTION AND USES: Used as a non-caustic alkali in conditions of acidosis, and to neutralize the acid of the gastric juice in hyperacidity and gastric ulcer. May be used intravenously in cases of extreme acidosis. Externally, used as a mild alkaline wash.

DOSAGE: 1 Gm. or 15 grains (U. S. P.). For intravenous injection a 6 per cent. solution sterilized by boiling and thus partly converted into the normal carbonate has been recommended. One thousand Cc. of such a solution may be injected, but great care must be taken that none of the liquid gets outside the veins lest necrosis of the tissues occur.

Liquor Sodae et Menthae (Liq. Sod. et Menth.), Solution of Soda and Mint, N. F. (Mistura Sodae et Menthae, N. F. III, Soda Mint).—Sodium bicarbonate (5%), aromatic spirit of ammonia (1%) and spearmint water.

USES: Antacid and carminative.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Sal Vichyanum Factitium (Sal Vichy. Fact.), Artificial Vichy Salt, N. F.—Potassium carbonate (3.85%), magnesium sulphate (8%), sodium chloride (7.7%) and sodium bicarbonate (84.6%).

USES: Alkaline saline cathartic.

Sal Vichyanum Factitium Effervescens (*Sal Vichy. Fact. Eff.*), *Effervescent Artificial Vichy Salt*, N. F. (*Pulvis Salis Vichyani Factitii Effervescens*, N. F. III).—Artificial vichy salt (25%) in a mixture of sodium bicarbonate (48.5%), tartaric acid (16.45%) and citric acid (yielding sodium citrate and tartrate).

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Sal Vichyanum Factitium Effervescens cum Lithio (*Sal Vichy. Fact. Eff. c. Lith.*), *Effervescent Artificial Vichy Salt with Lithium*, N. F. (*Pulvis Salis Vichyani Factitii Effervescens cum Lithio*, N. F. III).—Artificial vichy salt (25%), lithium citrate (8.33%) in a mixture of sodium bicarbonate (44.14%), tartaric acid (12.54%) and citric acid (yielding sodium tartrate and citrate).

USES: Irrational laxative mixture. The lithium serves no useful purpose.

DOSAGE: 6 Gm. or $1\frac{1}{2}$ drachms (N. F.).

Trochisci Sodii Bicarbonatis (*Troch. Sod. Bicarb.*), *Troches of Sodium Bicarbonate*, U. S. P.—Each troche contains sodium bicarbonate (0.18 Gm.) with sugar, nutmeg and mucilage of tragacanth.

**Sodii Boras* (*Sod. Bor.*), *Sodium Borate*, U. S. P. (*Borax*, *Sodium Tetraborate*, *Sodium Pyroborate*).— $\text{Na}_2\text{B}_4\text{O}_7 + 10\text{H}_2\text{O}$.

Colorless crystals or white powder, odorless and having a sweetish, alkaline taste. Soluble in water (1:15) and insoluble in alcohol.

ACTION AND USES: Antiseptic detergent and alkaline. Used in the form of solution as a wash both externally and in the body cavities. Seldom used internally.

DOSAGE: 0.75 Gm. or 12 grains (U. S. P.).

Liquor Sodii Boratis Compositus (*Liq. Sod. Bor. Co.*), *Compound Solution of Sodium Borate*, N. F. (*Dobell's Solution*).—Sodium borate and sodium bicarbonate (each 1.5%) and liquefied phenol (0.3%) in glycerin and water.

USES: Mild antiseptic.

Mel Rosae et Sodii Boratis (*Mel Ros. et Sod. Bor.*), *Honey of Rose and Sodium Borate*, N. F. (*Honey of Rose with Borax*).—Sodium borate (10%), glycerin and honey of rose.

USES: Alkaline demulcent.

Mel Sodii Boratis (*Mel Sod. Bor.*), *Honey of Sodium Borate*, N. F. (*Mel Boracis*, *Honey and Borax*).—Sodium borate (10%) in glycerin and clarified honey.

USES: Alkaline demulcent.

Sodii Boro-Benzoas (Sod. Boro-Benz.), Sodium Boro-Benzoate, N. F.—Sodium borate (43%) and sodium benzoate (57%).

ACTION AND USES: Feeble alkaline antiseptic.

DOSAGE: 2 Gm. or 30 grains (N. F.).

***Sodii Bromidum** (Sod. Brom.), Sodium Bromide, U. S. P.—NaBr.

White, odorless crystals or powder, having a saline taste. Freely soluble in water (1:1.1) and soluble in alcohol (1:16).

ACTION AND USES: Used as a nerve sedative and cerebral depressant. Practically identical with potassium bromide in action and uses, but somewhat less irritating.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Sodii Bromidi (Elix. Sod. Brom.), *Elixir of Sodium Bromide*, N. F.—Sodium bromide (1.75%) in syrup, water and aromatic elixir. Absolute alcohol content about 7.5 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

***Sodii Cacodylas** (Sod. Cacodyl.), Sodium Cacodylate, U. S. P.—Sodium dimethylarsenate.

White, odorless, deliquescent crystals or powder. Very soluble in water (1:0.5) and freely soluble in alcohol (1:2.5).

ACTION AND USES: Is slowly decomposed in the tissues into trivalent arsenic. The action is thus more gradual and less toxic than that of other arsenic compounds. Produces garlic breath. Used especially by hypodermic injection, in anemias, etc.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

***Sodii Carbonas Monohydratus** (Sod. Carb. Monohyd.), Monohydrated Sodium Carbonate, U. S. P.— $\text{Na}_2\text{CO}_3 + \text{H}_2\text{O}$.

White, odorless, granular powder having a strongly alkaline taste. Freely soluble in water (1:3) and insoluble in alcohol. Incompatible with acids and acid salts and with the salts of the heavy metals and alkaloids.

ACTION AND USES: Antacid and detergent; employed in medicine chiefly in the preparation of alkaline baths.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

***Sodii Chloridum** (Sod. Chlorid.), Sodium Chloride, U. S. P.—NaCl.

Colorless crystals or white powder, odorless, and having taste. Freely soluble in water (1:2.8), and only slightly in alcohol.

ACTION AND USES: Used for preparing physiologic salt solution. Given by mouth in large doses, is emetic and in proper dilutions, laxative. Excessive use as a seasoning for foods is particularly harmful in nephritis.

DOSAGE: Emetic, 15 Gm. or 4 drachms (U. S. P.).

Liquor Sodii Chloridi Physiologicus (Liq. Sod. Chlor. Physio.), *Physiological Solution of Sodium Chloride*, U. S. P. (Physiological Salt Solution, Normal Salt Solution).—Sodium chloride (0.85%) in water.

USES: An indifferent vehicle, used especially for intravenous injections.

Sal Kissingense Factitium (Sal Kissingen. Fact.), *Artificial Kissingen Salt*, N. F.—Potassium chloride (2.8%), sodium chloride (59.4%), magnesium sulphate (20%), sodium bicarbonate (17.8%).

USES: Alkaline saline cathartic.

Sal Kissingense Factitium Effervescens (Sal Kissingen. Fact. Eff.), *Effervescent Artificial Kissingen Salt*, N. F. (Pulvis Salis Kissingensis Factitii Effervescens, N. F. III).—Artificial Kissingen salt (40%), sodium bicarbonate (40.6%), tartaric acid (9.4%) and citric acid (yielding sodium tartrate and citrate).

DOSAGE: 5.5 Gm. or 80 grains (N. F.).

Sodii Citras (Sod. Cit.), *Sodium Citrate*, U. S. P.—The hydrated salt.

White, odorless granular powder or small crystals, having a cooling saline taste. Freely soluble in water (1:1.3); insoluble in alcohol.

ACTION AND USES: Similar to those of potassium citrate.

DOSAGE: 1 Gm. or 15 grains.

Liquor Sodii Citratis (Liq. Sod. Cit.), *Solution of Sodium Citrate*, N. F. (Mistura Sodii Citratis, Potio Rivierii).—Sodium citrate (3.5%) in water.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Liquor Sodii Citro-Tartratis Effervescens (Liq. Sod. Citro-Tart. Eff.), *Effervescent Solution of Sodium Citro-Tartrate*, N. F.—Sodium tartrate (about 8%) with syrup of citric acid (about 12%) and water.

DOSAGE: The contents of one bottle, about 350 Cc. or nearly 12 fluidounces (N. F.).

Sodii Cyanidum (Sod. Cyanid.), *Sodium Cyanide*, U. S. P.—NaCN.

White masses or powder, odorless when perfectly dry; deliquescent in the air and exhaling the odor of hydrocyanic acid. Freely soluble in cold water.

ACTION AND USES: Sodium cyanide replaces potassium cyanide, U. S. P. VIII. The alkaline cyanides are rarely administered internally, but their use in solution has been suggested as a substitute for diluted hydrocyanic acid.

Sodii Glycerophosphas (Sod. Glycerophos.), Sodium Glycerophosphate, U. S. P. (Sodium Glycerinophosphate).—Hydrated sodium glycerophosphate. Represents not less than 68 per cent. of the anhydrous salt.

White, odorless plates, scales or powder, having a saline taste. Very soluble in hot or cold water and nearly insoluble in alcohol.

ACTION AND USES: Has been recommended in solutions as a nerve tonic in all kinds of wasting and nervous diseases. The weight of evidence indicates that the glycerophosphates are not superior to ordinary phosphates and that the claims for them as nerve tonics are, therefore, unfounded.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

*Elixir Glycerophosphatum Compositum (Elix. Glycerophos. Co.), Compound Elixir of Glycerophosphates, N. F. (Compound Solution of Glycerophosphates).—*Solution of sodium glycerophosphate (4%), calcium glycerophosphate (1.6%), ferric glycerophosphate (0.3%), soluble manganese glycerophosphate (0.2%), quinine glycerophosphate (0.1%), strychnine glycerophosphate (0.015%), lactic acid (1%) and compound spirit of cardamom, in alcohol, glycerin and water. Absolute alcohol content about 12 per cent.

USES: A highly complex and therefore irrational preparation of the inactive glycerophosphates.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

*Liquor Sodii Glycerophosphatis (Liq. Sod. Glycerophos.). Solution of Sodium Glycerophosphate, U. S. P.—*Anhydrous sodium glycerophosphate (not less than 50%) in water.

DOSAGE: 0.35 Cc. or 6 minims (U. S. P.).

***Sodii Hydroxidum (Sod. Hydrox.), Sodium Hydroxide, U. S. P. (Caustic Soda, Sodium Hydrate).—**NaOH (not less than 90%).

White, odorless masses or hard, brittle sticks. Very soluble in water (1:0.9) and very soluble in alcohol.

ACTION AND USES: Practically identical with those of potassium hydroxide.

***Liquor Sodii Hydroxidi** (*Liq. Sod. Hydrox.*), *Solution of Sodium Hydroxide*, U. S. P. (Liquor Sodae, Solution of Soda).—NaOH (about 5%) in water.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Sodii Hypophosphis (*Sod. Hypophos.*), *Sodium Hypophosphite*, U. S. P.— $\text{NaH}_2\text{PO}_2 + \text{H}_2\text{O}$.

Colorless plates or white powder, odorless and having a saline taste. Freely soluble in water (1:1) and soluble in alcohol.

ACTION AND USES: The hypophosphites have been prescribed as "tonics," but all reliable evidence indicates that they are inert.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Sodii Hypophosphitis (*Elix. Sod. Hypophos.*), *Elixir of Sodium Hypophosphite*, N. F.—Sodium hypophosphite (3.5%), hypophosphorous acid (0.4%), and aromatic elixir. Absolute alcohol content about 21 per cent.

USES: Objectionable alcoholic preparation of an inactive drug.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Syrupus Sodii Hypophosphitis (*Syr. Sod. Hypophos.*), *Syrup of Sodium Hypophosphite*, N. F.—Sodium hypophosphite (3.5%), and hypophosphorous acid (0.2%) in syrup.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Sodii Indigotindisulphonas (*Sod. Indigotin.*), *Sodium Indigotindisulphonate*, U. S. P. (Indigo Carmine).

Blue powder or dark purple paste. Sparingly soluble in water and almost insoluble in alcohol.

ACTION AND USES: Coloring agent.

***Sodii Iodidum** (*Sod. Iod.*), *Sodium Iodide*, U. S. P.—NaI.

Colorless crystals or white powder, odorless and having a saline taste. Very soluble in water (1:0.55) and freely soluble in alcohol (1:2).

ACTION AND USES: Practically identical with those of potassium iodide, but less disagreeable in taste; perhaps somewhat less irritating locally.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.).

***Sodii Nitris** (*Sod. Nitris*), *Sodium Nitrite*, U. S. P.— NaNO_2 .

Colorless crystals or white or nearly white masses, sticks or powder, odorless and having a mild saline taste. Deliquescent on exposure to air, being gradually converted into sodium nitrate. Freely soluble in water (1:1.5), but only sparingly soluble in alcohol.

ACTION AND USES: Vasodilator, similar to nitroglycerin. The action is probably somewhat slower and more prolonged.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Sodii Perboras (Sod. Perbor.), Sodium Perborate, U. S. P.
— $\text{NaBO}_3 + 4\text{H}_2\text{O}$ (about 86.5%). Should contain not less than 9 per cent. of available oxygen.

White, odorless granules or powder, having a saline taste. Soluble in water.

ACTION AND USES: Antiseptic, deodorant and bactericide; in general, similar to hydrogen peroxide.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.); may be applied as a dusting powder or in a 2 per cent. solution.

Sodii Phenolsulphonas (Sod. Phenolsulph.), Sodium Phenolsulphonate, U. S. P. (Sodium Sulphocarbolate).—The hydrated salt.

Colorless, odorless, somewhat efflorescent crystals or granules, having a cooling, saline bitter taste. Freely soluble in water (1:4.2), and only slightly soluble in alcohol (1:140).

ACTION AND USES: Phenolsulphonates were introduced with the idea that they would be antiseptic like phenol, but less toxic. Actually, they are practically inert.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

***Sodii Phosphas (Sod. Phos.), Sodium Phosphate, U. S. P.**
— $\text{Na}_2\text{HPO}_4 + 12\text{H}_2\text{O}$, representing about 41.5 per cent. of the anhydrous salt.

Colorless, odorless, efflorescent crystals or powder, having a cooling, saline taste. Freely soluble in water (1:2.7); insoluble in alcohol.

ACTION AND USES: Saline cathartic, similar to magnesium and sodium sulphate. The taste is less disagreeable.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Sodii Phosphas Exsiccatus (Sod. Phos. Exsic.), Exsiccated Sodium Phosphate, U. S. P.— Na_2HPO_4 .

White powder which absorbs moisture readily. Freely soluble in water (1:8.1), but insoluble in alcohol.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Liquor Sodii Phosphatis Compositus (Liq. Sod. Phos. Co.), Compound Solution of Sodium Phosphate, N. F.—Sodium phosphate (100%) and citric acid in glycerin and water.

USES: Laxative.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

***Sodii Phosphas Effervescens** (*Sod. Phos. Eff.*), *Effervescent Sodium Phosphate*, U. S. P.—Exsiccated sodium phosphate (20%) in a mixture of sodium bicarbonate (47.7%), tartaric acid (25.2%) and citric acid, the last three yielding sodium citrate and sodium tartrate.

DOSAGE: 10 Gm. or 2½ drachms (U. S. P.).

***Sodii Salicylas** (*Sod. Salicyl.*), *Sodium Salicylate*, U. S. P.— $C_6H_4(OH)(COONa)$.

White or nearly white, odorless or nearly odorless powder having a sweet, saline taste. Very soluble in water (1:0.9) and freely soluble in alcohol (1:9.2). Incompatible with acids, with acid salts and with solutions of many of the alkaloids, particularly quinine, which precipitates as the salicylate.

ACTION AND USES: The salt usually employed to secure the constitutional action of salicylic acid.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Elixir Sodii Salicylatis (*Elix. Sod. Salicyl.*), *Elixir of Sodium Salicylate*, N. F.—Sodium salicylate (8.5%), syrup, water and aromatic elixir. Absolute alcohol content about 7 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Sodii Salicylatis Compositum (*Elix. Sod. Salicyl. Co.*), *Compound Elixir of Sodium Salicylate*, N. F.—Sodium salicylate (8%), fluidextract of cimifuga, (3.2%), fluidextract of gelsemium (1.6%) and potassium iodide (1.5%) in aromatic elixir. Absolute alcohol content about 23 per cent.

USES: An irrational rheumatism mixture, constituting an invitation to haphazard use.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Liquor Ferri Salicylatis (*Liq. Ferr. Salicyl.*), *Solution of Ferric Salicylate*, N. F. (Salicylated Mixture of Iron).—Solution containing ferric salicylate with a large excess of sodium salicylate [obtained by the interaction of sodium salicylate (12.5%) and tincture of ferric citrochloride (12.5%)], and ammonium citrate, methyl salicylate, glycerin and water.

USES: "Rheumatism remedy." Ferric salicylate has no special advantage over the sodium salt.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

***Sodii Sulphas** (*Sod. Sulph.*), *Sodium Sulphate*, U. S. (*Glauber's Salt*).— $Na_2SO_4 + 10H_2O$.

Colorless, odorless, efflorescent crystals, having a bitter taste. Freely soluble in water (1:1), but insoluble in al

ACTION AND USES: Saline cathartic, more disagreeable but less active than magnesium sulphate, over which it has no material advantage.

DOSAGE: 15 Gm. or 4 drachms (U. S. P.).

Sal Carolinum Factitium (Sal Carol. Fact.), Artificial Carlsbad Salt, N. F.—This is available in two forms, the dry amorphous form and the crystalline form, the latter representing about 80 per cent. of water of crystallization. The dry amorphous form represents potassium sulphate (2%), sodium chloride (18%), sodium bicarbonate (36%) and hydrated sodium sulphate (100%).

USES: Alkaline cathartic.

Sal Carolinum Factitium Effervescens (Sal Carol. Fact. Eff.), Effervescent Artificial Carlsbad Salt, N. F. (Pulvis Salis Carolini Factitii Effervescens, N. F. III).—Artificial Carlsbad salt (26.6%) in a mixture of sodium bicarbonate (40%), tartaric acid (15.7%) and citric acid, the last three yielding sodium citrate and sodium tartrate.

ACTION AND USES: Cathartic. Its high sodium citrate and tartrate content should be borne in mind.

DOSAGE: 6 Gm. or 1½ drachms (N. F.).

***Sodii Sulphis Exsiccatus (Sod. Sulphis Exsic.), Exsiccated Sodium Sulphite, U. S. P.**— Na_2SO_3 (not less than 90%).

White, odorless powder, having a cooling, saline sulphurous taste. Freely soluble in water (1:3.2), but only sparingly soluble in alcohol.

ACTION AND USES: Antiseptic; principally used externally as a wash.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

***Sodii Thiosulphas (Sod. Thiosulph.), Sodium Thiosulphate, U. S. P. ("Sodium Hyposulphite").**— $\text{Na}_2\text{S}_2\text{O}_3 + 5\text{H}_2\text{O}$.

Colorless, odorless crystals, having a cooling, afterward bitter taste. Very soluble in water (1:0.5) and insoluble in alcohol.

ACTION AND USES: Used externally in the form of a lotion as application for ringworm and other parasitic diseases. Internally, rarely as cathartic.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Solanum (Solan.), Solanum, N. F. (Horsenettle Berries).—Ripe fruit.

ACTION AND USES: Rarely used as a sedative in convulsions and epilepsy; probably ineffective.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Solani (*Fldext. Solan.*), *Fluidextract of Solanum*, N. F.—*Solanum* (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Sparteinae Sulphas (*Sparteinae Sulph.*), **Sparteine Sulphate**, U. S. P.

Colorless, odorless, hygroscopic crystals or powder, having a slightly saline and somewhat bitter taste. Freely soluble in water (1:1.1) and in alcohol (1:3).

ACTION AND USES: Has been widely exploited as a diuretic, but on insufficient evidence. Is of little use and is now less frequently employed than formerly. Large doses slow and weaken the heart.

DOSAGE: 0.01 Gm. or $\frac{1}{6}$ grain (U. S. P.).

SPECIES.—Mixtures of vegetable drugs used as teas or poultices. For the official mixtures, see *Species Emollientes*, N. F., under *Althaea Folia*; *Species Laxativae*, N. F., under *Senna*, and *Species Pectorales*, under *Althaea*.

Spigelia (*Spigelia*), **Spigelia**, U. S. P. (Pinkroot).—Rhizome and roots.

ACTION AND USES: Obsolete anthelmintic for ascarides. Inferior to *santonin*.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Fluidextractum Spigeliae (*Fldext. Spigel.*), *Fluidextract of Spigelia*, U. S. P. (*Fluidextract of Pinkroot*).—*Spigelia* (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

SPIRITUS, SPIRITS.—Alcoholic solutions of either gaseous, liquid, or solid volatile substances.

For the official spirits, see under the names of the respective principal constituents. For *Spiritus Cinnamomi*, see under *Oleum Cassiae*; for *Spiritus Odoratus*, under *Oleum Bergamottae*. For the following, see under the respective oils: *Spiritus Amygdalae Amarae*, *Spiritus Anisi*, *Spiritus Aurantii Compositus*, *Spiritus Cardamomi Compositus*, *Spiritus Juniperi*, *Spiritus Juniperi Compositus*, *Spiritus Lavandulae*, *Spiritus Menthae Piperitae*, *Spiritus Menthae Viridis*, *Spiritus Myrciae Compositus*, *Spiritus Sinapis*.

SPIRITUS OLEORUM VOLATILIUM, SPIRITS OF VOLATILE OILS.—Volatile oil (6.5%) in alcohol.

Staphisagria (Staphisag.), **Staphisagria**, U. S. P. (Stavesacre).—Seeds.

ACTION AND USES: To destroy pediculi. The contained alkaloid, delphinine, is toxic, resembling aconitine in its physiologic action.

DOSAGE: Used as an ointment or lotion.

Fluidextractum Staphisagria (Fldext. Staphisag.), *Fluidextract of Staphisagria*, U. S. P. (Fluidextract of Stavesacre).—Staphisagria (100%). Absolute alcohol content about 80 per cent.

STILI DILUBILES (STIL. DILUB.), **PASTE PENCILS**, (Unna Pencils).—Paste pencils are for the direct application of medicinal agents to the skin in dermatologic practice. The paste consists of starch, dextrin, tragacanth and sugar with sufficient water to form a plastic mass. For the official paste pencils, see *Stili Acidi Salicylici Dilubiles*, under *Acidum Salicylicum*.

Stillingia (Stilling.), **Stillingia**, U. S. P. (Queen's Root).

ACTION AND USES: Has been used as a sialagogue and expectorant. Has no advantage over senega; is now less frequently used than formerly.

The compound fluidextract and syrup are antiquated "alteratives," at one time used especially against syphilis and scrofula. They are complex and inefficient, and therefore irrational and harmful.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Fluidextractum Stillingiae (Fldext. Stilling.), *Fluidextract of Stillingia*, U. S. P.—Stillingia (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Fluidextractum Stillingiae Compositum (Fldext. Stilling. Co.), *Compound Fluidextract of Stillingia*, N. F.—Stillingia (25%), corydalis (25%), blue flag (12.5%), sambucus (12.5%), chimaphila (12.5%), prickly ash berries (6.2%) and coriander (6.2%) in glycerin, alcohol and water. Absolute alcohol content about 40 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Syrupus Stillingiae Compositus (Syr. Stilling. Co.), *Compound Syrup of Stillingia*, N. F.—Compound fluidextract of stillingia (25%) in glycerin and syrup. Absolute alcohol content about 10 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Stramonium** (*Stramon.*), **Stramonium**, U. S. P. (*Jamestown Weed*, *Jimson Weed*).—Leaves, containing not less than 0.25 per cent. of alkaloids.

ACTION AND USES: Similar to those of belladonna, over which it has no advantage. Often used in "asthma powders."

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

Extractum Stramonii (*Ext. Stramon.*), *Extract of Stramonium*, U. S. P.—Two forms: Pilular extract of stramonium and powdered extract of stramonium. One Gm. of extract represents about 4 Gm. of stramonium and yields about 1 per cent. of alkaloids.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.).

Fluidextractum Stramonii (*Fldext. Stramon.*), *Fluidextract of Stramonium*, N. F.—Stramonium (100%), yielding about 0.25 per cent. of alkaloids. Absolute alcohol content about 55 per cent.

DOSAGE: 0.05 Cc. or 1 minim (N. F.).

Tinctura Stramonii (*Tr. Stramon.*), *Tincture of Stramonium*, U. S. P.—Stramonium (10%), yielding about 0.025 per cent. of stramonium alkaloids. Absolute alcohol content about 40 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.).

Unguentum Stramonii (*Ung. Stramon.*), *Stramonium Ointment*, U. S. P.—Pilular extract of stramonium (10%), diluted alcohol, hydrous wool fat and benzoinated lard.

Strontii Bromidum (*Stront. Brom.*), **Strontium Bromide**, U. S. P.— $\text{SrBr}_2 + 6\text{H}_2\text{O}$.

Colorless, odorless crystals, having a bitter, saline taste. Very soluble in water (1: 0.35); soluble in alcohol.

ACTION AND USES: Similar to those of sodium bromide over which it has no definite advantage.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Strontii Carbonas (*Stront. Carb.*), **Strontium Carbonate**, N. F.— SrCO_3 .

White, odorless, tasteless powder. Insoluble in water and in alcohol. Decomposed by acids.

ACTION AND USES: Of use only in the making of salts of strontium.

Strontii Iodidum (*Stront. Iod.*), **Strontium**
— $\text{SrI}_2 + 6\text{H}_2\text{O}$.

Colorless crystals or white powder or crystalline crusts, turning yellow on exposure to light; odorless, deliquescent and having a bitter saline taste. Very soluble in water (1:0.2); somewhat less soluble in alcohol.

ACTION AND USES: Similar to, but without definite advantage over potassium iodide.

DOSAGE: 0.3 Gm. or 5 grains (U. S. P.).

Strontii Salicylas (Stront. Salicyl.), Strontium Salicylate, U. S. P.— $\text{Sr}(\text{C}_6\text{H}_4.\text{OH}.\text{COO})_2 + 2\text{H}_2\text{O}$.

White, odorless powder, having a somewhat sweet, saline taste. Soluble in water (1:19) and in alcohol (1:61).

ACTION AND USES: Identical with those of sodium salicylate, over which it has no advantage.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

***Strophanthinum (Strophanthin.), Strophanthin, U. S. P.**—A glucoside or mixture of glucosides obtained from strophanthus.

White or yellowish powder. Very soluble in water and in diluted alcohol.

ACTION AND USES: Like those of digitalis. Strophanthin acts more rapidly and is excreted sooner, but its absorption from the alimentary canal is so variable that its administration by mouth is inadvisable.

DOSAGE: Daily by mouth, 0.001 Gm. or $\frac{1}{60}$ grain (U. S. P.); intravenous, 0.00075 Gm. or $\frac{1}{80}$ grain (U. S. P.). Caution should be used if patient has been taking digitalis.

***Strophanthus (Strophanth.), Strophanthus, U. S. P.**—Ripe seeds.

ACTION AND USES: Strophanthus and tincture of strophanthus have properties similar to those of the glucoside strophanthin. The absorption of strophanthus from the alimentary canal is so variable that administration by mouth is not advisable.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

***Tinctura Strophanthi (Tr. Strophanth.), Tincture of Strophanthus, U. S. P.** (Strophanthi Tinctura, P. I.).—Strophanthus (10%) with the oil removed. Biologically assayed by producing death in frogs. Absolute alcohol content about 92 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.). Physicians who prescribe it should specify the biologically assayed preparation.

Strychnina (Strych.), Strychnine, U. S. P.—An alkaloid obtained from *nux vomica*.

Colorless, odorless crystals or white powder. Extremely bitter in very dilute solutions. Very slightly soluble in water (1: 6420) and slightly soluble in alcohol (1: 136).

ACTION AND USES: Stimulates the reflex activity of the spinal cord, but produces little or no effect on the higher nervous centers; is also used as a bitter tonic, generally in the form of a preparation of *nux vomica*.

As strychnine is so slightly soluble, it is generally prescribed as one of its salts. Of these, the nitrate meets every need.

DOSAGE: 0.0015 Gm. or $\frac{1}{40}$ grain (U. S. P.).

Elixir Pepsini, Bismuthi et Strychninae (Elix. Pepsin. Bism. et Strych.), Elixir of Pepsin, Bismuth and Strychnine, N. F.—Strychnine (0.0175%), tartaric acid, and elixir of pepsin and bismuth. Absolute alcohol content about 10 per cent.

USES: An irrational preparation.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Pilulae Antidyspepticae (Pil. Antidyspep.), Antidyspeptic Pills, N. F.—Each pill contains strychnine (0.0016 Gm., or $\frac{1}{40}$ grain), ipecac and pilular extract of belladonna leaves (each 0.0065 Gm., or $\frac{1}{40}$ grain), mass of mercury, and compound extract of colocynth (each 0.13 Gm., or 2 grains).

USES: An objectionably named and misdirected mixture.

DOSAGE: 1 pill (N. F.).

Strychninae Acetas, Strychnine Acetate.

Liquor Strychninae Acetatis (Liq. Strych. Acet.), Solution of Strychnine Acetate, N. F. (Hall's Solution of Strychnine).—Strychnine acetate (about 0.209%), alcohol, compound tincture of cardamom and water. Absolute alcohol content about 23 per cent.

USES: Superfluous strychnine solution.

DOSAGE: 0.6 Cc. or 10 minims (U. S. P.).

Strychninae Glycerophosphas (Strych. Glycerophos.), Strychnine Glycerophosphate, N. F. (Strychnine Glycerinophosphate).

White, odorless crystals or powder, with a taste at first sweet, afterward intensely bitter. Slightly soluble in water (1: 350), and in alcohol (1: 310).

ACTION AND USES: Superfluous strychnine salt.

DOSAGE: 0.0015 Gm. or $\frac{1}{40}$ grain (N. F.).

Strychninae Nitras (Strych. Nit.), Strychnine Nitrate, U. S. P.

Colorless, odorless crystals or white powder. Soluble in water (1:42) and somewhat less soluble in alcohol.

ACTION AND USES: Same as those of strychnine.

DOSAGE: 0.0015 Gm. or $\frac{1}{40}$ grain (U. S. P.).

Strychninae Sulphas (Strych. Sulph.), Strychnine Sulphate, U. S. P.

Colorless or white crystals or white powder, odorless; efflorescent in dry air. Soluble in water (1:32) and in alcohol (1:81).

USES: Same as those of strychnine. Has no advantage over the nitrate.

DOSAGE: 0.0015 Gm. or $\frac{1}{40}$ grain (U. S. P.).

Elixir Cinchonae Alkaloidorum, Ferri, Bismuthi et Strychninae (Elix. Cinchon. Ferr. Bism. et Strych.), Elixir of Cinchona Alkaloids, Iron, Bismuth and Strychnine, N. F. (Elixir Cinchonae, Ferri, Bismuthi et Strychninae, N. F. III, Elixir of Calisaya, Alkaloidal, with Iron, Bismuth and Strychnine).—Strychnine sulphate (0.0175%), elixir of cinchona alkaloids, iron and bismuth (about 99%) and water. Absolute alcohol content about 18 per cent.

USES: An irrational mixture supposed to contain six active ingredients.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Cinchonae Alkaloidorum, Ferri et Strychninae (Elix. Cinchon. Ferr. et Strych.), Elixir of Cinchona Alkaloids, Iron and Strychnine, N. F. (Elixir Cinchonae, Ferri et Strychninae, N. F. III, Elixir of Calisaya Alkaloidal, with Iron and Strychnine).—Strychnine sulphate (0.0175%), elixir of cinchona alkaloids and iron (about 99%) and water. Absolute alcohol content about 18 per cent.

USES: A complex and irrational "tonic."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Strychninae Valeras (Strych. Valer.), Strychnine Valerate, N. F.

White, crystalline powder, with a valerian-like odor intensely bitter taste. Sparingly soluble in water, soluble in

ACTION AND USES: Superfluous strychnine salt.

DOSAGE: 0.0015 Gm. or $\frac{1}{40}$ grain (N. F.).

Elixir Strychninae Valeratis (Elix. Strych. Valer.), *Elixir of Strychnine Valerate*, N. F. (Elixir Strychninae Valerianatis, N. F. III).—Strychnine valerate (0.0175%), aromatic elixir (about 96.5%), water, tincture of vanilla and compound tincture of cudbear. Absolute alcohol content about 22 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Styrax (Styrax), **Storax**, U. S. P. (Liquid Storax).—A balsam.

ACTION AND USES: Closely resembles balsam of Peru, over which it has no advantage. An ointment (1:4) has been used as a parasiticide in scabies and other parasitic infections.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Succus Citri (Suc. Cit.), **Lime Juice**, N. F.

ACTION AND USES: Used especially in prevention and treatment of scurvy.

Succus Pomorum (Suc. Pomor.), **Fresh Apple Juice**, N. F.

ACTION AND USES: Used in pharmacy in the making of crude ferric malate.

***Sulphonethylmethanum** (Sulphonethylmeth.), **Sulphonethylmethane**, U. S. P. (Trional).—Diethylsulphonemethylethylmethane.

Colorless, lustrous, odorless, crystalline scales, having a bitter taste in aqueous solutions. Slightly soluble in water (1:200) and in alcohol.

ACTION AND USES: A hypnotic producing in ordinary doses no other symptoms than sleep. The sleep comes on in about an hour, though in some cases it may be much longer delayed.

DOSAGE: 0.75 Gm. or 12 grains (U. S. P.), best administered in hot milk or other hot drinks.

***Sulphonmethanum** (Sulphonmeth.), **Sulphonmethane**, U. S. P. (Sulphonal).—Diethylsulphonedimethylmethane.

Colorless, odorless and nearly tasteless crystals or powder. Slightly soluble in water (1:365) and soluble in alcohol (1:60).

ACTION AND USES: Hypnotic and sedative. Has properties similar to those of sulphonethylmethane, but usually somewhat more slowly.

DOSAGE: 0.75 Gm. or 12 grains (U. S. P.). If administered in hot milk or other hot drinks.

***Sulphur Lotum (Sulphur Lot.), Washed Sulphur, U. S. P.—S.**

Fine, yellow, odorless, tasteless powder. Insoluble in water and in alcohol.

ACTION AND USES: Used locally in parasitic diseases of the skin and as a mild cathartic especially in hemorrhoids.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Trochisci Sulphuris et Potassii Bitartratis (Troch. Sulphur. et Pot. Bitart.), Troches of Sulphur and Potassium Bitartrate, N. F. (Troches of Sulphur and Cream of Tartar).—

Each troche contains washed sulphur (0.3 Gm. or 5 grains) and potassium bitartrate (0.06 Gm. or 1 grain) with tragacanth, sugar and oil of orange.

DOSAGE: 1 troche (N. F.). Dose so small as to be practically worthless.

***Sulphur Praecipitatum (Sulphur Praec.), Precipitated Sulphur, U. S. P. (Lac Sulphuris, Milk of Sulphur).—S.**
Made by precipitating a solution of calcium sulphide with hydrochloric acid.

Fine, pale yellow, odorless, tasteless powder.

ACTION AND USES: Same as those of washed sulphur; more active and irritant.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

***Sulphur Sublimatum (Sulphur Sublim.), Sublimed Sulphur, U. S. P. (Flowers of Sulphur).—S.**

Fine, yellow powder having a slight, characteristic odor and a faintly acid taste.

ACTION AND USES: Same as those of washed sulphur.

DOSAGE: 4 Gm. or 1 drachm (U. S. P.).

Petroxolinum Sulphuratum (Petrox. Sulphur.). Sulphurated Petroxolin, N. F. (Sulphurated Petrox).—Sublimed sulphur (3%) in linseed oil, oleic acid and liquid petroxolin.

Petroxolinum Sulphuratum Compositum (Petrox. Sulphur. Co.), Compound Sulphurated Petroxolin, N. F. (Compound Sulphurated Petrox).—Sulphurated petroxolin (10%), in oil of cade, thymol, eucalyptol, oil of turpentine and liquid petroxolin.

***Unguentum Sulphuris (Ung. Sulphur.), Sulphur Ointment, U. S. P.**—Sublimed sulphur (15%) in benzoinated lard.

Unguentum Sulphuris Alkalinum (Ung. Sulphur. Alk.), Alkaline Sulphur Ointment, N. F.—Sublimed sulphur (20%) and potassium carbonate (10%) in water and benzoinated lard.

Unguentum Sulphuris Compositum (*Ung. Sulphur. Co.*), *Compound Sulphur Ointment*, N. F. (Wilkinson's Ointment, Hebra's Itch Ointment).—Sublimed sulphur (15%), oil of cade (15%) and precipitated calcium carbonate in soft soap and lard.

Sulphuris Iodidum (*Sulphur. Iod.*), **Sulphur Iodide**, N. F.—Washed sulphur (20%) and iodine combined by means of heat.

Brittle, grayish black masses, having a metallic luster, an odor of iodine and a somewhat acrid taste. Almost insoluble in water; alcohol dissolves the iodine, leaving the sulphur.

ACTION AND USES: Used in parasitic skin diseases; antiquated.

Sumbul (*Sumbul*), **Sumbul**, U. S. P. (*Musk-root*).—Rhizome and roots.

ACTION AND USES: Formerly exploited as a stimulant and antispasmodic in hysterical conditions. Effect probably psychic.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Extractum Sumbul (*Ext. Sumbul*), *Extract of Sumbul*, U. S. P.—A pilular extract.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Fluidextractum Sumbul (*Fldext. Sumbul*), *Fluidextract of Sumbul*, U. S. P. (*Fluidextract of Musk-root*).—Sumbul (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Tinctura Sumbul (*Tr. Sumbul*), *Tincture of Sumbul*, N. F.—Sumbul (10%) in alcohol and water. Absolute alcohol content about 60 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

SUPPOSITORIA, SUPPOSITORIES, U. S. P.—Suppositories are solid bodies of various weights and shapes adapted for introduction into the different orifices of the human body and melting readily at body heat.

Suprarenalum Siccum (*Supraren. Sicc.*), **Dried Suprarenals**, U. S. P. (*Glandulae Suprarenales Siccae*, U. S. P., VIII, *Desiccated Suprarenal Glands*).—The suprarenal glands of animals which are used for food by man, cleaned, dried, freed from fat, and powdered, and containing epinephrine (about 0.5%), the active principle of the suprarenal

yellowish-brown powder, having a slight, characteristic fatty soluble in water.

ACTION AND USES: The action of dried suprarenals is virtually that of epinephrine, but more variable and less conveniently employed.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

SYRUPI, SYRUPS.—Syrups are aqueous solutions of medicinal substances rendered palatable by the addition of sugar. For the official syrups see under the names of the principal ingredients. For *Syrupus*, see under *Saccharum*; for *Syrupus Bromidorum*, under *Potassii Bromidum*; for *Syrupus Ficorum Compositus*, under *Ficus*; for *Syrupus Hypophosphitum* and *Syrupus Hypophosphitum Compositus*, under *Calcii Hypophosphis*; for *Syrupus Ipecacuanhae et Opii*, under *Opium*; for *Syrupus Phosphatum Compositus*, under *Acidum Phosphoricum*; for *Syrupus Phosphatum cum Quinina et Strychnina*, under *Quininae Hydrochloridum*, and for *Syrupus Pini Strobi Compositus cum Morphina*, under *Morphinae Sulphas*.

Talcum Purificatum (Talc. Purif.), Purified Talc, U. S. P.—

A purified native hydrous magnesium silicate, sometimes containing a small amount of aluminum silicate.

A fine white or nearly white, odorless, tasteless powder, which adheres to the skin and is slippery to the touch.

ACTION AND USES: Used as a dusting powder, and as a clarifying agent in pharmacy.

Tamarindus (Tamarind.), Tamarind, N. F.—The preserved pulp of the fruit.

ACTION AND USES: Mildly laxative. When infused with water, yields an agreeable acid drink.

DOSAGE: 15 Gm. or 4 drachms (N. F.).

Taraxacum (Tarax.), Taraxacum, U. S. P. (Dandelion).—Rhizome and roots.

ACTION AND USES: Has been used as a bitter tonic (without advantage over gentian) and as a mild laxative in habitual constipation (inferior to drugs like *cascara sagrada*); has no specific action on the liver.

DOSAGE: 10 Gm. or 2½ drachms (U. S. P.).

Elixir Taraxaci Compositum (Elix. Tarax. Co.), Compound Elixir of Taraxacum, N. F.—Fluidextract of taraxacum (3.5%), fluidextract of wild cherry (2%), fluidextract of glycyrrhiza, tincture of sweet orange peel, tincture of cinnamon, compound tincture of cardamom and aromatic elixir. Absolute alcohol content, about 28 per cent.

DOSAGE: 8 Cc. or 2 fluidrachms (N. F.).

Extractum Taraxaci (Ext. Tarax.), *Extract of Taraxacum*, U. S. P.—A pilular extract.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Fluidextractum Taraxaci (Fldext. Tarax.), *Fluidextract of Taraxacum*, U. S. P. (Fluidextract of Dandelion).—Taraxacum (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

Terebenum (Tereben.), **Terebene**, U. S. P.—Obtained from oil of turpentine by the action of concentrated sulphuric acid.

A colorless, thin liquid, having a thyme-like odor and an aromatic taste. Only slightly soluble in water; soluble in alcohol (1:3).

ACTION AND USES: Expectorant resembling oil of turpentine, but more pleasant in odor.

DOSAGE: 0.25 Cc. or 4 minims (U. S. P.).

***Terebinthina** (Terebinth.), **Turpentine**, N. F.—A solid oleo-resin obtained from pine.

ACTION AND USES: Rubefacient; without advantage over turpentine oil.

Terebinthina Laricis (Terebinth. Laric.), **Venice Turpentine**, N. F. (Larch Turpentine).—A viscid oleoresin obtained from European larch.

ACTION AND USES: Similar to those of turpentine; now seldom used.

Petroxolinum Terebinthinae Laricis (Petrox. Terebinth. Laric.), *Venice Turpentine Petroxolin*, N. F. (Venice Turpentine Petrox).—Venice turpentine (20%) in liquid petroxolin.

***Terpini Hydras** (Terpin. Hyd.), **Terpin Hydrate**, U. S. P.

Colorless, lustrous, nearly odorless crystals, having a slightly aromatic and somewhat bitter taste. Slightly soluble in water (1:200) and soluble in alcohol (1:13).

ACTION AND USES: Antiseptic, diaphoretic, diuretic; largely used as an expectorant in cases accompanied by free bronchial secretion.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Elixir Terpini Hydratis (Elix. Terpin. Hyd.), *Elixir Terpin Hydrate*, N. F.—Terpin hydrate (1.75%), sweet orange peel, spirit of bitter almond, alcc

syrup and water. Absolute alcohol content about 42 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Elixir Terpini Hydratis et Codeinae, N. F.—See under Codeina.

Elixir Terpini Hydratis et Diacetylmorphinae, N. F.—See under Diacetylmorphina.

Terra Silicea Purificata (Ter. Sil. Purif.), Purified Siliceous Earth, U. S. P. (Purified Kieselguhr, Purified Infusorial Earth).—Silica (SiO_2).

Fine, bulky, white or nearly white, odorless, tasteless powder. Insoluble in water, acids or dilute solutions of the alkalis.

ACTION AND USES: Used as a clarifying agent in pharmacy.

***Theobrominae Sodio-Salicylas** (Theobrom. Sodio-Sal.), Theobromine Sodio-Salicylate, U. S. P. (Diuretin).—Sodium theobromine and sodium salicylate in approximately molecular proportions. Contains not less than 46.5 per cent. of theobromine.

White, odorless powder, having a sweetish, saline and somewhat alkaline taste. Freely soluble in water (1:1), slightly soluble in alcohol. Incompatible with acids and has the other incompatibilities of salicylates.

ACTION AND USES: A soluble salt of theobromine, having much the same action as caffeine, except that it does not affect the central nervous system to any extent. Mainly used as a diuretic.

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

***Theophyllina** (Theophyll.), Theophylline, U. S. P. (Dimethyl-xanthine, Theocin).—An isomer of theobromine.

White, odorless powder, having a bitter taste. Soluble in water (1:100) and in alcohol (1:80).

ACTION AND USES: Has a diuretic action similar to that of theobromine, more powerful, but said to be not so lasting.

DOSAGE: 0.25 Gm. or 4 grains (U. S. P.).

Thuja (Thuja), Thuja, N. F. (Arbor Vitae).—Young twigs.

ACTION AND USES: Somewhat irritant; no well-defined action. Recommended as a febrifuge and as a stimulating agent in the aged.

Fluidextractum Thujae (*Fldext. Thuj.*), *Fluidextract of Thuja*, N. F.—Thuja (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Thymol (Thymol), Thymol, U. S. P.**—A phenol.

Large, colorless, translucent crystals, having an aromatic thyme-like odor, and a pungent, aromatic taste. Very slightly soluble in water (1:1,010); freely soluble in alcohol (1:1).

ACTION AND USES: Antiseptic and anthelmintic. Used chiefly against the hookworm.

DOSAGE: Antiseptic, 0.125 Gm. or 2 grains (U. S. P.); anthelmintic, 1 Gm. or 15 grains, per day (U. S. P.). In the treatment of hookworm disease it should be given in as finely divided state as possible in dosage of from 0.5 to 4 Gm. or from 8 grains to 1 drachm. No fats, oils or alcohol should be given at the same time, for fear of absorption of the drug.

The dosage may be regulated according to age as follows: Up to 5 years of age, 0.5 Gm. or 8 grains; up to 10, 1 Gm. or 15 grains; up to 15, 1.5 Gm. or 24 grains; up to 20, 2 Gm. or 30 grains; above 20, 3 to 4 Gm. or 45 grains to 1 drachm.

Nebula Thymolis (*Nebul. Thymol.*), *Thymol Spray*, N. F.—Thymol (1%) in light liquid petrolatum.

***Thymolis Iodidum (Thymol. Iod.), Thymol Iodide, U. S. P.** (Aristol).—Chiefly dithymol-diiodide. Should contain not less than 43 per cent. of iodine.

A reddish-brown or reddish-yellow, bulky powder, with a very slight aromatic odor. Insoluble in water or glycerin; only slightly soluble in alcohol.

ACTION AND USES: Antiseptic; used chiefly as a dusting powder. Less efficient than iodoform.

Thymus (Thymus), Thyme, N. F.—Tops.

ACTION AND USES: Aromatic and rubefacient; without advantage over camphor.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Thymi (*Fldext. Thym.*), *Fluidextract of Thyme*, N. F.—Thyme (100%). Absolute alcohol content about 20 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Thyroideum Siccum (Thyroid. Sicc.), Dried Thyroids, U. S. P.** (*Glandulae Thyroideae Siccæ*, U. S. P., VIII, *Desiccated Thyroid Glands*).—The thyroid glands of ani-

mals which are used for food by man, freed from connective tissue and fat, dried and powdered. Should contain about 0.2 per cent. of iodine.

A yellowish powder, with a slight characteristic odor.

ACTION AND USES: Used in thyroid deficiency; sometimes in obesity, in which it may be harmful.

DOSAGE: 0.1 Gm. or $1\frac{1}{2}$ grains (U. S. P.).

TINCTURAE, TINCTURES.—Tinctures with few exceptions are alcoholic or hydroalcoholic extractive preparations of vegetable drugs. The tinctures of potent drugs represent uniformly 10 Gm. of drug in 100 Cc. of the preparation, while tinctures of less potent drugs vary in strength but represent usually 20 Gm. of drug in 100 Cc. of the preparation. For the official tinctures see under the names of the respective drugs. For *Tinctura Amara*, see under *Gentiana*; for *Tinctura Antiperiodica* and *Tinctura Antiperiodica sine Aloe*, under *Quininae Bisulphas*; for *Tinctura Aromatica*, under *Cinnamomum Saigonicum*; for *Tinctura Ipecacuanhae et Opii* and for *Tinctura Pectoralis*, under *Opium*.

TINCTURAE AETHEREAE, ETHEREAL TINCTURES, N. F.—When not otherwise directed, ethereal tinctures are to be prepared representing 10 per cent. drug strength, extracted with a mixture of alcohol and ether. Of doubtful advantage over ordinary tinctures.

TINCTURAE MEDICAMENTORUM RECENTIUM, TINCTURES OF FRESH DRUGS, N. F. (*Tincturae Herbarum Recentium*, U. S. P. VIII).—When not otherwise directed, tinctures of fresh drugs are to be prepared by extracting 50 Gm. of the fresh drug with alcohol to make 100 Cc.

TOXITABELLAE, POISON TABLETS.—Tablets containing poisons.

Tragacantha (Trag.), Tragacanth, U. S. P. (*Gum Tragacanth*).—Mixed with 50 parts of distilled water, forms a smooth, nearly uniform, stiff, opalescent mucilage.

ACTION AND USES: Demulcent and ingredient in emulsions.

Glyceritum Tragacanthae (*Glycer. Trag.*), *Glycerite of Tragacanth, N. F.*—Tragacanth (about 10%) in glycerin and water.

Mucilago Tragacanthae (*Mucil. Trag.*), *Mucilage of Tragacanth, U. S. P.*—Tragacanth (6%) in glycerin (18%) and water.

Trifolium (Trifol.), Trifolium, N. F. (Red Clover Blossoms).

ACTION AND USES: Inert. Has been used in antisyphilitic and alterative compounds; also as an expectorant.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Trifolii (Fldext. Trifol.), Fluidextract of Trifolium, N. F.—Trifolium (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Trillium (Trill.), Trillium, N. F. (Beth Root).—Rhizome.

ACTION AND USES: Obsolete, formerly used as astringent and tonic.

DOSAGE: 2 Gm. or 30 grains (N. F.).

Fluidextractum Trillii (Fldext. Trill.), Fluidextract of Trillium, N. F.—Trillium (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

***Trinitrophenol (Trinitrophen.), Trinitrophenol, U. S. P. (Picric Acid).**

Pale yellow, odorless crystals, having an intensely bitter taste, and exploding when heated rapidly and when subjected to percussion. Trinitrophenol stains the skin an intense, permanent yellow. Soluble in water (1:78), and in alcohol (1:12).

ACTION AND USES: For the dressing of burns. Internally is highly toxic, producing nausea, vomiting and diarrhea, and stains the skin and mucous membranes a yellow color, simulating jaundice.

DOSAGE: Externally as saturated watery solution or as ointment.

Triticum (Tritic.), Triticum, U. S. P. (Couch Grass, Dog Grass).—Rhizome and roots.

ACTION AND USES: Formerly exploited as a diuretic in cystitis and irritable bladder. Now seldom employed.

DOSAGE: 8 Gm. or 2 drachms (U. S. P.).

Fluidextractum Tritici (Fldext. Tritic.), Fluidextract of Triticum, U. S. P. (Fluidextract of Couch Grass).—Triticum (100%). Absolute alcohol content about 15 per cent.

DOSAGE: 10 Cc. or 2½ fluidrachms (U. S. P.).

TRITURATIONES, TRITURATIONS.—M:

ing active drugs in powder form; or
substance (10%) with powdered

official trituration see Trituratio Elaterini, U. S. P., under *Elaterinum*.

TROCHISCI, TROCHES (Lozenges).—Tablets intended to be dissolved in the mouth for their local effect on the mucous membrane of the mouth and the throat. For the official troches see under the names of the principal ingredients.

Ulmus (*Ulmus*), Elm, U. S. P. (Elm Bark, Slippery Elm).—The inner bark.

ACTION AND USES: Obsolete demulcent.

Trochisci Ulmi (*Troch. Ulmi*), *Troches of Elm*, N. F.—Each troche contains elm, tragacanth, sugar and methyl salicylate.

USES: Demulcent expectorant.

DOSAGE: 1 troche (N. F.).

UNGUENTA, OINTMENTS.—Soft, fatty solids, of such consistency that they are readily spread at ordinary temperatures. For the official ointments, see under the names of the respective principal constituents. For Unguentum, see under *Adeps*; for Unguentum Diachylon, under *Plumbi Oxidum*, and for Unguentum Fuscum, under *Camphora*.

Uranii Nitras (*Uran. Nit.*), Uranium Nitrate, U. S. P. (*Uranyl Nitrate*).— $\text{UO}_2(\text{NO}_3)_2 + 6\text{H}_2\text{O}$.

Light, yellow, odorless, somewhat efflorescent, radioactive crystals, having a bitter astringent taste. Freely soluble in water (1:1.2); also soluble in alcohol or ether.

ACTION AND USES: Has been used, without adequate justification, in the treatment of diabetes and cancer. Solutions are poisonous and produce glucosuria when injected subcutaneously, even in small doses.

DOSAGE: 0.01 Gm. or $\frac{1}{8}$ grain (U. S. P.), best administered in simple solutions. Use with caution.

Uva Ursi (*Uva Ursi*), *Uva Ursi*, U. S. P. (Bearberry).—Leaves.

ACTION AND USES: Mild and slightly antiseptic diuretic used especially in vesical catarrh. Less effective than santal oil.

GE: 2 Gm. or 30 grains (U. S. P.).

-tum Uvae Ursi (*Flidext. Uvae Ursi*), *Fluidextract Ursi*, U. S. P.—*Uva ursi* (100%). Absolute \dagger about 25 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

***Valeriana (Valer.), Valerian, U. S. P.**—Rhizome and roots.

ACTION AND USES: Used as "antispasmodic" and "nerve sedative" in hysteria and other nervous excitations. Its influence is largely psychic, owing to its strong, persistent odor.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Fluidextractum Valerianae (Fldext. Valer.), Fluidextract of Valerian, N. F.—Valerian (100%). Absolute alcohol content about 70 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Valerianae (Tr. Valer.), Tincture of Valerian, U. S. P.—Valerian (20%), in alcohol and water. Absolute alcohol content about 69 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (U. S. P.).

**Tinctura Valerianae Ammoniata (Tr. Valer. Ammon.), Ammoniated Tincture of Valerian, U. S. P.*—Valerian (20%) in aromatic spirit of ammonia. Absolute alcohol content about 64 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Vanilla (Vanilla), Vanilla, N. F. (Vanilla Bean).—Cured fruit.

ACTION AND USES: Flavoring.

Tinctura Vanillae (Tr. Vanill.), Tincture of Vanilla, N. F.—Vanilla, sugar and diluted alcohol. Absolute alcohol content about 40 per cent.

Vanillinum (Vanillin.), Vanillin, U. S. P.—The odorous principle of vanilla, sometimes prepared synthetically.

Fine, white or nearly white, crystalline needles having the odor and taste of vanilla. Soluble in water (1:100) and freely soluble in alcohol or glycerin.

ACTION AND USES: Used only as a flavoring.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

Elixir Vanillini Compositum (Elix. Vanil. Co.), Compound Elixir of Vanillin, N. F.—Compound spirit of vanillin (2%), alcohol, glycerin, syrup, tincture of caramel and water. Absolute alcohol content about 8 per cent.

Spiritus Vanillini Compositus (Sp. Vanil. Co.), Compound Spirit of Vanillin, N. F.—Vanillin, oil of orange, oil of cardamom, oil of cinnamon and alcohol. Absolute alcohol content about 80 per cent.

Veratrina (Veratrin.), Veratrine, U. S. P. (Veratria).—A mixture of alkaloids from *sabadilla* seeds.

White or nearly white, odorless powder, intensely irritating to the nasal mucous membrane. Very slightly soluble in water (1:1760) and freely soluble in alcohol (1:2.8).

ACTION AND USES: In its action on peripheral nerve endings, it is said to resemble aconite, but has no material advantage over the latter. Has been applied (in an ointment) as analgesic in neuralgia. It is highly toxic, and caution should be observed in its use, especially where the skin is broken.

Oleatum Veratrinae (Oleat. Veratrin.), Oleate of Veratrine, N. F.—Veratrine (2%) in oleic acid and olive oil.

Unguentum Veratrinae (Ung. Veratrin.), Veratrine Ointment, N. F.—Veratrine (4%) in expressed oil of almond and benzoinated lard.

Veratrum Viride (Verat. Vir.), Veratrum Viride, U. S. P. (Green Hellebore, American Hellebore).—Rhizome and roots.

ACTION AND USES: Slows the heart and lowers blood pressure. The tincture has been recommended especially in eclampsia, but is now little used.

DOSAGE: 0.06 Gm. or 1 grain (U. S. P.).

*Fluidextractum Veratri Viridis (Fldext. Verat. Vir.), Fluidextract of Veratrum Viride, U. S. P. (Fluidextract of Green Hellebore).—*Veratrum viride (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 0.1 Cc. or 1½ minims (U. S. P.).

Tinctura Veratri Viridis (Tr. Verat. Vir.), Tincture of Veratrum Viride, U. S. P.—Veratrum viride (10%). Absolute alcohol content about 91 per cent.

DOSAGE: 0.5 Cc. or 8 minims (U. S. P.)

Verbasci Flores (Verbasc. Flor.), Mullein Flowers, N. F.—Corollas with stamens.

ACTION AND USES: Demulcent, without advantage over acacia, elm, etc.

DOSAGE: 8 Gm. or 2 drachms (N. F.).

Verbasci Folia (Verbasc. Fol.), Mullein Leaves, N. F.

ACTION AND USES: Same as those of the flowers.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Verbasci Foliae [should be Foliorum] (Fldext. Verbasc. Fol.), Fluidextract of Mullein Leaves.

N. F.—Mullein leaves (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Verbena (Verben.), Verbena, N. F. (Blue Vervain).—Plant deprived of its root.

ACTION AND USES: Obsolete; said to be tonic, emetic, expectorant and diaphoretic.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Fluidextractum Verbenae (Fldext. Verben.), Fluidextract of Verbena, N. F.—Verbena (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 1 Cc. or 15 minims (N. F.).

Viburnum Opulus (Viburn. Opul.), Viburnum Opulus, N. F. (Crampbark, High Bush Cranberry Bark).

ACTION AND USES: Slightly bitter; practically inert; was recommended as tonic, antispasmodic and alterative. The market supply is said to have been spurious for many years, consisting of mountain maple (*Acer spicatum*).

DOSAGE: 2 Gm. or 30 grains (N. F.).

Elixir Viburni Opuli Compositum (Elix. Viburn. Opul. Co.), Compound Elixir of Viburnum Opulus, N. F. (Compound Elixir of Crampbark).—Fluidextract of viburnum opulus (7.5%), fluidextract of trillium (15%), fluidextract of aletris (7.5%) and compound elixir of taraxacum. Absolute alcohol content about 35 per cent.

USES: An inefficient mixture sold for use in "female weakness."

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Fluidextractum Viburni Opuli (Fldext. Viburn. Opul.), Fluidextract of Viburnum Opulus, N. F.—Viburnum opulus (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 2 Cc. or 30 minims (N. F.).

Tinctura Viburni Opuli Composita (Tr. Viburn. Opul. Co.), Compound Tincture of Viburnum, N. F.—Viburnum opulus (3.5%), dioscorea (3.5%) and scutellaria (1%), with clove and Saigon cinnamon in glycerin, alcohol and water. Absolute alcohol content about 80 per cent.

USES: Irrational mixture used in menstrual disorders.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Viburnum Prunifolium (Viburn. Prun.), **Viburnum Prunifolium**, U. S. P. (Black Haw, Viburnum).—The dried bark.

ACTION AND USES: Has had considerable vogue as a "uterine sedative," in dysmenorrhea and habitual abortion. There is no good evidence that it has any action.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Elixir Viburni Prunifolii (Elix. Viburn. Prun.), *Elixir of Viburnum Prunifolium*, N. F. (Elixir of Black Haw).—Fluidextract of viburnum prunifolium (12.5%), compound tincture of cardamom and aromatic elixir. Absolute alcohol content about 29 per cent.

USES: An objectionable alcoholic preparation of viburnum.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Extractum Viburni Prunifolii (Ext. Viburn. Prun.), *Extract of Viburnum Prunifolium*, U. S. P. (Powdered Extract of Viburnum Prunifolium).—One Gm. extract represents 5 Gm. viburnum prunifolium.

DOSAGE: 0.5 Gm. or 8 grains (U. S. P.).

Fluidextractum Viburni Prunifolii (Fldext. Viburn. Prun.), *Fluidextract of Viburnum Prunifolium*, U. S. P. (Fluidextract of Black Haw).—Viburnum prunifolium (100%). Absolute alcohol content about 55 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

VINA, WINES.—Medicated wines are solutions of active medicaments or the active constituents of drugs in wine. For the official wines, see under the names of the respective principal constituents.

Vinum Xericum (Vin. Xeric.), **Sherry Wine**, N. F.—An alcoholic liquid made by fermenting the juice of fresh ripe grapes, the fruit of cultivated varieties of *Vitis*, freed from seeds, stems and skins, and fortifying with pure grape brandy. Absolute alcohol content about 20 per cent.

ACTION AND USES: Sherry wine has the action of alcohol.

***Virus Vaccinicum** (Virus Vaccin.), **Vaccine Virus**, U. S. P. (Glycerinated Vaccine Virus, Smallpox Vaccine, Jennerian Vaccine).—Prepared from the pustules of vaccinia from vaccinated cattle. The product must comply with the requirements established by the Secretary of the Treasury of the United States through the United States Public Health Service.

USES: Prophylactic vaccination against smallpox

Xanthoxylum (Xanthox.), Xanthoxylum, U. S. P. (Prickly Ash Bark).

ACTION AND USES: Bitter, without special advantage over gentian.

DOSAGE: 2 Gm. or 30 grains (U. S. P.).

Fluidextractum Xanthoxyli (Fldext. Xanthox.), Fluidextract of Xanthoxylum, U. S. P. (Fluidextract of Prickly Ash).—Xanthoxylum (100%). Absolute alcohol content about 60 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

Xanthoxylum Fructus (Xanthox. Fruct.), Prickly Ash Berries, N. F.

ACTION AND USES: Obsolete and probably worthless; has been used as stimulant, tonic, alterative, diaphoretic, sialagogue and carminative.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Zea (Zea), Zea, N. F. (Corn Silk).

ACTION AND USES: Probably valueless. Has been used in inflammatory conditions of the bladder.

DOSAGE: 4 Gm. or 1 drachm (N. F.).

Fluidextractum Zeae (Fldext. Zea.) Fluidextract of Zea, N. F.—Zea (100%). Absolute alcohol content about 40 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Zedoaria (Zedoar.), Zedoary, N. F.—Rhizome.

ACTION AND USES: Practically the same as those of ginger, over which it has no advantage.

DOSAGE: 1 Gm. or 15 grains (N. F.).

Tinctura Zedoariae Amara (Tr. Zedoar. Amar.), Bitter Tincture of Zedoary, N. F. (Compound Tincture of Zedoary).—Zedoary (25%), aloes (12.5%), rhubarb (6.2%), gentian (6.2%), agaric (6.2%) and saffron (6.2%) in glycerin, alcohol and water. Absolute alcohol content about 50 per cent.

USES: Bitter stomachic and laxative; needlessly cor and without special advantage over compound tinc gentian.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

***Zinci Acetas (Zinc. Acet.), Zinc Acetate, U. S. P.—**
 $\text{Zn}(\text{C}_2\text{H}_3\text{O}_2)_2 \cdot 2\text{H}_2\text{O}$.

Soft white crystals, having a faint vinegar odor and, in dilute solutions, an astringent metallic taste. Freely soluble in water (1:2.3), and soluble in alcohol (1:30).

ACTION AND USES: Used locally, like zinc sulphate, being somewhat less powerful.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Zinci Carbonas Praecipitatus (Zinc. Carb. Praec.), Precipitated Zinc Carbonate, U. S. P.

Impalpable, white, odorless, tasteless powder. Insoluble in water or alcohol.

ACTION AND USES: Similar to, but without advantage over zinc oxide.

***Zinci Chloridum (Zinc Chlor.), Zinc Chloride, U. S. P.**
 ZnCl_2 .

White or nearly white, granular powder, porcelain-like masses or molded pencils, odorless, very deliquescent and intensely caustic. Very soluble in water (1:0.25) and in alcohol (1:1.3).

ACTION AND USES: Antiseptic, astringent and escharotic.

***Liquor Zinci Chloridi (Liq. Zinc. Chlor.), Solution of Zinc Chloride, U. S. P.—** ZnCl_2 (about 50 per cent.).

***Zinci Oxidum (Zinc. Oxid.), Zinc Oxide, U. S. P.—** ZnO .

Fine, white or nearly white, odorless, tasteless powder. Insoluble in water or in alcohol.

ACTION AND USES: Antiseptic and astringent, widely used either alone or in combination with other substances as a dusting powder and as a protective and sedative in ointments.

Glycerogelatinum Zinci Durum (Glycerogel. Zinc. Dur.), Firm Zinc Glycerogelatin, N. F.—Zinc oxide (10%) in water, glycerin and glycerinated gelatin.

Glycerogelatinum Zinci Molle (Glycerogel. Zinc. Mol.), Soft Zinc Glycerogelatin, N. F.—zinc oxide (10%) in water, glycerin and glycerinated gelatin.

Mulla Zinci (Mull. Zinc.), Zinc Mull, N. F. (Unguentum Zinci Extensum, N. F. III).—Zinc oxide (10%) in benzoinated suet and benzoinated lard.

Pasta Zinci (Past. Zinc.), Zinc Paste, N. F. (Lassar's Zinc Paste, N. F.).—Zinc oxide (24%), salicylic acid (2%), starch

Moll.), Soft Zinc Paste, N. F.—Zinc oxide (25%), pre- and solution of

Pasta Zinci Sulphurata (*Past. Zinc. Sulphur.*), *Sulphurated Zinc Paste*, N. F. (Unna's Sulphurated Zinc Paste).—Zinc oxide (15%), precipitated sulphur (10%), purified siliceous earth and benzoinated lard.

**Unguentum Zinci Oxidi* (*Ung. Zinc. Ox.*), *Ointment of Zinc Oxide*, U. S. P. (Zinc Ointment).—Zinc oxide (20%) in benzoinated lard.

Zinci Phenolsulphonas (**Zinc. Phenolsulph.**), **Zinc Phenolsulphonate**, U. S. P. (Zinc Sulphocarbolate).—The hydrated salt.

Colorless, odorless crystals or granules having an astringent, metallic taste. Freely soluble in water (1:1.6), and in alcohol (1:1.8).

ACTION AND USES: Similar to, but less active than zinc sulphate, over which it has no advantage.

DOSAGE: 0.125 Gm. or 2 grains.

***Zinci Stearas** (**Zinc. Stear.**), **Zinc. Stearate**, U. S. P.—Chiefly zinc stearate and palmitate.

Fine, bulky, white, tasteless powder, having a faint characteristic odor. Insoluble in water or alcohol.

ACTION AND USES: Similar to, but without special advantage over zinc oxide.

Unguentum Zinci Stearatis (*Ung. Zinc. Stear.*), *Ointment of Zinc Stearate*, N. F.—Zinc stearate (50%) in white petrolatum.

USES: Without advantage over the oxide.

***Zinci Sulphas** (**Zinc. Sulph.**), **Zinc Sulphate**, U. S. P.— $\text{ZnSO}_4 + 7\text{H}_2\text{O}$.

Colorless, transparent crystals or granular powder, odorless and having an astringent, metallic taste. Very soluble in water (1:0.6) and freely soluble in glycerin (1:2.5); insoluble in alcohol.

ACTION AND USES: Astringent, styptic and emetic. Much used in eye washes and especially effective in that form of conjunctivitis caused by the Morax-Axenfeld bacillus.

DOSAGE: Emetic, 1 Gm. or 15 grains (U. S. P.) in diluted solution. Locally, 0.1 to 1 per cent. in collyria; 0.5 to 4 per cent. in injections (gonorrhea).

Liquor Zinci et Alumini Compositus (*Liq. Zinc. et Alumin. Co.*), *Compound Solution of Zinc and Aluminum*, N. F.—Zinc sulphate (20%), aluminum sulphate (20%), betanaphthol (0.06%) with oil of thyme in water.

USES: Needlessly complex astringent and antiseptic.

Liquor Zinci et Ferri Compositus (Liq. Zinc. et Ferr. Co.), *Compound Solution of Zinc and Iron*, N. F. (Deodorant Solution).—Zinc sulphate (20%), ferrous sulphate (20%), copper sulphate (6.5%), betanaphthol (0.06%) with oil of thyme and hypophosphorous acid in water.

USES: Needlessly complex antiseptic mixture.

Pulvis Antisepticus (Pulv. Antisept.), *Soluble Antiseptic Powder*, N. F. (Pulvis Antisepticus Solubilis).—Zinc sulphate (12.5%), salicylic acid (0.5%), phenol, eucalyptol, menthol and thymol (each 0.1%) and boric acid.

USES: Needlessly complex antiseptic mixture.

Zinci Valeras (Zinc. Valer.), *Zinc Valerate*, U. S. P. (Zinc Valerianate).

White scales or powder, having a valerian-like odor and a sweetish, astringent, metallic taste. Soluble in water (1:70), often leaving a residue, and in alcohol (1:22).

ACTION AND USES: Formerly deemed a "nerve sedative," but is without value.

DOSAGE: 0.125 Gm. or 2 grains (U. S. P.).

Elixir Zinci Valeratis (Elix. Zinc. Valer.), *Elixir of Zinc Valerate*, N. F. (Elixir Zinci Valerianatis, N. F. III).—Zinc valerate (1.75%), ammonium citrate, alcohol, spirit of bitter almond, compound tincture of cudbear, water and aromatic elixir. Absolute alcohol content about 28 per cent.

DOSAGE: 4 Cc. or 1 fluidrachm (N. F.).

Zincum (Zinc.), *Zinc*, U. S. P.—Zn.

Bluish-white, thin sheets, irregular, granulated pieces, molded pencils or powder. Insoluble in water or alcohol, but dissolves in diluted sulphuric or hydrochloric acid with the evolution of hydrogen.

ACTION AND USES: Reagent for the generation of hydrogen.

**Zingiber* (Zingib.), *Ginger*, U. S. P.—Rhizomes.

ACTION AND USES: Flavor, carminative, aromatic and stimulant to the gastro-intestinal tract, because of the irritating action of the volatile oil and resin. Probably of advantage over other pungent aromatics, such as the

DOSAGE: 1 Gm. or 15 grains (U. S. P.).

Fluidextractum Zingiberis (Fldext. Zingib.), *Fluid Ginger*, U. S. P.—Ginger (100%). Absolute alcohol content about 80 per cent.

DOSAGE: 1 Cc. or 15 minims (U. S. P.).

Oleoresina Zingiberis (*Oleores. Zingib.*), *Oleoresin of Ginger*, U. S. P.

USES: Highly irritant.

DOSAGE: 0.03 Gm. or $\frac{1}{2}$ grain (U. S. P.).

Syrupus Zingiberis (*Syr. Zingib.*), *Syrup of Ginger*, U. S. P.

Fluidextract of ginger (3%), in alcohol (2%) and syrup.

DOSAGE: 15 Cc. or 4 fluidrachms (U. S. P.).

**Tinctura Zingiberis* (*Tr. Zingib.*), *Tincture of Ginger*, U. S. P. (Tincture of Jamaica Ginger).—Jamaica ginger (20%) in alcohol. Absolute alcohol content about 91 per cent.

DOSAGE: 2 Cc. or 30 minims (U. S. P.).

INDEX

In this index preference has been given to the English names of the various drugs. The Latin names are used only when they differ materially in spelling from the English.

	PAGE		PAGE
Absinthium	5	Adhesive plaster—Continued	
Absorbent cotton (Gossypium purificatum)	103	rubber (Emplastrum elasticum)	182
Acacia	5	Adjuvans elixir (Elixir glycyrrhizae)	19
and morphine syrup	134	Adonis	16
mucilage	5	fluidextract	16
syrup	5	Aether	16
Aceta	5	aceticus	17
Acetanilid	5	nitrosus	17
powder, compound	6	Aethylis carbamas	17
Acetic acid	6	chloridum	17
acid, diluted	6	Aethylmorphinae hydrochloridum	17
acid, glacial	6	dum	17
ether	17	African chillies (Capsicum)	55
turpentine liniment	148	Agar	18
Acetomorphine (Diacetylmorphina)	78	Agar-Agar	18
Acetone	6	Agaric	18
Acetphenetidin	6	larch (Agaricus)	18
Acetum aromaticum	142	white (Agaricus)	18
opii	149	Aitken tonic pills (pillulae ferri, quinae, strychninae et arseni mites)	95
scillae	192	Albuminate of iron, solution	94
Acid, acetic acid, boric acid, citric acid, etc.: See Acetic acid, Boric acid, Citric acid, etc.		Alcohol	18
Aconite	14	absolutum (Alcohol dehydratum)	18
and chloroform liniment	15	dehydrated	18
extract	15	diluted	18
fluidextract	15	Aleppo galls (Galla)	98
powdered extract (Extractum aconiti)	15	Aletris	19
root	14	fluidextract	19
tincture	15	Alexandria senna or India senna (Senna)	194
Aconiti tuber (Aconitum)	14	Alkaline antiseptic solution	169
Aconitine	14	mixture of rhubarb	183
oleate	14	solution of tar	165
Actaea, compound syrup (Syrupus cimicifugae compositus)	64	sulphur ointment	215
Adeps	15	Allium	19
benzoinatus	15	Alspice (Pimenta)	163
lanae	15	oil (Oleum Pimentae)	146
lanae hydrosus	16	Almond, bitter oil	137
Adhesive plaster (Emplastrum resinae)	182	bitter, spirit (Spiritus Amygdalae amarae)	138
rosin (Emplastrum resinae)	182	bitter, water	137
		elixir, compound	137
		emulsion	29
		salik (Emulsio amygdalae)	29

	PAGE		PAGE
Almond—Continued		Ammonia—Continued	
oil, expressed	133	water	24
oil, sweet (Oleum amygdalae		water, stronger	24
expressum)	133	Ammoniated camphor wash	25
sweet	28	glycyrrhizin	103
Aloes	19	mercury	113
and asafetida pills	20	mercury ointment	113
and canella powder	21	tincture of ergot	82
and iron pills	20	tincture of gualac	105
and mastic pills	20	tincture of valerian	224
and myrrh pills	20	Ammonio-ferric citrate (Ferri et	
and myrrh tincture	21	ammonii citras)	87
and podophyllum compound		acetate	25
pills	21	acetate and iron solution	25
extract	20	acetate solution	25
iron, quinine and nux vomica		benzoate	25
pills	94	bromide	26
mercury and podophyllum		bromide elixir	26
pills	21	carbonate	26
mercury and scammony com-		chloride	27
pound pills	21	chloride mixture	27
pills	20	chloride troches	27
powdered extract (Extractum		citrate	27
aloes)	20	citrate and bismuth	41
tincture	21	citrate and iron	87
Aloin	21	citrate solution	27
compound pills	22	hypophosphite	27
strychnine and belladonna		hypophosphite syrup	27
compound pills	22	iodide	28
strychnine and belladonna		iodide liniment	28
pills	22	phosphate	28
Alum	23	salicylate	28
burnt (Alumen exsiccatum) ..	23	valerate	28
dried (Alumen exsiccatum) ..	23	valerate elixir	28
exsiccated	23	valerianate (Ammonii valeris)	28
Alumen	23	Amygdala dulcis	28
ustum (Alumen exsiccatum) ..	23	Amyl nitrite	29
Aluminum acetate	23	Amylum	29
acetate solution	23	Anethol	29
acetico-tartrate	24	Angelica fruit	29
acetico-tartrate solution	24	root	29
and zinc compound solution ..	230	root, fluidextract	30
chloride	24	seed (Angelicae fructus)	29
hydroxide	24	Anhydrous lanolin (Adeps lanae)	15
subacetate	24	Anisated powder of rhubarb and	
subacetate solution	24	magnesia	184
sulphate	24	solution of ammonia (Spiritus	
Althaea	22	ammoniae anisatus)	25
leaves	23	spirit of ammonia	25
syrup	22	Anise	30
American hellebore (Veratrum		elixir	29
viride)	225	oil (Oleum anisi)	138
spikenard (Aralla)	32	powder, compound (Pulvis	
wormseed oil (Oleum cheno-		Rhei et magnesia anisatus) ..	184
podii)	140	spirit	138
Aminoform (Hexamethylena-		water	138
mina)	107	Aniseed (Anisum)	30
Ammonia	24	Anisum	30
liniment	25	Antidiphtheric globulins (Serum	
solution, anisated (Spiritus		antidiphthericum purifica-	
ammoniae anisatus)	25	tum)	196
spirit, anisated	25	serum	196
spirit, aromatic	26	serum, dried	196

PAGE	PAGE
Antidiphtheric—Continued	Apple juice, fresh..... 214
serum, purified 196	Apples, ferrated extract..... 90
Antidyspeptic pills 212	ferrated extract, tincture..... 90
Antifebrin (Acetanilidum) 5	Aqua
Antimonial powder 30	See also under Water.
Antimonium oxysulphuratum	ammoniae 24
(Antimonium sulphuratum).	ammoniae fortior 25
sulphuratum 30	amygdalae amarae..... 137
Antimony and potassium tartrate	anisi 138
compound pills 31	aqua 32
oxide 30	aurantii florum 37
sulphurated 30	aurantii florum fortior..... 37
tartrated (Antimonii et potas-	camphorae 53
sii tartras) 30	chloroformi 62
wine 30	cinnamomi 140
Antiperiodic pills..... 180	creosoti 75
pills without aloes..... 180	destillata 32
tincture 177	destillata sterilisata 32
tincture without aloes..... 178	foeniculi 141
Antipyrine 31	hamamelidis 106
Antiseptic powder, soluble..... 231	hydrogenii dioxidii (Liquor
solution 7	hydrogenii dioxidii)..... 115
solution, alkaline 169	menthae piperitae 143
solution of pepsin 157	menthae viridis 143
Antitetanic globulins (Serum	phagedaenica flava (Lotio
antitetanicum purificatum)..... 196	flava)..... 111
serum 196	phagedaenica nigra (Lotio
serum, dried 197	nigra)..... 109
serum, purified 196	phenolata 159
Antitoxin, diphtheria (Serum	regia (Acidum nitrohydro-
antidiphthericum) 196	chloricum) 10
diphtheria, concentrated (Se-	rosae 185
rum antidiphthericum puri-	rosae fortior 185
ficatum) 196	sedativa (Lotio ammoniacalis
diphtheria, dried (Serum anti-	camphorata) 25
diphthericum siccum)..... 196	Aquae aromaticae 32
diphtheria, refined and con-	Aqueous elixir of glycyrrhiza..
centrated (Serum antidiph-	elixir of licorice (Elixir gly-
thericum purificatum) 196	cyrrhizae aquosum) 102
globulins, diphtheric (Serum	extract of ergot 82
antidiphthericum purifica-	fluidextract of cinchona..... 65
tum) 196	tincture of rhubarb..... 184
globulins, tetanus (Serum anti-	Aralla 32
tetanicum purificatum).... 196	fluidextract 32
tetanus (Serum antitetani-	Arbor vitae (Thuja)..... 219
cum) 196	Argenti nitras 32
tetanus, concentrated (Serum	nitras fusus 33
antitetanicum purificatum)..... 196	oxidum 33
tetanus, dried (Serum anti-	Aristol (Thymolis iodidum).... 220
tetanicum siccum) 197	Arnica 33
tetanus, refined and concen-	flowers (Arnica) 33
trated (Serum antitetani-	fluidextract 33
cum purificatum) 196	tincture 33
Ants, spirit of (Spiritus acidi	Aromatic camphor mixture..... 142
formici) 8	castor oil 146
Apil fructus 31	elixir 18
Aplol, liquid (Oleo-resina petro-	elixir of eriodictyon 83
selini) 159	elixir of glycyrrhiza 19
Apocynum 31	elixir of licorice (Elixir gly-
fluidextract 31	cyrrhizae aromaticum)..... 19
Apomorphine chloride (Apomor-	elixir of yerba santa (Elixir
phinae hydrochloridum).... 31	eriodictyl aromaticum)..... 83
hydrochloride 31	fluidextract 86

	PAGE		PAGE
Aromatic—Continued		Asarum	34
fluidextract of cascara sagrada	58	syrup, compound (Syrupus asari compositus)	34
fluidglycerate of cascara sagrada	58	Asclepias	35
oil spray	158	fluidextract	35
powder	66	Aspidium	35
powder of chalk	66	oleoresin	35
powder of chalk and opium	151	Aspidosperma	35
solution of pepsin	157	fluidextract	35
spirit of ammonia	26	Astringent mixture	77
sulphuric acid	13	Atophan (Acidum phenylephedrinicum)	11
syrup of eriodictyon	83	Atropine	35
syrup of rhubarb	184	oleate	36
syrup of senna	195	sulphate	36
syrup of yerba santa (Syrupus eriodictyli aromaticus)	83	Aurantii amari cortex	36
tincture	66	dulcis cortex	36
tincture of rhubarb	184	flores	37
vinegar	142	Auri et sodii chloridum	37
waters	32	Baccae spinæ cervinæ (Rhamnus cathartica)	182
Aromatized Iodoform	117	Baking soda (Sodii bicarbonas)	199
Aromatol (Nebula aromatica)	158	Balm of Gilead buds (Populi gemmae)	167
Arsenas sodii (Sodii arsenas)	198	Balsam of copaiba (Copaiba) of Peru	73
Arseni iodidum	33	of Tolu	37
trioxidum	33	poplar buds	167
Arsenic antidote (Ferri hydroxidum cum magnesii oxido)	88	Balsamum tranquillans (Oleum hyoscyami compositum)	115
bromide and gold solution	7	Baptisia	38
chloride, solution (Liquor acidi arsenosi)	34	fluidextract	38
iodide (Arseni iodidum)	33	Barker's post partum pills (Pillulae laxative post partum)	71
iron, quinine and strychnine pills, mild	95	Basham's mixture (Liquor ferri et ammonii acetatis)	25
iron, quinine and strychnine pills, stronger	95	Basilicon ointment (Ceratum resinae)	181
solution, Clemens'	168	Bateman's pectoral drops (Tinctura pectoralis)	152
solution, hydrochloric (Liquor acidi arsenosi)	34	Bay oil (Oleum myrciae)	145
trioxide	33	Bayberry bark	135
white (Arseni trioxidum)	33	powder, compound	135
Arsenous acid (Arseni trioxidum)	33	Bearberry (Uva ursi)	223
acid, solution	34	Beef	56
and mercuric iodide solution	33	and wine (Vinum carnis)	56
iodide	33	and iron, wine	57
oxide (Arseni trioxidum)	33	extract (Extractum carnis)	56
Artificial Carlsbad salt	207	wine	56
Carlsbad salt, effervescent	207	wine and iron (Vinum carnis et ferri)	57
Kissingen salt	202	Beeswax (Cera flava)	59
Kissingen salt, effervescent	202	Belladonna, aloin and strychnine pills	22
Vichy salt	189	aloin and strychnine pills, compound	22
Vichy salt, effervescent	200	leaves	38
Vichy salt with lithium, effervescent	200	leaves, extract	38
Asafoetida	34	leaves, tincture	38
and aloes pills	20	liniment	39
emulsion	34	ointment	38
gum (Asafoetida)	34	plaster	38
milk (Emulsio Asafoetidae)	34		
opium and magnesia mixture	128		
pills	34		
tincture	34		

PAGE	PAGE
Belladonna—Continued	Bismuth—Continued
root	strychnine, cinchona alkaloids,
root, fluidextract	and iron, elixir..... 213
tincture (Tinctura belladonnae	subcarbonate
foliorum)	42
Benne oil (Oleum sesami)....	subgallate
147	42
Benzaldehyde	subnitrate
39	42
Benzamine hydrochloride (Beta-	subsallylate
eucalae hydrochloridum)...	42
40	Bitter almond oil..... 137
Benzinum purificatum	almond spirit
39	138
Benzoic acid	almond water
7	137
Benzoin	apple (Colocynthis)
39	70
tincture	metallic pills (Pilulae ferri,
39	quininae, strychninae et ar-
tincture, compound	sen] fortiores)
39	95
Benzoinated lard	orange elixir
15	138
suet	orange flowers
197	37
Benzosulphinide	orange oil
40	138
Berberis	orange peel
40	36
fluidextract	orange peel, fluidextract.....
40	36
Bergamot, oil	orange peel, tincture.....
139	36
Bestuscheff's tincture (Tinctura	stomachic drops (Tinctura
ferri chloridi aetherea)....	amara)
87	100
Betaeucaline hydrochloride	tincture
40	100
Betanaphthol	tincture of zedoary.....
40	228
paste	wine of iron
40	88
petrox (Petroxolinum beta-	wood (Quassia)
naphtholis)	176
41	Bitterless syrup of quinidine
petroxolin	(Syrupus quinidinae)
41	177
Beth root (Trillium)	Bittersweet
222	80
Bichloride of mercury (Hydrar-	fluidextract
gyri chloridum corrosivum). ..	80
108	tablets (Toxatabellae hydrar-
gyri chloridi corrosivi)....	109
109	Biniodide of mercury (Hydrar-
gyri iodidum rubrum)	110
110	Birch, rectified empyroligneous
oil (Oleum betulae empyreu-	maticum rectificatum)
139	139
tar oil, rectified.....	139
Rismuth ammoniochlorate (Bis-	muth et ammonii chloras)....
41	and ammonium citrate
41	and pepsin, elixir.....
156	and sodium tartrate.....
41	betanaphthol
41	betanaphtholate (Bismuthi
betanaphtholas)	41
elixir	41
glycerite	41
iron and callosa, alkaloidal	elixir (Elixir cinchonae al-
41	kaloïdorum, ferri et bis-
41	muthi)
41	iron and cinchona alkaloids,
41	elixir
42	magma
42	milk (Magma bismuthi)....
42	oxide
42	solution
42	strychnine, and pepsin, elixir.
212	strychnine, callosa, alka-
212	loidal, and iron, elixir.....
213	

	PAGE		PAGE
Blood root—Continued		Buchu—Continued	
tincture (Tinctura sangul-		long	44
narlae)	189	short	44
Blue cohosh (Caulophyllum)...	59	Buckbean (Menyanthes)	133
flag	120	Buckthorn bark (Frangula)....	97
flag, fluidextract	120	bark, fluidextract (Fluidex-	
gum leaves (Eucalyptus).....	83	tractum frangulae)	97
mass (Massa hydrargyri).....	112	berries (Rhamnus Cathartica)	182
ointment (Unguentum hydrar-		berries, syrup (Syrupus rham-	
gyri dilutum)	112	ni catharticae)	183
pill (Massa hydrargyri)	112	Burdock root (Lappa)	124
vervaln (Verbena)	226	Burnt alum (Alumen exsicca-	
Boldo	43	tum)	23
fluidextract	43	sugar coloring (Caramel)....	55
leaves	43	Burow's solution (Liquor alu-	
Boneset (Eupatorium)	84	mini acetatis)	23
Boric acid (Acidum boricum)..	7	Butter of cacao (Oleum theo-	
Borax (Sodii boras)	200	bromatis)	148
and honey (Mel sodii boratis)	200	Butternut bark (Juglans).....	120
with honey of rose (Mel rosae			
et sodii boratis)	200	Cabbage rose (Rosa)	185
Boric acid	7	Cacao butter (Oleum theobro-	
acid, ointment	7	matis)	148
Boroglycerin, glycerite	7	prepared	44
suppositories	101	Cactus grandiflorus	45
Boro-Salicylated powder of talc		cactus grandiflorus, tincture	45
(Pulvis talci compositus)...	12	Cade, oil	139
Boulton's solution (Liquor iodi		petrox (Petroxollinum cadini).	139
phenolatus)	118	petroxollin	139
Brayera	43	Caffeine	45
infusion	43	citrated	45
Breast tea (Species pectorales).	22	citrated, effervescent	45
Bromauric acid	7	sodio-benzoate	45
Bromide and chloral, compound		sodio-salicylate	45
(Mistura chloralis et potas-		with effervescent potassium	
sii bromidi composita).....	61	bromide (Sal potassii bro-	
Bromides and potassium arsen-		midi effervescens compositus)	
ate solution (Liquor arseni-		calis clemens)	170
callis clemens)	168	Cajuput, oil	139
syrup	170	Calabar bean (Physostigma)...	161
three, elixir	26	bean, tincture (Tinctura phys-	
Bromine	43	ostigmatis)	162
solution	43	Calamine ointment	46
solution, Smith's (Liquor		prepared	46
bromi)	43	Calcined magnesla (Magnesi	
Bromoforn	43	oxidum)	129
Broom tops (Scoparius).....	193	Calcium and sodium glycerophosphates, elixir	47
Brown mixture (Mistura glycyrrhizae composita)	150	and sodium hypophosphates	
mustard (Sinapis nigra)....	197	syrup	48
ointment	53	bromide	46
plaster, camphorated	53	bromide, elixir	46
Bryonia	44	carbonate, precipitated	46
tincture	44	chloride	47
Bryony (Bryonia)	44	cinchona alkaloids and iron	
Buchu	44	lactophosphate elixir	49
and potassium acetate, elixir.	44	glycerinophosphate (Calcii	
elixir	44	glycerophosphas)	47
elixir, compound	44	glycerophosphate	47
fluidextract	44	hydrochlorophosphate, syrup..	50
fluidextract, compound	44	hydroxide, solution	52
leaves	44	hypophosphite	47
		hypophosphite elixir	48

	PAGE
Calcium—Continued	
hypophosphite syrup	48
iodide	49
iodide syrup	49
lactate	49
lactophosphate	49
lactophosphate and cod liver oil emulsion	144
lactophosphate and iron syrup	50
lactophosphate, cinchona alkaloids and iron elixir	50
lactophosphate elixir	49
lactophosphate syrup	50
oxide	52
oxysulphuret, solution (Liquor calcis sulphuratae)	51
phosphate and cod liver oil emulsion	144
phosphate, precipitated	50
sulphide, crude	50
Calendula	51
flowers	51
fluidextract	51
tincture	51
Calisaya, alkaloidal elixir (Elixir cinchonae alkaloidorum)	180
alkaloidal, ferrated elixir (Elixir cinchonae alkaloidorum et ferri)	92
alkaloidal, with hypophosphites, elixir (Elixir cinchonae alkaloidorum et hypophosphitum)	180
alkaloidal, with iron and pepsin, elixir (Elixir cinchonae alkaloidorum, ferri et pepsini)	156
alkaloidal, with iron and strychnine, elixir (Elixir cinchonae alkaloidorum, ferri et strychninae)	213
alkaloidal, with iron, bismuth and strychnine, elixir (Elixir cinchonae alkaloidorum, ferri, bismuthi, et strychninae)	213
bark (Cinchona)	64
iron and lactophosphate of lime, elixir (Elixir of cinchona alkaloids, iron and calcium lactophosphate)	50
Calomel (Hydrargyri chloridum mite)	109
and jalap (Pulvis hydrargyri chloridi mltis et jalapae)	110
and santonin troches (Trochisci santonini compositi)	190
Calumba	51
fluidextract	51
tincture	51
Calx	52
chlorinata	52

	PAGE
Calx—Continued	
sulphurata (Calcii sulphidum crudum)	50
Cambogia	52
Campho-menthol (Menthol camphoratum)	133
Camphor	52
and chloroform petrox (Petroxolinum chloroformi camphoratum)	62
and menthol (Menthol camphoratum)	133
and opium pills	151
cerate	53
liniment	53
mixture, acid	149
mixture, aromatic	142
mixture, Parrish's (Mistura camphorae aromatica)	142
monobromated	53
ointment	53
spirit	53
wash, ammoniated (Lotio ammoniacalis camphorata)	25
water	53
Camphorated brown plaster	53
chloral	61
chloroform petroxolin	62
menthol	133
motherplaster (Emplastrum fuscum camphoratum)	53
oil (Linimentum camphorae)	53
phenol petrox (Petroxolinum phenolis camphoratum)	160
phenol petroxolin	160
soap liniment	190
tincture of opium	152
Canada liniment (Linimentum opii compositum)	149
snake-root (Asarum)	34
Canadian hemp (Apocynum)	31
hemp, fluidextract (Fluidextractum apocyni)	31
Canella	53
and aloes powder	21
Cannabis	54
extract	54
fluidextract	54
indica	54
tincture	54
Cantharidal collodion	54
Cantharides	54
cerate	51
plaster	54
tincture	55
Capsicum	55
and myrrh, tincture	55
oleoresin	55
plaster	55
tincture	55
Caramel	55
tincture	55

	PAGE		PAGE
Caraway	57	Cataplasmata	59
oil	140	Cataria	59
Carawayseed (Carum)	57	Cathartic elixir, compound	97
Carbamate, ethyl	17	pills, compound	71
Carbamic acid ethyl ester		pills, vegetable	71
(Aethylis carbamas)	17	Catmint (Cataria)	59
Carbo ligni	55	Catnep	59
Carbolic acid (Phenol)	159	fluidextract	59
acid, glycerite (Glyceritum		Caulophyllum	59
phenolis)	159	fluidextract	59
acid, iodized (Phenol ioda-		Caustic potash (Potassii hy-	
tum)	160	droxidum)	172
acid, liquefied (Phenol lique-		soda (Sodii hydroxidum)	203
factum)	160	Cayenne pepper (Capsicum)	55
acid, ointment (Unguentum		Celery fruit	31
phenolis)	160	fruit, fluidextract	31
acid water (Aqua phenolata).		seed (ApII fructus)	31
Carbonized oil (Oleum phenola-		Centauri	59
tum)	159	Cera alba	59
solution of iodine (Liquor iodi		flava	59
phenolatus)	118	Cerate	59
Carbonate, strontium	210	camphor	53
Carbonis, pix (Pix lithanthracis)		cantharides	54
Cardamom, elixir, compound...	140	of lead subacetate	167
oil	139	simple (Ceratum)	59
seed	56	Cerates	59
spirit, compound	140	Ceratum resinae	181
tincture	56	resinae compositum	182
tincture, compound	56	Cerevisiae fermentum compres-	
Carlsbad salt, artificial	207	sum	60
salt, artificial, effervescent...	207	Cerium oxalate	60
Carminative, Dalby's (Mistura		Cetaceum	60
carminativa)	150	Ceylon cinnamon	66
mixture	150	Chalk and opium aromatic pow-	
Carmine	56	der	151
solution	56	drop (Creta praeparata)	75
Caro	56	mixture	76
Carrageen (Chondrus)	62	powder, aromatic	66
Carron oil (Linimentum calcis).		powder, compound	76
Carum	57	precipitated (Calci carbonas	
Caryophyllus	57	praecipitatus)	46
Cascara sagrada	57	prepared	75
elixir	57	with mercury	112
elixir, compound	57	Chalybeate pills (Pilulae ferri	
extract	57	carbonatis)	86
extract, powdered (Extractum		Chamomile, German (Matri-	
cascarae sagradae)	57	carla)	131
fluidextract	58	wild (Matricaria)	131
fluidextract, aromatic	58	Channing's solution (Liquor hy-	
fluidglycerate	58	drargyri et potassii iodidi).	110
fluidglycerate, aromatic	58	Chapman's dinner pills	20
Cascarilla	58	mixture (Mistura copalibae et	
Cassia fistula	58	opii)	73
oil (Oleum cassiae)	140	Charcoal (Carbo ligni)	55
Castanea	58	troches	56
Castile soap, white (Sapo)	190	Charta potassii nitratis	173
Castor oil	146	sinapis (Emplastrum sinapis).	197
aromatic	146	Chartae	60
emulsion	146	Chemical food (Syrupus phos-	
Cataplasma emollient (Species		phatum compositum)	11
emollientes)	23	Chenopodium, oil	140
of kaolin	121	Chestnut leaves (Castanea)	53
		leaves, fluidextract	53

	PAGE		PAGE
Chillies, African (<i>Capsicum</i>)...	55	Cinchona	64
Chimaphila	60	alkaloids and hypophosphites,	
fluidextract	60	elixir	180
Chionanthus	60	alkaloids and iron, elixir.....	92
fluidextract	60	alkaloids, elixir	180
Chirata	61	alkaloids, iron and bismuth,	
fluidextract	61	elixir	41
Chiretta (<i>Chirata</i>)	61	alkaloids, iron and calcium	
Chloral (<i>Chloralum hydratum</i>)..	61	lactophosphate, elixir	49
and bromide compound (<i>Mis-</i>		alkaloids, iron and pepsin	
tura chloralis et potassii		elixir	156
bromidi composita)	61	alkaloids, iron and strychnine	
and potassium bromide mix-		elixir	213
ture, compound	61	alkaloids, iron, bismuth and	
camphorated	61	strychnine elixir	213
hydrate (<i>Chloralum hydra-</i>		extract	64
tum)	61	fluidextract	64
hydrated	61	fluidextract, aqueous	65
chloride of lime (<i>Calx chlori-</i>		infusion	65
nata)	52	red	65
Chlorinated lime	52	tincture	65
potassa	167	tincture, compound	65
potassa, solution	167	yellow (<i>Cinchona</i>)	64
soda	197	Cinchonidine sulphate	65
soda solution	197	Cinchonine sulphate	65
Chlorine	62	Cineol (<i>Eucalyptol</i>)	83
solution, compound	62	Cinnamomum saigoncum	66
water (<i>Liquor chlori composi-</i>		zeylanicum	66
tus)	62	Cinnamon, Ceylon	66
Chloroform	61	oil	140
and aconite liniment.....	15	saigon	66
and morphine mixture, com-		spirit	140
pound	62	syrup	66
and opium mixture, compound	150	tincture	66
anodyne (<i>Mistura chloroformi</i>		water	140
et morphinae composita)...	62	Citrated caffeine	45
camphor and petrox (<i>Petroxo-</i>		caffeine, effervescent	45
linum chloroform cam-		Citric acid	8
phoratum)	62	acid syrup	8
liniment	62	Citrine ointment (<i>Unguentum</i>	
petroxolin, camphorated	62	hydrargyri nitratis)	111
spirit	62	Clarified honey	132
water	62	Clemens' solution of arsenic...	168
Chlorum	62	Clove	57
Chondrus	62	Cloves, oil (<i>Oleum caryophylli</i>)..	140
gelatin	63	Coal tar	165
mucilage	63	solution	165
Chromic acid (<i>Chromii trioxi-</i>		Cocaine	67
dum)	63	chloride (<i>Cocaina hydro-</i>	
anhydride (<i>Chromii trioxidum</i>)	63	chloridum)	67
Chromium trioxide	63	hydrochloride	67
Chrysarobin	63	oleate	67
ointment	63	Cocculus indicus	67
Churchill's tincture of iodine		indicus tincture	67
(<i>Tinctura iodi fortior</i>).....	119	Coccus	67
Cimicifuga	63	Cochia pills (<i>Pilulae colocyn-</i>	
extract	63	thidis compositae)	71
extract, powdered (<i>Extractum</i>		Cochineal	67
cimicifugae)	63	color	67
fluidextract	63	Cocillana	67
syrup, compound	64	fluidextract	68
tincture	64	Cocoa	68
		soluble (<i>Cacao praeeparata</i>)..	44

	PAGE		PAGE
Cod liver oil	144	Colocynth—Continued	
emulsion	144	extract, powdered, compound	
with calcium lactophosphate,		(Extractum colocynthis	
emulsion	144	compositum)	70
with calcium phosphate, emul-		pills, compound	71
sion	144	pulp (Colocynthis)	70
with egg, emulsion	145	Colombo (Calumba)	51
with hypophosphites, emulsion	144	Colophony (Resina)	181
with malt, emulsion	144	Colorless hydrastine solution	
with phosphate of lime, emul-		(Liquor hydrastinae compo-	
sion (Emulsum olei morr-		situs)	113
huae cum calcii phosphate) ..	144	Coltsfoot	85
with wild cherry, emulsion ..	144	leaves (Farfara)	85
Codeine	68	Columba (Calumba)	51
and terpin hydrate, elixir	68	Composition powder (Pulvis my-	
phosphate	68	ricae compositus)	135
sulphate	68	Compound acetanilid powder ..	6
syrup	68	anise powder (Pulvis rhel et	
Coffee	69	magnesiae anisatus)	184
fluidextract	69	cathartic elixir	97
roasted (Coffee tosta)	69	cathartic pills	71
Cola (Kola)	122	chalk powder	76
nuts (Kola)	122	chloral and bromide (Mistura	
Colchici semen	69	chloralis et potassii bromidi	
tinctura (Tinctura colchici		composita)	61
seminis)	70	croton oil liniment	149
Colchicine	70	decoction of sarsaparilla	191
Colchicum corm	69	effervescent salt of potassium	
corm extract	69	bromide	170
corm, extract, powdered (Ex-		effervescing powder	172
tractum colchici cormi)	69	elixir of almond	137
corm fluidextract	69	elixir of blackberry	186
corm wine	69	elixir of buchu	44
root (Colchici cormus)	69	elixir of cardamon	140
seed	69	elixir of cascara sagrada	57
seed, fluidextract	69	elixir of corydalis	74
seed tincture	70	elixir of crampbark (Elixir vi-	
seed wine	70	burni opuli compositum)	226
Cold cream (Unguentum aquae		elixir of formates	8
rosae)	186	elixir of glycerophosphates ..	203
Cole's dinner pills	20	elixir of orange (Vinum au-	
Colic root (Aletris)	19	rantii compositum)	36
root (Dioscorea)	79	elixir of pepsin and rennin	156
Collodion	176	elixir of sodium salicylate	206
blistering (Collodium canthar-		elixir of taraxacum	217
idatum)	54	elixir of vanillin	124
cantharidal	54	elixir of viburnum opulus	226
croton oil	148	extract of colocynth	70
flexible	176	extract of colocynth, powdered	
iodine	118	(Extractum colocynthis	
iodoform	117	compositum)	70
salicylic, compound	12	fluidextract of buchu	44
styptic	13	fluidextract of sarsaparilla ..	191
vesicating (Collodium can-		fluidextract of stillingiae	209
tharidatum)	54	gargle of gualac	105
Collodions	70	infusion of gentian	100
Colocynth	70	infusion of rose	186
and hyoscyamus pills	71	infusion of senna	194
and podophyllum pills	71	iron mixture	86
apple (Colocynthis)	70	laxative pills	22
extract	70	licorice powder (Pulvis gly-	
extract, compound	70	cyrrhizae compositus)	194
extract, powdered	70		

	PAGE
Compound—Continued	
liniment of mustard	147
liniment of opium	149
liniment of soft soap	190
menthol liniment	133
menthol spray	133
mixture of chloral and potas- sium bromide	61
mixture of chloroform and morphine	62
mixture of glycyrrhiza	150
mixture of opium and chloro- form	150
mixture of opium and rhubarb	150
mixture of rhubarb	183
oil of hyoscyamus	115
pancreatin powder	153
pills of aloes and podophyllum	21
pills of aloes, mercury and scammony	21
pills of aloin	22
pills of aloin, strychnine and belladonna	22
pills of antimony	31
pills of colocynth	71
pills of rhubarb	184
powder of bayberry	135
powder of gambir	98
powder of glycyrrhiza	194
powder of ipecac (Pulvis Ipe- cacuanhae et opii)	151
powder of jalap	120
powder of kino and opium	122
powder of rhubarb	184
powder of talc	12
resorcinol ointment	182
rosin cerate	182
salicylate collodion	12
solution of chlorine	62
solution of cresol	75
solution of glycerophosphates (Elixir glycerophosphatum compositum)	203
solution of hydrastine	113
solution of hypophosphites	48
solution of iodine	118
solution of phosphates	11
solution of sodium borate	200
solution of sodium phosphate	205
solution of zinc and aluminum	230
solution of zinc and iron	231
spirit of cardamon	140
spirit of ether	16
spirit of juniper	142
spirit of lavender (Tinctura lavandulae composita)	142
spirit of myrcia	145
spirit of orange	138
spirit of vanilla	224
sulphur ointment	216
sulphurated petrox (Petroxoli- num sulphuratum composi- tum)	215

	PAGE
Compound—Continued	
sulphurated petroxolin	215
syrup of actaea (Syrupus cimicifugae compositus)	64
syrup of asarum (Syrupus asari compositus)	34
syrup of cimicifuga	64
syrup of figs	195
syrup of hydrochlorophos- phates (Syrupus phosphat- um cum quinina et strych- nina)	179
syrup of hypophosphites	49
syrup of phosphates	11
syrup of sarsaparilla	191
syrup of senna	195
syrup of squill	192
syrup of stillingia	209
syrup of white pine	164
syrup of white pine with mor- phine	135
tar ointment	146
tincture of benzoin	39
tincture of cardamon	56
tincture of cinchona	63
tincture of cudbear	157
tincture of gambir	98
tincture of gentian	100
tincture of guaiac	105
tincture of jalap	120
tincture of kino (Tinctura kino et opii composita)	122
tincture of kino and opium	122
tincture of lavender	142
tincture of pale catechu (Tinctura gambir compos- ita)	98
tincture of viburnum	226
tincture of zedoary (Tinctura zedoariae amara)	228
troches of santalin	190
wine and rhubarb	185
wine of orange	36
Compressed yeast	60
Concentrated and refined diph- theria antitoxin (Serum antidiphthericum purifica- tum)	196
diphtheria antitoxin (Serum antidiphthericum purifica- tum)	196
tetanus antitoxin (Serum antitetanicum purificatum)	196
Condurango	72
fluidextract	72
Confection of rose	185
of senna	194
Confections	72
Conium	72
extract	72
fluidextract	72
maculatum (Conium)	72
Conserves (Confectiones)	72

	PAGE		PAGE
Convallaria flowers	72	Cream—Continued	
flowers, fluidextract	72	of tartar (Potassi bitartras) ..	169
root	72	of tartar, and sulphur troches	
root, fluidextract	73	(Trochisci sulphuris et po-	
Copaiba	73	tassi bitartratis)	215
and opium, mixture	73	Creasote (Creosotum)	75
mass	73	Creosote	75
mixture	73	carbonate	75
solidified (Massa copalbae)...	73	mull, salicylated	12
Copalva (Copaiba)	73	petrox (Petroxolinum creosoti)	75
Copper sulphate	77	petroxolin	75
Coptis	73	water	75
fluidextract	73	Cresol	75
Cordial, blackberry	186	solution, compound	75
Godfrey's (Mistura opii et		Creta praeparata	75
sassafras)	150	Crocus	76
Cordials	73	Croton oil	148
Coriander	74	collodion	148
oil	141	liniment	149
seeds (Coriandrum)	74	liniment, compound	149
Corn silk (Zea)	228	Crude calcium sulphide	50
starch (Amylum)	29	malate of iron (Extractum	
Cornus	74	ferri pomatum)	90
fluidextract	74	malate of iron tincture (Tinc-	
Corrigens elixir (Elixir erio-		tura ferri pomata)	90
dictyli aromaticum)	83	Cubeb	76
syrupus (Syrupus eriodictyli		fluidextract	76
aromaticus)	83	oil	141
Corrosive mercuric chloride	108	oleoresin	76
mercuric chloride mull	109	tincture	76
mercuric chloride tablets, pol-		troches	76
son	109	Cubeba (Cubeba)	76
sublimate (Hydrargyri chlori-		Cudbear	157
dum corrosivum)	108	tincture	157
sublimate tablets (Toxibellae		tincture, compound	157
hydrargyri chloridi corro-		Culver's root (Leptandra)	124
sivi)	109	Cupric sulphate (Copri sulphae)	77
Corydalis	74	Curassao, replacing elixir (Elixir	
elixir, compound	74	aurantii amari)	138
fluidextract	73	Curled dock (Rumex)	187
Cotarnine chloride (Cotarninae		Cusso (Brayera)	43
hydrochloridum)	74	Cypripedium	77
hydrochloride	74	fluidextract	77
Cotton, absorbent (Gossypium		Cystamin (Hexamethylenamina)	107
purificatum)	103	Cystogen (Hexamethylenamina).	107
purified	103		
root bark	103	Dalby's carminative (Mistura	
root bark, fluidextract	103	carminativa)	150
stoptic	86	Damiana	77
Cottonseed oil	141	fluidextract	77
Couch grass (Triticum)	222	Dandelion (Taraxacum)	217
fluidextract (Fluidextractum		fluidextract (Fluidextractum	
tritici)	222	taraxaci)	218
Coumarin	74	Deadly nightshade leaves (Bel-	
Cow's milk	123	ladonnae folia)	38
Crampbark (Viburnum opulus).	226	nightshade root (Belladonnae	
elixir, compound (Elixir vi-		radix)	39
burni opuli compositum)...	226	Decoction of sarsaparilla, com-	
Cranberry bark, high bush (Vi-		pound	191
burnum opulus)	226	Decoctions	77
Cranesbill (Geranium)	101	Decolorized tincture of iodine ..	118
Cream, cold (Unguentum aquae		Deer musk (Moschus)	135
rosae)	186	Dehydrated alcohol	13

PAGE		PAGE	
Delphinium	77	Dinner pills	20
Deodorant solution (Liquor zinci et ferri compositus)	231	Chapman's	20
Deodorized opium	151	Cole's	20
opium tincture	152	Hall's	20
Dermatol (Bismuthi subgallas) ..	42	Lady Webster's (Pululae aloes et mastiches)	20
Dermatologic pastes	155	Dionin (Aethylmorphinae hydro- chloridum)	17
Deshler's salve (Ceratum resinae compositum)	182	Dioscorea	79
Desiccated hypophysis	116	fluidextract	80
pituitary body (Hypophysis sicca)	116	Diphtheria antitoxin (Serum antidiphthericum)	196
suprarenal glands (Supra- renalum siccum)	216	antitoxin, concentrated (Serum antidiphthericum purifica- tum)	196
thyroid glands (Thyroideum siccum)	220	antitoxin, dried (Serum anti- diphthericum siccum)	196
Dewees' carminative (Mistura magnesiae, asafoetidae et opii)	128	antitoxin globulins (Serum antidiphthericum purifica- tum)	196
tincture of gualac (Tinctura gualaci composita)	105	antitoxin, refined and concen- trated (Serum antidiph- thericum purificatum)	196
Dextrin, white	78	Distilled extract of witch hazel (Aqua hamamelidis)	106
Dextrinated paste	78	water	32
Diacetylmorphine	78	water, sterilized	32
and terpin hydrate, elixir	78	Diuretin (Theobrominae sodio- salicylas)	219
chloride (Diacetylmorphinae hydrochloridum)	78	Dobell's solution (Liquor sodii boratis compositus)	200
hydrochloride	78	Dog grass (Triticum)	222
Diachylon ointment	166	Dogwood bark (Cornus)	74
plaster (Emplastrum plumbi) ..	166	Donovan's solution (Liquor ar- seni et hydrargyri iodidi) ..	33
Diarrhea mixture, Squibb's (Mistura opii et chloroform composita)	150	Dover's powder (Pulvis ipe- cacuanhae et opii)	151
Diastase	78	syrup (Syrupus Ipecacuanhae et opii)	151
Digitalis	79	tincture (Tinctura Ipecacu- anhae et opii)	152
fluidextract	79	Dried alum (Alumen exsicca- tum)	23
infusion	79	antidiphtheric serum	196
leaves (Digitalis)	79	antitetanic serum	197
quinine and opium pills	151	diphtheria antitoxin (Serum antidiphthericum siccum) ..	196
squill and mercury pills	79	ferrous sulphate (Ferri sul- phas exsiccatus)	93
tincture	79	suprarenals	216
Diluted acetic acid	6	tetanus antitoxin (Serum anti- tetanicum siccum)	197
alcohol	18	thyroids	220
hydriodic acid	9	Drop chalk (Creta Praeparata) ..	75
hydrobromic acid	9	Drosera	80
hydrochloric acid	3	fluidextract	80
hydrocyanic acid	9	Drugs, fresh tinctures	221
hypophosphorous acid	10	Dulcamara	80
iodine petroxolin	118	Dwarf pine needles, oil of	146
mercurial ointment	112	pine oil (Oleum pini pumili- onis)	146
nitrohydrochloric acid	10		
nitromuriatic acid (Acidum nitrohydrochloricum dilu- tum)	10		
phosphoric acid	11		
prussic acid (Acidum hydro- cyanicum dilutum)	9		
solution of lead subacetate ..	167		
sulphuric acid	13		
Dimethyl-ketone (Acetonum) ..	6		
Dimethylxanthine (Theophyl- lina)	219		

	PAGE		PAGE
Eau sédative de raspail (Loto ammoniacalis comphorata)	25	Elixir—Continued	
Echinacea	20	of almond, compound	137
fluidextract	80	of ammonium bromide	26
Effervescent artificial Carlsbad salt	207	of ammonium valerate	28
artificial Kissingen salt	202	of anise	29
artificial Vichy salt	200	of bismuth	41
artificial Vichy salt with lithium	200	of bitter orange	138
citrate caffeine	45	of black haw (Elixir viburni prunifolii)	227
potassium bromide with caffeine (Sal potassii bromidi effervescens compositus)	170	of blackberry, compound	186
potassium citrate	171	of buchu	44
salt of lithium citrate	126	of buchu and potassium acetate	44
salt of potassium bromide	170	of buchu compound	44
salt of potassium bromide, compound	170	of calcium and sodium glycerophosphates	47
salts, granular	188	of calcium bromide	46
sodium phosphate	206	of calcium hypophosphite	48
solution of magnesium sulphate	129	of calcium lactophosphate	49
solution of sodium citro-tartrate	202	of callisaya, alkaloidal ferrated (Elixir cinchonae alkaloidorum et ferri)	92
Effervescing powder, compound	172	of callisaya, alkaloidal, with hypophosphites (Elixir cinchonae alkaloidorum et hypophosphitum)	180
Egg albumen, fresh	152	of callisaya, alkaloidal, with iron and bismuth (Elixir cinchonae alkaloidorum, ferri et bismuthi)	41
and cod liver oil emulsion	145	of callisaya, alkaloidal, with iron and pepsin (Elixir cinchonae alkaloidorum, ferri et pepsini)	156
fresh	153	of callisaya, alkaloidal, with iron and strychnine (Elixir cinchonae alkaloidorum, ferri et strychninae)	213
yolk, fresh	152	of callisaya, alkaloidal, with iron, bismuth and strychnine (Elixir cinchonae alkaloidorum, ferri, bismuthi et strychninae)	213
yolk, glycerite	153	of callisaya, iron and lactophosphate of lime (Elixir cinchonae alkaloidorum, ferri et calcii lactophosphatis)	50
Elsenzucker (Ferri oxidum saccharatum)	91	of cardamon, compound	140
Elaeasacchara (Oleosacchara)	137	of cascara sagrada	57
Elaterin	80	of cascara sagrada compound	57
trituration	81	of cinchona alkaloids	180
Elder flowers (Sambucus)	188	of cinchona alkaloids and hypophosphites	180
Elecampane (Inula)	117	of cinchona alkaloids and iron	92
Electuaries (Confectiones)	72	of cinchona alkaloids, iron and bismuth	41
Elixir adjuvans (Elixir glycyrrhizae)	19	of cinchona alkaloids, iron and calcium lactophosphate	49
amygdalae compositum	137	of cinchona alkaloids, iron and pepsin	156
aromaticum	18	of cinchona alkaloids, iron and strychnine	213
aurantii amari	138		
aurantiorum compositum (Vinum aurantii compositum)	36		
callisaya, alkaloidal (Elixir cinchonae alkaloidorum)	180		
cathartic, compound	97		
corrigena (Elixir eriodictyi aromaticum)	83		
corydalis compositum	74		
curassao, replacing (Elixir aurantii amari)	138		
ferri pyrophosphatis	93		
ferri pyrophosphatis, quinae et strychninae	93		
glycerophosphatum (Elixir calcei et sodii glycerophosphatum)	47		
humuli	108		
laxative (Elixir cascarae sagradae compositum)	57		

	PAGE
Elixir—Continued	
of cinchona alkaloids, iron, bismuth and strychnine.....	213
of corydalis, compound.....	74
of crampbark, compound (Elixir viburni opuli compositum)	226
of eriodictyon, aromatic.....	83
of ferric hypophosphite.....	89
of ferric phosphate.....	92
of ferric pyrophosphate.....	93
of ferric pyrophosphate, quinine and strychnine.....	93
of formates.....	8
of formates compound.....	8
of gentian.....	100
of gentian and ferric phosphate.....	92
of gentian and iron.....	87
of gentian, glycerinated.....	100
of gentian with tincture of ferric citro-chloride (Elixir gentinae et ferri).....	87
of glycerophosphates compound.....	203
of glycyrrhiza.....	19
of glycyrrhiza, aqueous.....	102
of glycyrrhiza, aromatic.....	19
of guarana.....	106
of hops.....	108
of hypophosphites.....	48
of hypophosphites and iron.....	89
of iron lactate.....	90
of iron, quinine and strychnine.....	87
of licorice (Elixir glycyrrhizae).....	19
of licorice, aqueous (Elixir glycyrrhizae aquosum).....	102
of licorice, aromatic (Elixir glycyrrhizae aromaticum).....	19
of lithium bromide.....	125
of lithium citrate.....	126
of lithium salicylate.....	126
of orange, compound (Vinum aurantii compositum).....	36
of pepsin.....	156
of pepsin and bismuth.....	156
of pepsin and iron.....	156
of pepsin and rennin compound.....	156
of pepsin, bismuth and strychnine.....	212
of phosphorus.....	161
of phosphorus and nuxvomica.....	161
of potassium acetate.....	168
of potassium acetate and juniper.....	168
of potassium bromide.....	170
of pyrophosphate of iron (Elixir ferri pyrophosphatis).....	93

	PAGE
Elixir—Continued	
of quinine valerate and strychnine.....	181
of sodium bromide.....	201
of sodium hypophosphite.....	204
of sodium salicylate.....	206
of sodium salicylate compound.....	206
of strychnine valerate.....	214
of taraxacum compound.....	217
of terpin hydrate.....	218
of terpin hydrate and codeine.....	68
of terpin hydrate and diacetylmorphine.....	78
of three bromides.....	26
of vanillin compound.....	224
of viburnum opulus compound.....	226
of viburnum prunifolium.....	227
of yerba santa, aromatic (Elixir eriodictyli aromaticum).....	83
of zinc valerate.....	231
pepsini.....	156
pepsini, bismuth et strychninae.....	212
pepsini et bismuthi.....	156
pepsini et ferri.....	156
pepsini et rennini compositum.....	156
phosphori.....	161
phosphori et nuclei vomicae.....	161
potassii acetatis.....	168
potassii acetatis et juniperi.....	168
potassii bromidi.....	170
quininae valeratis et strychninae.....	181
quininae valerianatis et strychninae.....	181
red (Elixir aromaticum rubrum).....	19
red aromatic.....	19
ramni purshianae (Elixir cascarae sagradae).....	57
ramni purshianae co. (Elixir cascarae sagradae compositum).....	57
rubi compositum.....	186
simple (Elixir aromaticum).....	18
sodii bromidi.....	201
sodii hypophosphitis.....	204
sodii salicylatis.....	206
sodii salicylatis compositum.....	206
strychninae valeratis.....	214
strychninae valerianatis (Elixir strychninae valeratis).....	214
taraxaci compositum.....	217
terpini hydratis.....	218
terpini hydratis et codeinae.....	68
terpini hydratis cum heroina (Elixir terpin hydratis et diacetylmorphinae).....	78
terpini hydratis et diacetylmorphinae.....	78
trium bromidorum.....	20

	PAGE		PAGE
Elixir—Continued		Emulum—Continued	
vanillin compositum	224	olei terebinthinae	148
viburni opuli compositum	226	Epsom salt (Magnesii sulphas)	129
viburni prunifolii	227	Ergot	82
zinci valeratis	231	extract	82
zinci valerianatis (Elixir zinci valeratis)	231	extract, aqueous	82
Elixirs	81	fluidextract	82
Elm	223	of rye (Ergota)	82
bark (Ulmus)	223	tincture, ammoniated	82
troches	223	Eriodictyon	83
Emetine hydrochloride	81	elixir, aromatic	83
Emollient cataplasm (Species emollientes)	23	fluidextract	83
species	23	syrup, aromatic	83
Emplastra	82	Eserine salicylate (Physostigminae salicylas)	162
Emplastrum		Essence of peppermint (Spiritus menthae piperitae)	143
See also under Plaster.		Essentia pepsini (Elixir pepsini et rennini compositum)	156
belladonnae	38	Ether	16
cantharidis	54	acetic	17
capsici	55	nitrous	17
elasticum	182	nitrous, spirit	17
fusculum camphoratum	53	petroleum (Benzinum purificatum)	39
plumbi	166	spirit	16
resinae	182	spirit, compound	16
saponis	190	Ethereal oil	137
sinapis	197	tincture of ferric chloride	87
Empyroligneous oil of birch, rectified (Oleum betulae empyreumaticum rectificatum)	139	tinctures	221
Emulsion of almond	29	Ethyl acetate (Aether aceticus)	17
of asafetida	34	carbamate	17
of castor oil	146	chloride	17
of cod liver oil	144	ester, carbamic acid (Aethylis carbamas)	17
of cod liver oil with calcium lactophosphate	144	nitrate (Aether nitrosus)	17
of cod liver oil with calcium phosphate	144	urethane (Aethylis carbamas)	17
of cod liver oil with egg	145	Ethylmorphine chloride (Aethylmorphinae hydrochloridum)	17
of cod liver oil with hypophosphites	144	hydrochloride	17
of cod liver oil with malt	144	Eucaine (Betaeucainae hydrochloridum)	40
of cod liver oil with phosphate of lime (Emulum olei morrhuae cum calcei phosphate)	144	chloride (Betaeucainae hydrochloridum)	40
of cod liver oil with wild cherry	144	Eucalyptol	83
of oil of turpentine	148	petrox (Petroxolinum eucalyptolis)	83
of petrolatum	158	petroxolin	83
Emulsions	82	spray	83
Emulum amygdalae	29	Eucalyptus	83
olei morrhuae	144	fluidextract	84
olei morrhuae cum calcei lactophosphate	144	leaves (Eucalyptus)	83
olei morrhuae cum calcei phosphate	144	oil	141
olei morrhuae cum hypophosphitibus	144	Eugenol	84
olei morrhuae cum malto	144	Euonymus	84
olei morrhuae cum pruno virginian	144	extract	84
olei morrhuae cum vitello	145	extract, powdered (Extractum euonymi)	84
olei ricini	146	fluidextract	84
		Eupatorium	84
		fluidextract	84
		Euphorbia pluvifera	84
		pluvifera, fluidextract	84

INDEX

249

	PAGE
European goat's rue (Galega) ..	98
Expectorant, Stokes'	26
Expressed oil of almond	138
Exsiccated alum	23
ferrous sulphate	93
sodium arsenate	198
sodium phosphate	205
sodium sulphite	207
Extract, Goulard's (Liquor plumbi subacetatis)	167
of aconite	15
of aconite, powdered (Extrac- tum aconiti)	15
of aloes	20
of aloes, powdered (Extractum aloes)	20
of apples, ferrated	90
of apples, ferrated, tincture ..	90
of beef	56
of belladonna leaves	38
of cannabis	54
of cascara sagrada	57
of cascara sagrada, powdered (Extractum cascarae sa- grada)	57
of cicicifuga	63
of cicicifuga, powdered (Ex- tractum cicicifugae)	63
of cinchona	64
of colchicum corm	69
of colchicum corm, powdered (Extractum colchici cormi) ..	69
of colocynth	70
of colocynth, compound	70
of colocynth, powdered (Ex- tractum colocynthidis)	70
of colocynth compound, pow- dered (Extractum colocyn- thidis compositum)	70
of conium	72
of ergot	82
of ergot, aqueous	82
of euonymus	84
of euonymus, powdered (Ex- tractum euonymi)	84
of gelsemium	99
of gelsemium, powdered (Ex- tractum gelsemii)	99
of gentian	100
of glycyrrhiza	102
of glycyrrhiza, pure	102
of golden seal (Extractum hy- drastis)	114
of hematoxylon	106
of hydrastis	114
of hydrastis, powdered (Ex- tractum hydrastis)	114
of hyoscyamus	115
of ignatia	116
of ignatia, powdered (Extrac- tum ignatiae)	116
of jalap	120

Extract—Continued

	PAGE
of krameria	122
of krameria, powdered (Ex- tractum krameriae)	122
of leptandra	124
of leptandra, powdered (Ex- tractum leptandrae)	124
of licorice (Extractum gly- cyrrhizae)	102
of malt	129
of nux vomica	136
of nux vomica, powdered (Ex- tractum nucis vomicae) ..	136
of opium	149
of opium, powdered (Extrac- tum opii)	149
of oxgall	85
of oxgall, powdered (Extrac- tum felis bovis)	85
of physostigma	162
of physostigma, powdered (Ex- tractum physostigmatis) ..	162
of podophyllum	167
of quassia	176
of quassia, powdered (Extrac- tum quassiae)	176
of rhubarb	183
of rhubarb, powdered (Ex- tractum rhei)	183
of stramonium	210
of sumbul	216
of taraxacum	218
of viburnum prunifolium	227
of viburnum prunifolium, pow- dered (Extractum viburni prunifolii)	227
of witch hazel, distilled (Aqua hamamelidis)	106
Extracts	85
solid	85
Extractum	
See also under Extract.	
carnis	56
euonymi	84
felis bovis	85
ferri pomatum	90
rhamni purshianae (Extrac- tum cascarae sagradae) ..	57
rhei	183
False unicorn (Helionias)	107
Farfara	85
Fel bovis	85
Fennel	96
oil	141
seed	96
water	141
Fermented milk	123
Ferrated elixir of calisaya, alka- loidal (Elixir cinchonae alkaloidorum et ferri)	92
extract of apples	80
wine of wild cherry	174

	PAGE		PAGE
Ferri carbonas	85	Ferric—Continued	
carbonas saccharatus	86	phosphate	91
cinchonae et calcei lactophos-		phosphate elixir	92
phatis, elixir (Elixir cin-		phosphate, soluble (Ferri phos-	
chonae alkaloidorum, ferri		phas)	91
et calcei lactophosphatis) ..	50	pyrophosphate	92
et ammonii citras	87	pyrophosphate elixir	93
et quinae citras	88	pyrophosphate, quinine and	
et quinae citras solubilis		strychnine elixir	93
(Ferri et quinae citras) ..	88	pyrophosphate, soluble (Ferri	
iodidi syrupus (Syrupus ferri		pyrophosphas)	92
iodidi)	90	salicylate solution	206
iodidum	90	subsulphate	93
lactas	90	subsulphate solution	93
lactophosphas	90	sulphate, basic solution (Li-	
malas	90	quor ferri subsulphatis) ..	93
malas crudus (Extractum ferri		sulphate solution	94
pomatum)	90	Ferrous carbonate	85
oxidum saccharatum	91	carbonate mass	85
protochloridum	92	carbonate pills	86
tersulphas	94	chloride	92
Ferric acetate	85	chloride solution	92
acetate solution	85	chloride syrup	92
and gentian phosphate elixir.	92	iodide	90
chloride	86	iodide pills	90
chloride solution	86	iodide syrup	90
chloride, tincture, ethereal ..	87	lactate	90
chloride, tasteless tincture		malate	90
(Tinctura ferri citro-chlor-		sulphate	93
idi)	87	sulphate, dried (Ferri sulphas	
chloride tincture	86	exsiccatus)	93
citrate	87	sulphate, exsiccated	93
citrate, soluble (Ferri et am-		sulphate, granulated	94
monii citras)	87	sulphate, precipitated (Ferri	
citrate solution	87	sulphas granulatus)	94
citro-chloride tincture	87	Ferruginous pills (Pilulae ferri	
citro-chloride tincture with		carbonatis)	86
gentian elixir (Elixir gen-		Ferrum	94
tianae et ferri)	87	albuminatum	94
glycerinophosphate (Ferri gly-		oxydatum saccharatum (Ferri	
cerophosphas)	88	oxidum saccharatum)	91
glycerophosphate	88	peptonatum	94
hydrate with magnesia (Ferri		reductum (Ferrum reductum) ..	95
hydroxidum cum magnesi		Ficus	96
oxido)	88	Fig	96
hydroxide	88	Flgs, syrup compound	195
hydroxide (Magma ferri hy-		Firm zinc glyceragelatin	229
droxidi)	89	Fish berry (Coccus indicus) ..	67
hydroxide magma	89	Flaxseed (Linum)	124
hydroxide with magnesium		oil (Oleum lini)	143
oxide	88	Fleming's solution (Liquor cal-	
hypophosphite	89	cis sulphuratae)	51
hypophosphite elixir	89	Flexible collodion	176
hypophosphite solution	89	Flies, Russian (Cantharis)	54
hypophosphite syrup	89	Spanish (Cantharis)	54
nitrate	91	Flowers of sulphur (Sulphur	
nitrate solution	91	sublimatum)	215
oxide, soluble (Ferri oxidum		Fluidextract, aromatic	66
saccharatum)	91	of aconite	15
oxychloride	91	of adonis	16
oxychloride solution	91	of aletris	19
oxysulphate	91	of angelica root	30
oxysulphate solution	91	of apocynum	31

INDEX

251

	PAGE
Fluidextract—Continued	
of aralia	32
of arnica	33
of asclepias	35
of aspidosperma	35
of baptisia	38
of belladonna root	39
of berberis	40
of bitter orange peel	36
of bittersweet	80
of black cohosh (Fluidextractum cimicifuga)	63
of black haw (Fluidextractum viburni prunifolii)	227
of black snakeroot (Fluidextractum cimicifuga)	63
of blue flag	120
of boldo	43
of buchu	44
of buchu compound	44
of buckthorn bark (Fluidextractum frangulae)	97
of calendula	51
of calumba	51
of Canadian hemp (Fluidextractum apocyni)	31
of cannabis	54
of cascara sagrada	58
of cascara sagrada, aromatic	58
of catnep	59
of caulophyllum	59
of celery fruit	31
of chestnut leaves	58
of chimaphila	60
of chiononthus	60
of chirata	61
of cimicifuga	63
of cinchona	64
of cinchona, aqueous	65
of cocillana	68
of coffee	69
of colchicum corm	69
of colchicum seed	69
of condurango	72
of conium	72
of convallaria flowers	72
of convallaria root	73
of coptis	73
of cornus	74
of corydalis	74
of cotton root bark	103
of couch grass (Fluidextractum tritici)	222
of cubeb	76
of cyripedium	77
of damiana	77
of dandelion (Fluidextractum taraxaci)	218
of digitalis	79
of dioscorea	80
of drosera	80
of echinacea	80
of ergot	82

	PAGE
Fluidextract—Continued	
of eriodictyon	83
of eucalyptus	84
of euonymus	84
of eupatorium	84
of euphorbia pilulifera	84
of frangula	97
of fucus	97
of galega	98
of gelsemium	99
of gentian	100
of geranium	101
of ginger	231
of glycyrrhiza	102
of green hellebore (Fluidextractum veratri viridis)	225
of grindelia	104
of guarana	106
of hamamelis leaves	106
of hellanthemum	107
of helonias	107
of henbane (Fluidextractum hyoscyami)	115
of hops	108
of hydrangea	108
of hydrastis	114
of hyoscyamus	115
of ipecac	119
of jaborandi (Fluidextractum pilocarpi)	163
of jalap	120
of juglans	121
of juniper berries	121
of kava	121
of kola	122
of krameria	123
of lappa	124
of leptandra	124
of lobelia	126
of lupulin	127
of matico	131
of mezereum	134
of mullein leaves	225
of musk root (Fluidextractum sumbul)	216
of nux vomica	136
of paracoto	154
of pareira	155
of parsley root	159
of phytolacca	162
of pilocarpus	163
of pinkroot (Fluidextractum spigelliae)	208
of podophyllum	167
of pomegranate	103
of prickly ash (Fluidextractum xanthoxyl)	228
of quassia	176
of quebracho (Fluidextractum aspidospermatis)	25
of quercus	176
of rhamnus cathartica	183
of rhubarb	183

	PAGE		PAGE
Fluidextract—Continued		Fluidextractum—Continued	
of rhus glabra	185	turnerae (Fluidextractum da-	77
of rose	186	mlanae)	225
of rubus	187	verbasci foliae	231
of rumex	187	zingiberis	58
of sabal	187	Fluidglycerate of cascara sa-	58
of sanguinaria	189	grada	102
of sarsaparilla	191	of cascara sagrada, aromatic .	123
of sarsaparilla compound . . .	191	of glycyrrhiza	102
of saw palmetto (Fluidextrac-	187	of krameria	102
tum sabal)	193	of licorice (Fluidglyceratum	183
of scoparius	193	glycyrrhizae)	96
of scutellaria	194	of rhubarb	96
of senecio	194	Fluidglycerates	96
of senna	194	Foeniculum	96
of serpentaria	195	Formaldehyde	96
of solanum	208	solution	96
of spigella	208	Formamin (Hexamethylenamina)	107
of squill	192	Formates elixir	8
of staphisagria	209	elixir, compound	8
of stavesacre (Fluidextractum	209	Formic acid	8
staphisagria)	209	acid spirit	8
of stillingia	209	Formin (Hexamethylenamina) .	107
of stillingia compound	209	Fowler's solution (Liquor potas-	168
of stramonium	210	sil arsenitis)	79
of sumbul	216	Foxglove (Digitalis)	21
of taraxacum	218	Francis' triplex pills (Pillulae	97
of thuja	220	aloes, hydrargyri et scam-	97
of thyme	220	monii compositae)	118
of trifolium	222	Frangula	214
of trillium	222	fluidextract	221
of triticum	222	Fraxinus	153
of uva ursi	223	French mixture (Liquor iodi	152
of valerian	224	phenolatus)	152
of veratrum viride	225	Fresh apple juice	107
of verberna	226	drugs, tinctures	97
of viburnum opulus	226	egg	8
of viburnum prunifolium . . .	227	egg albumen	13
of wild cherry	174	egg yolk	60
of xanthoxylum	228	Fringe tree bark (Chionanthus)	107
of zea	228	Frost-weed (Helianthemum) .	97
Fluidextracts	96	Fucus	97
Fluidextractum		fluidextract	98
See also under Fluidextract.		Galangal	98
apil fructi	31	Galega	98
aurantii amari	36	fluidextract	98
castaneae	58	Galla	9
catariae	59	Gallie acid	13
corni	74	Gallotannic acid (Acidum tan-	98
coto (Fluidextractum para-	154	nicum)	98
coto)	80	Gambir	98
dulcamarae	103	powder, compound	98
gossypii corticis	103	tincture, compound	99
granati	108	troches	52
humuli	120	Gamboge	52
iridis versicoloris	159	pipe (Cambogia)	105
petroselinii radicis	174	Gargle of gualac compound .	19
uni virginianae	183	Garlic	19
labrae	185	syrup	133
.....	192	Gaultheriae, oleum (Methylis	
		alicylas)	

INDEX

253

	PAGE
Gelatin	99
glycerinated	99
Gelatinum chondri	63
Gelsemium	99
extract	99
extract, powdered (Extractum gelsemii)	99
fluidextract	99
tincture	99
Gentian	99
and ferric phosphate elixir	92
and iron elixir	87
and rhubarb tincture	185
elixir	100
elixir, glycerinated	100
extract	100
fluidextract	100
infusion, compound	100
tincture, compound	100
with tincture of ferric chloro-chloride, elixir of (Elixir gentianae et ferri)	87
Geranium	101
fluidextract	101
German chamomile (Matricaria)	131
Gilead buds, balm of (Populi gemmae)	167
Ginger	231
fluidextract	231
oleoresin	232
syrup	232
tincture (Tinctura zingiberis)	232
wild (Asarum)	34
Glacial acetic acid	6
Glandulae suprarenales siccae (Suprarenalium siccum)	216
Glandulae thyroideae siccae (Thyroideum siccum)	220
Glauber's salt (Sodii sulphas)	206
Glonoin (Glycerylis nitras)	101
pills (Pillulae glycerylis nitratris)	102
spirit (Spiritus glycerylis nitratris)	102
Glucose	101
liquid (Glucosum)	101
syrup (Glucosum)	101
Glusidum (Benzosulphinidum)	40
Glycerin	101
suppositories	101
Glycerinated elixir of gentian	100
gelatin	99
vaccine virus (Virus vaccini-cum)	227
Glycerite of blsmuth	41
of boroglycerin	7
of carbolic acid (Glyceritum phenolis)	159
of egg yolk	153
of golden seal (Glyceritum hydrastis)	114
of gualae	105
of hydrastis	114

	PAGE
Glycerite—Continued	
of pepsin	157
of phenol	159
of starch	29
of tannic acid	13
of tannin (Glyceritum acidi tannici)	13
of tar	165
of tragacanth	221
Glycerites	101
Glyceritum amyli	29
pilis liquidae	165
vitelli	153
Glycerogelatinum acidi salicylici	12
iodoformi	117
zinci durum	229
zinci molle	229
Glycerol (Glycerinum)	101
Glycerophosphates elixir, compound	203
solution, compound (Elixir glycerophosphatum compositum)	203
Glyceryl trinitrate	101
trinitrate spirit	102
Glycerylis nitras	101
Glyconin (Glyceritum vitelli)	153
Glycyrrhiza	102
elixir	19
elixir, aqueous	102
elixir, aromatic	19
extract	102
extract, pure	102
fluidextract	102
fluidglycerate	102
mixture, compound	150
powder, compound	194
syrup	103
Glycyrrhizin, ammoniated	103
Goat's rue, European (Galega)	98
Godfrey's cordial (Mistura opii et sassafras)	150
Gold and arsenic bromide solution	7
and sodium chloride	37
Golden seal (Hydrastis)	114
extract (Extractum hydrastis)	114
glycerite (Glyceritum hydrastis)	114
tincture (Tinctura hydrastis)	114
Goldthread (Coptis)	73
Gossypii cortex	103
Gossypium purificatum	103
stypticum	86
Goulard's cerate (Ceratum plumbi subacetatis)	167
extract (Liquor plumbi subacetatis)	167
Granatum	103
Granular effervescent salts	188
ferrous sulphate	94
opium	151

	PAGE		PAGE
Gray powder (Hydrargyrum cum creta)	112	Hebra's ich ointment (Unguentum sulphuris compositum)	216
Green hellebore (Veratrum viride)	225	Hellianthemum fluidextract	107
hellebore, fluidextract (Fluidextractum veratri viridis)	225	Hellebore, American (Veratrum viride)	225
soap, tincture (Linimentum saponis molles)	190	green (Veratrum viride)	225
Gregory's powder (Pulvis rhei compositus)	184	Helonias	107
Griffith's mixture (Mistura ferri composita)	86	fluidextract	107
Grindelia	104	Hematoxylon extract	106
fluidextract	104	Hemlock, poison (Conium)	72
Gualac	104	Henbane (Hyoscyamus)	115
gargle, compound	105	fluidextract (Fluidextractum hyoscyami)	115
glycerite	105	tincture (Tinctura hyoscyami)	115
mixture	105	Heroin (Diacetylmorphina)	78
resin (Gualacum)	104	hydrochloride (Diacetylmorphinae hydrochloridum)	78
tincture	105	Hexamethylenamine	107
tincture, ammoniated	105	(Hexamethylenamina)	107
tincture, compound	105	Hexamethylene-Tetramine (Hexamethylenamina)	107
tincture, Dewees' (Tinctura gualaci composita)	105	Hiera picra (Pulvis aloes et Canellae)	21
wood	104	High bush cranberry bark (Viburnum opulus)	226
lignum	104	Hive syrup (Syrupus scillae compositus)	192
Gualacal	104	Hoffmann's anodyne (Spiritus aetheris compositus)	16
carbonate	104	drops (Spiritus aetheris compositus)	16
petrox (Petroxolinum gualacolis)	105	Homatropine bromide (Homatropinae hydrobromidum)	107
petroxolin	105	hydrobromide	107
Gualacum	104	Honey	132
Guarana	105	and borax (Mel sodii boratis)	200
elixir	106	clarified	132
fluidextract	106	of rose	132
Gum arabic (Acacia)	5	of rose and sodium borate	200
asafetida (Asafoetida)	34	of rose with borax (Mel rosae et sodii boratis)	200
benjamin (Benzoinum)	39	of sodium borate	200
myrrh (Myrrha)	136	Honeys	132
senegal (Acacia)	5	Hope's mixture (Mistura camphorae acida)	149
tragacanth (Tragacantha)	221	Hops	108
Gun cotton, soluble (Pyroxylinum)	175	elixir	108
Gutta percha	106	fluidextract	108
solution	106	tincture	108
Guttae pectorales (Tinctura pectoralis)	152	Horsenettle berries (Solanum)	207
Guy's pills (Pillulae digitalis, scillae et hydrargyri)	79	Hot drops (Tinctura capsici et myrrhae)	55
Haematoxylon	106	Humulus	108
Hall's dinner pills	20	Hydrangia	108
solution of strychnine (Liquor strychninae acetatis)	212	fluidextract	108
Hamamelis	106	Hydrargyri chloridum corrosivum	108
leaves	106	chloridum mite	109
leaves, fluidextract	106	iodidum flavum	110
water	106	iodidum rubrum	110
Warshorn liniment (Linimentum ammoniacae)	25	nitras	110
magnesia (Magnesii oxiponderosum)	129	oxidum flavum	110
um oxide	129		

PAGE	PAGE
Hydrargyri—Continued	Hypophosphites—Continued
oxidum rubrum 111	and cinchona alkaloids, elixir. 180
salicylas 111	and iron elixir 89
unguentum (Unguentum hy-	elixir 48
drargyri dilutum) 112	solution 48
Hydrargyrum 112	solution, compound 48
ammoniatum 113	syrup 48
cum creta 112	syrup, compound 49
Hydrastine 113	with cod liver oil emulsion . . . 144
chloride (Hydrastinae hydro-	Hypophosphorous acid 10
chloridum) 113	acid, diluted 10
hydrochloride 113	Hypophysis, desiccated 116
solution, colorless (Liquor hy-	solution 116
drastinae compositus) 113	
solution, compound 113	Ignatia 116
Hydrastis 114	amara (Ignatia) 116
extract 114	extract 116
extract, powdered (Extractum	extract, powdered (Extractum
hydrastis) 114	ignatiae) 116
fluidextract 114	tincture 116
glycerite 114	India senna or Alexandria senna
tincture 114	(Senna) 194
Hydrated chloral 61	Indian berry (Cocculus indicus)
Hydriodic acid, diluted 9	tobacco (Lobelia) 126
acid, syrup 9	Indigo carmine (Sodii Indigotin-
Hydrobromic acid, diluted 9	disulphonas) 204
Hydrochloric acid 9	Infusa 116
acid, syrup 9	Infused oils 137
solution of arsenic (Liquor	Infusion of brayera 43
acidi arsenosi) 34	of cinchona 65
hydrochlorophosphates syrup,	of digitalis 79
compound (Syrupus phos-	of gentian compound 100
phatum cum quina et	of rose compound 186
strychnina) 179	of senna compound 194
Hydrocyanic acid, diluted 9	of wild cherry 174
Hydrogen dioxide 114	Infusions 116
dioxide solution 115	Infusum pruni virginianae 174
peroxide (Hydrogenii diox-	Inula 117
idum) 114	Inunction, menthol 132
peroxide solution (Liquor hy-	menthol, compound 133
drogenii dioxidi) 115	Inunctions 117
Hydrous wool fat (Adeps lanae	Iodi carbolatus, liquor (Liquor
hydrosus) 16	iodi phenolatus) 118
Hyoscine hydrobromide (Scopo-	Iodine 117
laminae hydrobromidum) . . . 193	collodion 118
Hyoscyami collum (Hyoscyamus)	ointment 119
tinctura (Tinctura hyoscyami)	petrox 5% (Petroxolinum iodi
tinctura) 115	dilutum) 118
Hyoscyamine bromide (Hyoscy-	petrox 10% (Petroxolinum
aminae hydrobromidum) 115	iodi) 118
hydrobromide 115	petroxolin 118
Hyoscyamus 115	petroxolin, diluted 118
and colocynth pills 71	solution, carbolyzed (Liquor
extract 115	iodi phenolatus) 118
fluidextract 115	solution, compound 118
oil, compound 115	solution, phenolated 118
tincture 115	tincture 118
Hypophosphite of iron, solution	tincture, Churchill's (Tinctura
of (Liquor ferri hypophos-	iodi fortior) 119
phitis) 89	tincture, decolorized 118
Hypophosphites and calisaya,	tincture, stronger 118
alkaloidal, elixir (Elixir cin-	Iodized phenol 189
chonae alkaloidorum et	
hypophosphitum) 180	

	PAGE		PAGE
Iodoform	117	Iron—Continued	
aromatized	117	bismuth, strychnine and cin-	
colloidion	117	chona alkaloids elixir.....	213
glycerogelatin	117	bitter wine	88
ointment	117	by hydrogen (Ferrum reduc-	
petrox (Petroxolinum iodo-		tum)	95
formi)	117	calisaya alkaloidal and	
petroxolin	117	strychnine elixir (Elixir	
Iodotannin syrup	118	cinchona alkaloidorum,	
Iodum	117	ferri et strychninae).....	213
Ipecac	119	calisaya and lactophosphate	
and opium powder	151	of lime elixir (Elixir cin-	
and opium syrup	151	chona alkaloidorum, ferri	
and opium tincture	152	et calcii lactophosphatis)...	50
fluidextract	119	cinchona alkaloids and calcium	
powder, compound (Pulvis ipe-		lactophosphate elixir	49
cacuanhae et opii)	151	cinchona alkaloids and strychnine	
syrup	119	elixir	213
wine	119	citrate, wine of (Vinum	
Ipecacuanhae radix	119	ferri)	88
Iris	119	citro-chloride	87
versicolor	120	crude malate (Extractum ferri	
Irish moss (Chondrus)	62	pomatum)	90
moss gelatin (Gelatinum chon-		elixir of pyrophosphate of	
dri)	63	iron (Elixir ferri pyrophos-	
moss mucilage (Mucilago chon-		phatis)	93
dri)	63	hypophosphite, solution (Li-	
Iron	94	quor ferri hypophosphitis)...	89
albuminate	94	lactate (Ferri lactas)	90
albuminate, solution	94	lactate elixir	90
and aloe pills	20	lactophosphate	90
and ammonium acetate, solu-		lactophosphate syrup	90
tion	25	malate (Ferri malas).....	90
and ammonium citrate	87	mixture compound	86
and beef wine	57	pepsin and calisaya, alka-	
and calcium lactophosphate,		loidal, elixir (Elixir cin-	
syrup	50	chona alkaloidorum ferri	
and cinchona alkaloids elixir,	92	et pepsini)	156
and gentian elixir	87	pepsin and cinchona alka-	
and hypophosphites elixir...	89	loids elixir	156
and manganese iodide syrup..	90	peptonate	94
and manganese, peptonate,		perchloride (Ferri chloridum)..	86
solution	95	perchloride solution (Liquor	
and pepsin elixir	156	ferri chloride)	86
and protochloride solution		protochloride (Ferri proto-	
(Liquor ferri protochloridi)	92	chloridum)	92
and quinine citrate	88	protochloride, syrup (Syrupus	
and quinine citrate, soluble		ferri protochloridi)	92
(Ferri et quinae citras)...	88	protosulphate (Ferri sulphas)	
and zinc solution, compound.	231	Quevenne's (Ferrum reduc-	
beef and wine (Vinum carnis		tum)	95
et ferri)	57	quinine, aloe, and nux vomica	
bismuth and calisaya, alka-		pills	94
loidal, elixir (Elixir cin-		quinine and strychnine elixir.	
chona alkaloidorum, ferri		quinine and strychnine, syrup	
et bismuthi)	41	of the phosphates of.....	92
bismuth and cinchona alka-		quinine, strychnine and ar-	
loids elixir	41	senic pills, mild.....	95
bismuth, calisaya, alkaloidal,		quinine, strychnine and ar-	
and strychnine, elixir (Elixir		senic pills, stronger.....	95
cinchona alkaloidorum,		reduced	95
ferri, bismuthi et strychni-			
nae)	213		

PAGE	PAGE
Iron—Continued	
saccharated oxide of, syrup (Syrupus ferri saccharati solubilis)	91
salicylated mixture (Liquor ferri salicylatis)	206
sesquichloride (Ferri chlori- dum)	86
soluble oxide of, syrup (Syrupus ferri saccharati solubilis)	91
solution of peptonate of,	94
tersulphate (Ferri tersulphas) tersulphate solution (Liquor ferri tersulphatis)	94
tincture of crude malate, (Tinctura ferri pomata)....	90
tincture, tasteless (Tinctura ferri citro-chloridi)	87
wine	88
Jaborandi (Pilocarpus)	163
fluidextract (Fluidextractum pilocarpi)	163
Jackson's pectoral syrup (Syrupus morphinae et acaciae)	134
Jalap	120
and calomel (Pulvis hydrar- gyri chloridi mitis et Ja- lapae)	110
and mild mercurous chloride powder	110
extract	120
fluidextract	120
powder, compound	120
resin	120
tincture	120
tincture, compound	120
Jamaica ginger tincture (Tinc- tura zingiberis)	232
James' powder (Pulvis anti- monialis)	30
Jamestown weed (Stramonium). Janeway's pills (Piliulae aloes et podophylli compositae) ..	210
Javelle water (Liquor potassae chlorinatae)	167
Jennerian vaccine (Virus vac- cincum)	227
Jimson weed (Stramonium)....	210
Juglans	120
fluidextract	121
Juniper and potassium acetate elixir	168
berries	121
berries, fluidextract	121
berries, oil (Oleum juniper) ..	141
oil	141
spirit	142
spirit, compound	142
tar, oil (Oleum cadinum)....	139
Juniperus	121
Kaolin	121
cataplasm	121
Kava	121
fluidextract	121
Kava Kava (Kava)	121
Kentish's ointment (Linimentum terebinthinae)	148
Kermes mineral (Antimonium sulphuratum)	30
Kieselguhr, purified (Terra si- licea purificata)	219
Kino	121
and opium powder, com- pound	122
and opium tincture, compound composita, tinctura (Tinctura kino et opii composita)....	122
compositus, pulvis (Pulvis kino et opii compositus)....	122
tincture	122
tincture, compound (Tinctura kino et opii composita)....	122
Kissingen salt, artificial	202
artificial, effervescent	202
Kola	122
fluidextract	122
nuts (Kola)	122
Koussou (Brayera)	43
Krameria	122
extract	122
extract, powdered (Extractum krameriae)	122
fluidextract	123
fluidglycerate	123
syrup	123
tincture	123
Kumyss (Lac fermentatum) ...	123
Labarraque's solution (Liquor sodae chlorinatae)	197
Lac fermentatum	123
sulphuris (Sulphur praecip- itatum)	215
vaccinum	123
Lactic acid	10
Lactophosphate of lime, calisaya, and iron, elixir (Elixir of cinchona alkaloids, iron and calcium lactophosphate) ...	50
Lactose (Saccharum lactis)....	188
Lactucarium	123
syrup	123
tincture	124
Lady slipper root (Cypripedium) Webster's dinner pill (Piliulae aloes et mastiches)	77
Lafayette mixture (Mistura co- palbae)	20
Lamotte's drops (Tinctura ferri chloridi aetherea)	73
Lanolin (Adeps lanae hydraus) anhydrous (Adeps lanae)....	87
	15

	PAGE		PAGE
Lapis calaminaris (Calamina praeparata)	46	Licorice—Continued	
Lappa	124	elixir, aqueous (Elixir gly- cyrrhizae aquosum)	102
fluidextract	124	elixir, aromatic (Elixir gly- cyrrhizae aromaticum)	19
Larch agaric (Agaricus)	18	extract (Extractum glycyrr- rhizae)	102
turpentine (Terebinthina la- ricis)	218	fluidglycerate (Fluidglycera- tum glycyrrhizae)	102
Lard	15	powder, compound (Pulvis glycyrrhizae compositus)...	194
benzoinated	15	syrup (Syrupus glycyrrhizae)	103
Larkspur seed	77	Life root (Senecio)	193
tincture	78	Light magnesia (Magnesii oxi- dum)	129
Lassar's mild resorcinol paste (Pasta resorcinolis mitis) ..	182	Lignum vitae (Gualaci lignum).	104
naphthol paste (Pasta beta- naphtholis)	40	Lily-of-the-valley flowers (Con- vallariae flores)	72
stronger resorcinol paste (Pasta resorcinolis fortis) ..	182	root (Convallariae radix)	72
zinc paste (Pasta zinci)	229	Lime (Calx)	52
Laudanum (Tinctura opii)	152	chloride (Calx chlorinata) ...	52
Sydenham's (Tinctura opii crocata)	152	chlorinated	52
Laughing gas (Nitrogenii mo- noxidum)	136	elixir of calisaya, iron and lactophosphate of lime (Elixir of cinchona alka- loids, iron and calcium lactophosphate)	49
Lavender oil	142	juice	214
spirit	142	juice and pepsin	157
spirit, compound (Tinctura lavandulae composita) ...	142	liniment	52
tincture, compound	142	potassa with	172
Laxative elixir (Elixir cascarae sagradae compositum)	57	soda with	198
pills, compound	22	sulphurated (Calci sulphidum crudum)	50
pills, post partum	71	sulphurated, solution	51
species	195	water (Liquor calcis)	52
Lead acetate	165	Limonis cortex	124
and opium lotion	165	Liniment, acetic turpentine....	148
and opium pills	151	ammonia	25
and opium wash (Lotio plumbi et opii)	16	belladonna	39
carbonate	166	camphor	53
iodide	166	camphorated soap	190
oxide	166	Canada (Linimentum opii com- positum)	149
oxide, red	166	chloroform	62
plaster	166	hartshorn (Linimentum am- moniae)	25
red (Plumbi oxidum rubrum)	166	lime	52
subacetate	167	of aconite and chloroform....	15
subacetate solution	167	of ammonium iodide	28
subacetate solution, diluted..	167	of croton oil	149
subacetate, cerate	167	of croton oil, compound	149
sugar (Plumbi acetas)	165	of mustard compound	147
water (Liquor plumbi sub- acetatis dilutus)	167	of opium compound	149
white (Plumbi carbonas)	166	of soft soap	190
Lemon oil	143	of soft soap compound	190
peel	124	St. John Long's (Linimentum terebinthinae aceticum) ...	148
peel tincture	124	soap	190
Leptandra	124	Stokes' (Linimentum tere- binthinae aceticum)	148
extract	124	turpentine	148
extract, powdered	124		
fluidextract	124		
Licorice (Glycyrrhiza)	102		
elixir (Elixir glycyrrhizae)...	19		

	PAGE
Liniment—Continued	
volatile (Linimentum am-	
moniae)	25
Liniments	124
Linimentum aconiti et chloro-	
formi	15
album (Linimentum tere-	
binthinae aceticum)	148
ammonii iodidi	28
calcis	52
saponis mollis	190
saponis mollis compositum ..	190
sinapis compositum	147
terebinthinae	148
terebinthinae aceticum	148
tiglli	149
tiglli compositum	149
Linseed	124
oil	143
oil, raw (Oleum lini)	143
Linum	124
Liquefied carbolic acid (Phenol	
liquefactum)	160
phenol	160
Liquid apiol (Oleo-resina petro-	
selini)	159
glucose (Glucosum)	101
opodeldoc (Linimentum sa-	
ponis)	190
paraffini (Petrolatum liqul-	
dum)	158
petrolatum	158
petrox (Petroxolinum liqul-	
dum)	158
petroxolin	158
storax (Styrax)	214
Liquor	
See also under Solution.	
acidi arsenosi	34
alumni acetatis	23
alumni acetatis (Liquor	
alumni subacetatis)	24
alumni acetico-tartratis	24
alumni subacetatis	24
ammoniae anisatus (Spiritus	
ammoniae anisatus)	25
ammonii acetatis	25
ammonii citratis	27
antisepticus	7
antisepticus alkalinus	169
arseni et hydrargyri iodidi ..	33
arsenicalis (Liquor potassii	
arsenitis)	168
arsenicalis Clemens	168
arsenicalis Fowleri (Liquor	
potassii arsenitis)	168
arsenicalis Pearsoni (Liquor	
sodii arsenatis, Pearson) ..	198
auri et arseni bromidi	7
bismuthi	42
bromi	43
Burovi (Liquor alumni	
acetatis)	23

	PAGE
Liquor—Continued	
calcis	52
calcis sulphuratae	51
carmini	56
chlori compositus	62
cocci	67
cresolis compositus	75
ferri acetatis	85
ferri albuminati	94
ferri chloridi	86
ferri citratis	87
ferri et ammonii acetatis	25
ferri hypophosphitis	89
ferri nitratis	91
ferri oxychloridi	91
ferri oxysulphatis	91
ferri peptonati	94
ferri peptonati cum mangano	
(Liquor ferri peptonati et	
mangani)	95
ferri peptonati et mangani ..	95
ferri protochloridi	92
ferri salicylatis	206
ferri subsulphatis	93
ferri tersulphatis	94
formaldehydi	96
guttae perchae	106
hydrargyri et potassii iodidi ..	110
hydrargyri nitratis	110
hydrastinae compositus	113
hydrogenii dioxidi	115
hypophosphitum	48
hypophosphitum compositus ..	48
hypophysis	116
iodi carbolatus (Liquor iodi	
phenolatus)	118
iodi compositus	118
iodi phenolatus	118
magnesi citratis	128
magnesi sulphatis efferves-	
cens	129
pancreaticus (Liquor pancrea-	
tini)	153
pepsini	157
pepsini antisepticus	157
pepsini aromaticus	157
phosphatum acidus	11
phosphatum compositus	11
phosphori	161
pilis alkalinus	165
pilis carbonis	165
plumbi subacetatis	167
plumbi subacetatis dilutus ..	167
potassae (Liquor potassii hy-	
droxii)	172
potassae chlorate (Liquor po-	
tassae chlorinatae)	167
potassae chlorinatae	167
potassii arsenatis et bromidi	
(Liquor arsenicalis clemens)	168
potassii citratis	171
potassii hydroxidi	172
sodae (Liquor sodii hydroxidi)	204

	PAGE		PAGE
Liquor—Continued		Magmas	128
sodae chlorinatae	197	Magnesia (Magnesii oxidum)	129
sodae et menthae	199	and rhubarb powder, anisated	184
sodii arsenatis	198	asafoetida and opium mixture	128
sodii arsenatis, Pearson	198	calcined (Magnesii oxidum)	129
sodii boratis compositus	200	heavy (Magnesii oxidum pon-	
sodii chloridi physiologicus	202	derosum)	129
sodii citratis	202	light (Magnesii oxidum)	129
sodii citro-tartratis efferves-		magma	128
cens	202	milk (Magma magnesiae)	128
sodii glycerophosphatis	203	with ferric hydrate (Ferri hy-	
sodii hydroxidi	204	dioxidum cum magnesii	
sodii phosphatis compositus	205	oxido	88
strychninae acetatis	212	Magnesium carbonate	128
zinci chloridi	229	chloride	128
zinci et aluminii compositus	230	citrate	128
zinci et ferri compositus	231	citrate solution	128
Liquores	125	hydroxide	128
Liquorice root (Glycyrrhiza)	102	oxide	129
Litharge (Plumbi oxidum)	166	oxide heavy	129
Lithium bromide	125	oxide with ferric hydroxide	88
bromide elixir	125	sulphate	129
carbonate	125	sulphate solution, effervescent	129
citrate	126	Male fern (Aspidium)	35
citrate, effervescent salt	126	oleoresin (Oleoresina aspidii)	35
citrate elixir	126	Mallow leaves	130
salicylate	126	Malt	129
salicylate elixir	126	and cod liver oil emulsion	144
with effervescent artificial		extract	129
vichy salt	200	Malvae folia	130
Liver of sulphur (Potassa sul-		Mandrake (Podophyllum)	167
phurata)	168	Manganese and iron peptonate	
Lobelia	126	solution	95
fluidextract	126	and sodium citrate (Mangani	
tincture	127	citras solubilis)	130
Logwood (Haematoxylon)	106	citrate, soluble	130
London paste (Soda cum calce)	198	dioxide, precipitated	130
Long buchu	44	glycerophosphate, soluble	130
Lotio ammoniacalis camphorata	25	hypophosphite	130
flava	111	iodide and iron, syrup	90
nigra	109	sulphate	131
Lotion of lead and opium	165	Manganous glycerinophosphate,	
Vlemineckx' (Liquor calcis sul-		soluble (Mangani glycerophos-	
phuratae)	51	phosphas solubilis)	130
yellow	111	Manna	131
Lotions	127	syrup	131
Lozenges (Trochisci)	223	Marigold (Calendula)	51
Lugol's solution (Liquor Iodi		Marsh mallow leaves (Althaeae	
compositus)	118	folia)	23
Lunar caustic (Argenti nitras		mallow root (Althaea)	22
fusus)	33	trefoll (Menyanthes)	133
Lupulin	127	Mass of copaiba	73
fluidextract	127	of ferrous carbonate	85
oleoresin	127	of mercury	112
Lycopodium	127	Vallet's (Massa ferri carbon-	
		atis)	85
Mace	127	Masses	131
Macrotys (Cimicifuga)	63	Mastic	
Madderwort (Absinthium)	5	and aloes pills	
Magma bismuthi	42	Matico	
ferri hydroxidi	89	fluidextract	
magnesiae	128	leaves (Matico)	

INDEX

261

PAGE	PAGE
Matricaria	131
May apple rhizome (Podophyl- lum)	167
Meadow anemone (Pulsatilla) ..	174
Mel	132
boracis (Mel sodii boratis) ...	200
depuratum	132
rosae	132
rosae et sodii boratis	200
sodii boratis	200
Mellot	132
Mellita	132
Mentha piperita	132
viridis	132
Menthol	132
and camphor (Menthol cam- phoratum)	133
camphorated	133
inunction	132
inunction compound	133
petrox (Petroxolinum men- tholis)	133
petroxolin	133
spray	133
spray compound	133
Menyanthes	133
Mercurial ointment	112
ointment, diluted	112
Mercuric chloride (Hydrargyri chloridum corrosivum)	108
chloride, corrosive	108
chloride, corrosive, poison tab- lets	109
chloride mull, corrosive	109
iodide (Hydrargyri iodidum rubrum)	110
iodide and arsenous solution. .	33
iodide, red	110
nitrate	110
nitrate ointment	111
nitrate solution	110
oxide, red	111
oxide, red, ointment	111
oxide, yellow	111
oxide, yellow, ointment	111
salicylate	111
subsalicylate (Hydrargyri sal- icylas)	111
Mercurous chloride (Hydrargyri chloridum mite)	109
chloride and jalap, mild, pow- der	110
chloride, mild	109
iodide (Hydrargyri iodidum flavum)	110
iodide, yellow	110
Mercury	112
ammoniated	113
ammoniated	113
Mercury—Continued	
bichloride (Hydrargyri chlori- dum corrosivum)	108
biniodide (Hydrargyri iodidum rubrum)	110
digitalis, and squill, pills.	79
iodide, red (Hydrargyri iodi- dum rubrum)	110
iodide, yellow (Hydrargyri iodidum flavum)	110
mass	112
oleate	112
perchloride (Hydrargyri chlori- dum corrosivum)	108
petrox (Petroxolinum hydrar- gyri)	112
petroxolin	112
podophyllum and aloes pills. .	21
protochloride (Hydrargyri chloridum mite)	109
protiodide (Hydrargyri iodi- dum flavum)	110
subchloride (Hydrargyri chlo- ridum mite)	109
with chalk	112
Metallic pills (Pilulae ferri, quininae, strychninae et ar- seni fortiores)	95
Methyl salicylate	133
salicylate petrox (Petroxoli- num methylis salicylatis) ..	133
salicylate petroxolin	133
Methylene blue (Methylthioninae chloridum)	133
Methylmorphine (Codeina)	68
Methylthioninae hydrochloridum (Methylthioninae chloridum) .	133
Methylthionine chloride	133
Methysticum (Kava)	121
Mezereon (Mezereum)	134
Mezereum	134
fluidextract	134
Mild mercurous chloride	109
mercurous chloride and jalap powder	110
pills of iron, quinine, strych- nine and arsenic	95
resorcinol paste	182
resorcinol paste, Lassar's (Pasta resorcinolis mitis) ..	182
Milk, cow's	123
fermented	123
of almond (Emulsum amyg- dalaе)	29
of asafetida (Emulsion asa- foetidae)	34
of bismuth (Magma bismuthi) of magnesia (Magma magne- sia)	42
fermented	128
sulphur (Sulphur praecipit- um)	215
(Saccharum lactis)	188

	PAGE		PAGE
Mindererus spirit (Liquor ammonii acetatis)	25	Mixture—Continued	
Mineral oil (Petroleum liquidum)	158	of oil of tar	145
Mint and soda, solution	199	of opium and chloroform compound	150
soda (Liquor sodae et menthae)	199	of opium and rhubarb compound	150
Mistura		of opium and sassafras	150
See also under Mixture.		of rhubarb, alkaline	183
adstringens et escharotica (Mistura adstringens)	77	of rhubarb and soda (Mistura rhei composita)	183
ammonii chloridi	27	of rhubarb compound	142
antidysenterica (Mistura camphorae acida)	149	Parrish's camphor (Mistura camphorae aromatica)	142
chloroformi et cannabis indicae composita (Mistura chloroformi et morphinae composita)	62	Squibb's diarrhea (Mistura opii et chloroformi composita)	150
cretae	76	sun cholera (Mistura opii et rhei composita)	150
olei picis	145	tar (Mistura olei picis)	145
opii alkalina (Mistura opii et sassafras)	150	Villate's (Mistura adstringens)	77
pectoralis Stokes	26	Mixtures	134
picis liquidae (Mistura olei picis)	145	Molded silver nitrate	32
sassafras et opii (Mistura opii et sassafras)	150	Monkshood (Aconitum)	14
sodae et menthae (Liquor sodae et menthae)	199	Monobromated camphor	53
sodii citratis (Liquor sodii citratis)	202	Monrel's solution (Liquor ferri subsulphatis)	93
solvens simplex (Mistura ammonii chloridi)	27	Morphine	134
Mixture, acid camphor	149	and acacia syrup	134
aromatic camphor	142	and chloroform mixture, compound (Morphinae hydrochloridum)	134
astringent	77	hydrochloride	135
Basham's (Liquor ferri et ammonii acetatis)	25	sulphate	135
brown (Mistura glycyrrhizae composita)	150	with compound syrup of white pine	135
carminative	150	Moschus	135
chalk	76	Motherplaster, camphorated (Emplastrum fuscum camphoratum)	53
Chapman's (Mistura copaibae et opii)	73	Mother's salve (Unguentum fuscum)	53
copaiba	73	Mucilage of acacia	5
copaiba and opium	73	of chondrus	63
Griffith's (Mistura ferri composita)	86	of Irish moss (Mucilago chondri)	63
Hope's (Mistura camphorae acida)	149	of sassafras pith	181
iron, compound	86	of tragacanth	221
Lafayette (Mistura copaibae)	73	Mucilages	135
of ammonium chloride	27	Mull, corrosive mercuric chloride	109
of chloral and potassium bromide compound	61	salicylic acid	12
of chloroform and morphine compound	62	zinc	229
of glycyrrhiza compound	150	Mulla creosoti salicylata	12
of gualac	105	Mullein flowers	995
of iron, salicylate (Liquor ferri salicylatis)	206	leaves (Verbasci folia)	
of magnesia, asafetida and opium	128	leaves, fluidextract	
		Mulls	
		Muriatic acid (Acidum chloricum)	
		Musk	
		deer (Moschus) ..	

	PAGE		PAGE
Musk—Continued		Nitromuriatic acid—Continued	
root (Sumbul)	216	acid, diluted (Acidum nitro-	
root, fluidextract (Fluidextrac-		hydrochloricum dilutum)...	10
tum sumbul)	216	Nitrous ether	17
tincture	135	ether spirit	17
tonquin (Moschus)	135	oxide (Nitrogenii monoxidum)	136
Mustard, black	197	Normal salt solution (Liquor	
brown (Sinapis nigra).....	197	sodii chloridi physiologicus)	202
liniment, compound	147	Nutgall	98
oil (Oleum sinapis volatile)..	147	ointment	98
oil, volatile	147	tincture	98
paper (Emplastrum sinapis)..	197	Nutmeg (Myristica)	136
plaster	197	oil (Oleum myristicae).....	145
spirit	147	Nux vomica	136
white	197	and phosphorus elixir	161
yellow (Sinapis alba).....	197	extract	136
Mutton suet (Sevum praepara-		extract, powdered (Extractum	
tum)	197	nucis vomicae)	136
Myrcia oil	145	fluidextract	136
spirit, compound	145	iron, quinine and aloes pills..	94
Myrica	135	tincture	137
Myristica	136	Oil, anise (Oleum anisi).....	138
oil	145	benne (Oleum sesami)	147
Myrrh	136	camphorated (Linimentum	
and aloes pills	20	camphorae)	53
and aloes tincture	21	carbolyzed (Oleum phenola-	
and capsicum tincture.....	55	tum)	159
gum (Myrrha)	136	cassia (Oleum cassiae)	140
tincture	136	castor	146
Naphthol (Betanaphthol)	40	castor, aromatic	146
paste, Lassar's (Pasta beta-		castor, emulsion	146
naphtholis)	40	cod liver	144
Nebula aromatica	158	cod liver, emulsion	144
eucalyptolis	83	cod liver, with calcium lacto-	
mentholis	133	phosphate emulsion	144
mentholis composita	133	cod liver, with calcium phos-	
thymolis	220	phate emulsion	144
Nebulae	136	cod liver, with egg emulsion..	145
Neroli, oil (Oleum aurantii flo-		cod liver, with hypophosphites	
rum)	138	emulsion	144
Neutralizing cordial (Mistura		cod liver, with malt emulsion.	144
rhei alkalina)	183	cod liver, with phosphate of	
Niemeyer pills for dropsy (Pilu-		lime emulsion (Emulsium olei	
lae digitalis, scillae et hy-		morrhuae cum calci phos-	
drargyri)	79	phate)	144
pills for phthisis (Pilulae opil,		cod liver, with wild cherry	
digitalis et quiniinae)	151	emulsion	144
Night blooming cereus (Cactus		cottonseed	141
grandiflorus)	45	croton	148
Nitre, sweet spirit (Spiritus		croton, collodion	148
aetheris nitrosi)	17	croton, liniment	149
Nitric acid	10	croton, liniment compound....	149
Nitrogen monoxide	136	etheral	137
Nitroglycerin (Glycerylis nitras)	101	linseed	143
pills	109	linseed, raw (Oleum lini)....	143
spir		mineral (Petrolatum liqui-	
		dum)	158
		mustard (Oleum sinapis vola-	
		tile)	147
		(Oleum pimentae)	146
		mustard	138
		mustard (Oleum	
		mustard	140

	PAGE		PAGE
Oil—Continued		Oil—Continued	
of anise	138	of wintergreen (Methylis salicylas)	133
of bay (Oleum myrciae)	145	olive	145
of bergamot	139	phenolated	159
of birch, rectified empyroligneous (Oleum betulae empyreumaticum rectificatum) ..	139	phosphorated	161
of birch tar, rectified	139	pimento (Oleum pimentae) ..	146
of bitter almond	137	rosemary (Oleum rosmarini) ..	146
of bitter orange	138	santalwood (Oleum santali) ..	146
of cade	139	sesame	147
of cajuput	139	spray, aromatic	158
of caraway	140	tar, rectified (Oleum picis liquidae rectificatum)	145
of cardamom	139	teel (Oleum sesami)	147
of chenopodium	140	turpentine (Oleum terebinthinae)	147
of cinnamon	140	turpentine, rectified (Oleum terebinthinae rectificatum) ..	147
of clove	140	Oils, infused	137
of coriander	141	volatile spirits	208
of cubeb	141	Oil-Sugars	137
of dwarf pine needles	146	Ointment	15
of eucalyptus	141	basilicon (Ceratum resinae) ..	181
of fennel	141	belladonna	38
of flaxseed (Oleum lini)	143	blue (Unguentum hydrargyri dilutum)	112
of hyoscymus, compound	115	brown	53
of juniper	141	calamine	46
of juniper berries (Oleum juniperi)	141	camphor	53
of juniper tar (Oleum cadinum)	139	chrysarobin	63
of lavender	142	citrine (Unguentum hydrargyri nitratis)	111
of lemon	143	diachylon	166
of mustard, volatile	147	Hebra's itch (Unguentum sulphuris compositum)	216
of myrcia	145	iodine	119
of myristica	145	iodoform	117
of neroli (Oleum aurantii florum)	138	Kentish's (Linimentum terebinthinae)	148
of nutmeg (Oleum myristicae) ..	145	mercurial	112
of orange	138	mercurial, diluted	112
of orange flowers	138	nutgall	98
of peppermint	143	of ammoniated mercury	113
of pimenta	146	of boric acid	7
of rosemary	146	of carbolic acid (Unguentum phenolis)	160
of sandalwood (Oleum santali) ..	146	of lead iodide	166
of santal	146	of mercuric nitrate	111
of sassafras	147	of phenol	160
of spearmint	143	of potassium iodide	173
of star anise	138	of red mercuric oxide	111
of sweet almond (Oleum amygdalae expressum)	138	of rose water	186
of sweet birch (Methylis salicylas)	133	of tannic acid	14
of sweet orange (Oleum aurantii)	138	of yellow mercuric oxide	111
of tar mixture	145	of zinc oxide	230
of tar, rectified	145	of zinc stearate	230
of teaberry (Methylis salicylas)	133	petrolatum (Petrolatum)	158
of theobroma	148	resorcinol, compound	182
of thyme	148	simple (Unguentum)	15
of turpentine	147	stramonium	210
of turpentine emulsion	148	sulphur	215
of turpentine, rectified	147	sulphur, alkaline	215
		sulphur compound	216

	PAGE
Ointment—Continued	
tar	165
tar, compound	146
veratrine	225
white precipitate (Unguentum hydrargyri ammoniati)	113
Wilkinson's (Unguentum sulphuris compositum)	216
zinc (Unguentum zinci oxidi) ..	230
Ointments	223
Olea infusa	137
Oleate of aconitine	14
of atropine	36
of cocaine	67
of mercury	112
of quinine	177
of veratrine	225
Oleates	137
Oleic acid	11
Oleo-balsamic mixture	142
Oleoresin of aspidium	35
of capsicum	55
of cubeb	76
of ginger	232
of lupulin	127
of male fern (Oleoresina aspidii)	35
of parsley fruit	159
of pepper	164
Oleoresina petroselinii	159
piperis	164
zingiberis	232
Oleoresins	137
Oleosacchara	137
Oleum	
See also under Oil.	
aethereum	137
amygdalae amarae	137
amygdalae expressum	138
aurantii	138
aurantii amari	138
aurantii corticis (Oleum aurantii)	138
aurantii florum	138
bergamottae	139
betulae (Methylis salicylas) ..	133
betulae empyreumaticum rectificatum	139
carli	140
caryophylli	140
foeniculi	141
gaultheriae (Methylis salicylas)	133
gossypii seminis	141
Jecoris aselli (Oleum morrhuae)	144
lini	143
menthae piperitae	143
menthae viridis	143
morrhuae	144
picis liquidae (Oleum picis liquidae rectificatum)	145

	PAGE
Oleum—Continued	
picis liquidae rectificatum....	145
pinii pumilionis	146
ricini	146
ricini aromaticum	146
rusci rectificatum (Oleum rusci betulae empyreumaticum rectificatum)	139
sinapis volatile	147
terebinthinae	147
terebinthinae rectificatum....	147
tiglli	148
Olive oil	145
Opium	149
and camphor pills	151
and chalk powder, aromatic... and chloroform mixture, compound	151
and copaiba mixture	150
and ipecac powder	73
and ipecac syrup	151
and ipecac tincture	152
and kino powder, compound... and kino tincture, compound..	122
and lead lotion	122
and lead pills	165
and lead wash (Lotio plumbi et opii)	151
and rhubarb mixture, compound	165
and sassafras mixture	150
deodorized	150
deodorized tincture	151
digitalis and quinine pills....	152
extract	151
extract, powdered (Extractum opii)	149
granulated	149
liniment, compound	151
magnesia, and asafetida mixture	149
powdered	128
tincture	150
tincture, camphorated	152
vinegar	152
with saffron tincture	149
Opodeldoc (Linimentum saponato-camphoratum)	152
liquid (Linimentum saponis) ..	190
solid (Linimentum saponata-camphoratum)	190
Orange, elixir, compound (Vinum aurantii compositum)	36
flower water	37
flower water, stronger	37
flowers, bitter	37
flowers oil	138
flowers syrup	37
oil	138
peel, bitter	36
peel, bitter, fluidextract	36
peel, bitter, tincture	36

	PAGE		PAGE
Orange—Continued		Paste—Continued	
peel, sweet	36	Lassar's naphthol (Pasta	
peel, sweet, tincture	37	betanaphtholis)	40
spirit, compound	138	Lassar's stronger resorcinol	
syrup	36	(Pasta resorcinolis fortis) ..	182
wine, compound	36	Lassar's zinc (Pasta zinci) ..	229
Ordeal bean (Physastigma)	161	London (Soda cum calce)	198
Oregon grape root (Berberis)	40	pencils	209
Orphol (Bismuthi betanaph-		resorcinol, mild	182
tholas)	41	resorcinol, strong	182
Orris	119	Unna's soft zinc (Pasta zinci	
root (Iris)	119	mollis)	229
Ovi albumen recens	152	Unna's sulphurated zinc	
vitellum recens	152	(Pasta zinci sulphurata) ..	230
Ovum gallinaceum	153	zinc	229
Oxgall	85	zinc, soft	229
extract	85	zinc, sulphurated	230
extract, powdered (Extractum		Pastes, dermatologic	155
fellis bovis)	85	Pearson's solution of sodium	
Oxygen	153	arsenate	198
Oxymel of Squill	192	Pectoral drops (Tinctura pec-	
toralis)		drops, Bateman's (Tinctura	
Pale catechu (Gambir)	98	pectoralis)	152
catechu tincture, compound		species	22
(Tinctura gambir com-		tincture	152
posita)	98	Pectoralis, syrupus (Syrupus	
Pancreatic solution (Liquor		morphinae et acaciae)	134
pancreatini)	153	Pelletierine tannate	155
Pancrreatin	153	Pellitory root (Pyrethrum) ..	175
powder, compound	153	tincture (Tinctura pyrethri) ..	175
solution	153	Pencils, paste	209
Papaveris fructus	153	Unna (Still diluables)	209
Papers	60	Pepo	155
Papoose root (Caulophyllum) ..	59	Pepper	164
Paracoto	153	black (Piper)	164
fluidextract	154	cayenne (Capsicum)	55
tincture	154	oleoresin	164
Paraffin	154	Peppermint	132
liquid (Petrolatum liquidum) ..	158	essence (Spiritus menthae	
Paraform (Paraformaldehydum) ..	154	piperitae)	143
Paraformaldehyde	154	oil	143
Paraldehyde	154	spirit	143
Paregoric (Tinctura opii cam-		troches	143
phorata)	152	water	143
Pareira	155	Pepsin	155
brava (Pareira)	155	and bismuth elixir	156
fluidextract	155	and iron elixir	156
Parrish's camphor mixture (Mis-		and lime juice	157
tura camphorae aromatica) ..	142	and rennin elixir, compound ..	156
Parsley fruit	159	bismuth and strychnine elixir	
fruit, oleoresin	159	cinchona alkaloids and iron	
root	159	elixir	156
root, fluidextract	159	elixir	156
seed (Petroselinum)	159	glycerite	157
Pasque flower (Pulsatilla)	174	iron and calisaya, alkaloidal,	
Passion flower	155	elixir (Elixir cinchonae	
flower tincture	155	alkaloidorum, ferri et pep-	
vine (Passiflora)	155	sini)	
Paste, betanaphthol	40	saccharated	
dextrinated	78	solution	
Lassar's mild resorcinol		solution, antiseptic	
(Pasta resorcinol mitis)	182		

PAGE	PAGE
Pepsin—Continued	Phenol—Continued
solution, aromatic 157	iodized 160
wine 157	liquefied 160
Peptonate of iron and manga-	ointment 160
nese solution 95	petrox (Petroxolinum phe-
of iron solution 94	nolis) 160
Peptonizing powder (Pulvis	petrox, camphorated (Petrox-
pancreatini compositus) . . . 153	olinum phenolis camphora-
Perchloride of mercury (Hy-	tum) 160
drargyri chloridum corro-	Phenolated oil 159
sivum) 108	solution of iodine 118
Perfumed spirit 139	water 159
Peroxide, hydrogen (Hydrogenii	Phenolphthalein 160
dioxidum) 114	troches 160
Persio 157	Phenyl salicylate 160
Petrolatum 158	Phenylcinchoninic acid 11
album 158	Phenyldimethylpyrazoylon (Anti-
emulsion 158	pyrina) 31
liquid 158	Phenylis salicylas 160
ointment (Petrolatum) 158	Phenyl - quinoline - carboxylic
saponatum liquidum (Petrox-	acid (Acidum phenylcincho-
olinum liquidum) 158	ninicum) 11
saponatum spissum (Petroxoli-	Phosphate of lime with cod liver
num spissum) 159	oil emulsion (Emulsum olei
white 158	morrhuae cum calcii phos-
Petroleum benzin, purified. . . . 39	phate) 144
ether (Benzinum purificatum) . 39	of iron, quinine and strych-
jelly (Petrolatum) 158	nine, syrup of 92
jelly, white (Petrolatum al-	quinine and strychnine syrup.
bum) 158	solution, acid 11
Petroselinii radix 159	solution, compound 11
Petroselinum 159	syrup, compound 11
Petrox, liquid (Petroxolinum	Phosphorated oil 161
liquidum) 158	Phosphates, acid solution of. . . 11
solid (Petroxolinum spissum) . 159	Phosphoric acid 11
Petroxolin, liquid 158	acid, diluted 11
solid 159	Phosphorus 161
Petroxollins 159	and nux vomica elixir. 161
Petroxollinum betanaphtholis . . 41	elixir 161
cadini 139	pills 161
chloroformi camphoratum. . . . 62	solution 161
creosoti 75	solution, Thompson's (Liquor
eucalyptolis 83	phosphori) 161
gualacolis 105	Physiological salt solution
hydrargyri 112	(Liquor sodii chloridi
iodi 118	physiologicus) 202
iodi dilutum 118	solution of sodium chloride. . 202
iodoformi 117	Physostigma 161
mentholis 133	extract 162
methyllis salicylatis 133	extract, powdered (Extractum
phenolis 160	physostigmatis) 162
phenolis camphoratum 160	tincture 162
picis 145	Physostigmine salicylate 162
spissum 159	Phytolacca
sulphuratum 215	fluid ^{ext}
sulphuratum compositum 215	Pier
terebinthinae laricis 218	Pill
Pheasant's eye (Adonis) 16	
Phenacetin (Acetphenetidinum). . 6	
Phenazone (Antipyrina)	
Phenol	
glycerite	
iodatum	

	PAGE		PAGE
Pills—Continued		Pills—Continued	
antidyspeptic	212	of colocynth, compound.....	71
antiperiodic	180	of digitalis, squill and mer-	
antiperiodic, without aloes ...	180	cury	79
Barker's post partum (Pilulae		of ferrous carbonate.....	86
laxativae post partum).....	71	of ferrous iodide	90
bitter metallic (Pilulae ferri,		of glonoin (Pilulae glycerylis	
quininae, strychninae et		nitratil)	102
arseni fortiores)	95	of iron, quinine, aloes and	
Blaud's (Pilulae ferri car-		nux vomica	94
bonatis)	86	of iron, quinine, strychnine	
cathartic, compound	71	and arsenic, mild	95
cathartic, vegetable	71	of iron, quinine, strychnine	
chalybeate (Pilulae ferri car-		and arsenic, stronger	95
bonatis)	86	of nitroglycerin	102
cochia (Pilulae colocynthidis		of opium and camphor.....	151
compositae)	71	of opium and lead	151
dinner	20	of opium, digitalis and quinine	
dinner, Chapman's	20	151
dinner, Cole's	20	of phosphorus	161
dinner, Hall's	20	of rhubarb	184
dinner, Lady Webster's (Pilu-		of rhubarb, compound	184
lae aloes et mastiches).....	20	Plummer's (Pilulae antimonii	
ferruginous (Pilulae ferri car-		compositae)	31
bonatis)	86	post partum laxative.....	71
Francis' triplex (Pilulae		quadruplex (Pilulae ferri,	
aloes, hydrargyri et scam-		quinine, aloes et nucis	
monii compositae)	21	vomicae)	94
Guy's (Pilulae digitalis, scil-		quatuor (Pilulae ferri, qui-	
lae et hydrargyri)	79	ninae, aloes et nucis vom-	
Janeway's (Pilulae aloes et		icae)	94
podophylli compositae)	21	triplex (Pilulae aloes, hy-	
laxative, compound	22	drargyri et podophylli)	21
metallic (Pilulae ferri, qui-		Warburg's (Pilulae antiperi-	
ninae, strychninae et arseni		odicae)	180
fortiores)	95	Warburg's, without aloes	
Niemeyer, for dropsy (Pilulae		(Pilulae antiperiodicae sine	
digitalis, scillae et hydrar-		aloe)	180
gyri)	79	Pilocarpine chloride (Pilocar-	
Niemeyer, for phthisis (Pilulae		pinae hydrochloridum)	162
opli, digitalis et quinae).....	151	hydrochloride	162
of aloes	20	nitrate	163
of aloes and asafetida.....	20	Pilocarpus	163
of aloes and iron	20	fluidextract	163
of aloes and mastic	20	Pilulae	
of aloes and myrrh	20	See also under Pills.	
of aloes and podophyllum,		ad prandium	20
compound	21	cocclae (Pilulae colocynthidis	
of aloes, mercury and podo-		compositae)	71
phyllum	21	glycerylis nitratil)	102
of aloes, mercury and scam-		metallorum (Pilulae ferri,	
mony, compound	21	quininae, strychninae et	
of aloin, compound	22	arseni fortiores)	95
of aloin, strychnine and bella-		metallorum amarae (Pilulae	
donna	22	ferri, quinae, strychninae	
of aloin, strychnine and bella-		et arseni fortiores)	95
donna, compound	22	quadruplices (Pilulae ferri,	
of antimony, compound	31	quininae, aloes et nucis	
of asafetida	34	vomicae)	94
of colocynth and hyoscyamus.		triplices (Pilulae aloes, hy-	
of colocynth and podophyllum		drargyri et podophylli)	21
	71	Pimento (Pimenta)	163
	71	oil (Oleum pimentae)	146
		Pimpinel root (Pimpinella),...	164

	PAGE		PAGE
Pimpinella	164	Poplar buds, balsam	167
tincture	164	Poppy capsules	153
Pine bark, white	164	syrup	153
needles, dwarf, oil	146	Populi gemmae	167
oil, dwarf (Oleum pini pu- millionis)	146	Post partum laxative pills	71
tar (Pix liquida)	164	partum pills, Barker's (Pilu- lae laxative post partum) ..	71
Pinkroot (Spigelia)	208	Potassa, chlorinated	167
fluidextract (Fluidextractum spigeliae)	208	chlorinated, solution	167
Pinus alba	164	cum calce	172
Pipe gamboge (cambogia)	52	solution (Liquor potassii hy- droxidi)	172
Piper	164	sulphurated	168
Pipsissewa (Chimaphila)	60	with lime	172
Pituitary body, desiccated (Hy- pophysis sicca)	116	Potassium acetate	168
solution (Liquor hypophysis) ..	116	acetate and buchu, elixir of ..	44
Pix carbonis (Pix Lithanthracis)	165	acetate and juniper elixir	168
liquida	164	acetate elixir	168
lithanthracis	165	and sodium tartrate	171
Plaster, adhesive (Emplastrum resinae)	182	arsenate	168
adhesive, rubber (Emplastrum elasticum)	182	arsenate and bromides solu- tion (Liquor arsenicalls clemens)	168
belladonna	38	arsenite	168
camphorated brown	53	arsenite solution	168
cantharides	54	bicarbonate	169
capsicum	55	bitartrate	169
diachylon (Emplastrum plum- bi)	166	bitartrate and sulphur troches bromide	215
lead	166	bromide and chloral mixture, compound	61
mustard	197	bromide, effervescent salt	170
rosin	182	bromide, effervescent salt, compound	170
rosin adhesive (Emplastrum resinae)	182	bromide elixir	170
rubber	182	bromide with caffeine, efferv- escent (Sal potassi bromidi effervescens compositus) ..	170
soap	190	carbonate	170
Plasters	82	chlorate	171
Pleurisy root (Asclepias)	35	chlorate troches	171
Plumbi acetat	165	chloride	171
carbonas	166	citrate	171
iodidum	166	citrate, effervescent	171
oxidum	166	citrate solution	171
oxidum rubrum	166	hydrate (Potassii hydroxidum)	172
subacetat	167	hydroxide	172
Plummer's pills (Pilulae anti- monii compositae)	31	hydroxide solution	172
Podophyllin (Resina Podophylli)	167	hypophosphite	172
Podophyllum	167	iodide	172
aloes and mercury pills	21	iodide and mercury, solution ..	110
and aloes pills, compound	21	iodide ointment	
and colocynth pills	71	iodohydrargyrate solution quor hydrargyri et iodidi)	
extract of	167	nitrate	
fluidextract	167	nitrate pa	
resin	167	permangan	
Poison hemlock (Conium)	72	sulphate	
tablets	221	tartrate	
tablets of corrosive mercuric chloride	109	Potio	
Poke root (Phytolacca)	162	citr	
Pomegranate	103	Poultice	
bark (Granatum)	103		
fluidextract	103		

	PAGE		PAGE
Powder, acetanilid, compound...	6	Powdered—Continued	
anise, compound (Pulvis rhei et magnesiae anisatus)	184	extract of euonymus (Extrac- tum euonymi)	84
antimonial	30	extract of gelsemium (Extrac- tum gelsemi)	99
aromatic	66	extract of hydrastis (Extrac- tum hydrastis)	114
chalk, compound	76	extract of ignatia (Extractum ignatiae)	116
composition (Pulvis myricae compositus)	135	extract of krameria (Extrac- tum krameriae)	122
Dover's (Pulvis ipecacuanhae et opii)	151	extract of leptandra (Extrac- tum leptandrae)	124
Dover's syrup (Syrupus ipe- cacuanhae et opii)	151	extract of nux vomica (Ex- tractum nucis vomicae)	136
Dover's tincture (Tinctura ipecacuanhae et opii)	152	extract of opium (Extractum opii)	149
gray (hydrargyrum cum creta)	112	extract of oxgall (Extractum fellis bovis)	85
Gregory's (Pulvis rhei com- positus)	184	extract of physostigma (Ex- tractum physostigmatis)	162
James' (Pulvis antimonialis) ..	30	extract of quassia (Extractum quassiae)	176
licorice, compound (Pulvis glycyrrhizae compositum) ..	194	extract of rhubarb (Extrac- tum rhei)	183
of aloes and canella	21	extract of viburnum pruni- folium (Extractum viburni prunifolii)	227
of bayberry compound	135	opium	150
of chalk and opium, aromatic ..	151	Powders	175
of chalk, aromatic	66	Precipitated calcium carbonate ..	46
of gambir, compound	98	calcium phosphate	50
of glycyrrhiza, compound	194	chalk (Calci carbonas prae- cipitatus)	46
of ipecac and opium	151	ferrous sulphate (Ferri sul- phas granulatus)	94
of ipecac, compound (Pulvis ipecacuanhae et opii)	151	manganese dioxide	130
of jalap, compound	120	sulphur	215
of kino and opium, compound ..	122	zinc carbonate	229
of mild mercurous chloride and jalap	110	Prepared cacao (Cacao praepa- rata)	44
of rhubarb and magnesia, anisated	184	calamine	46
of rhubarb, compound	184	chalk	75
of talc, boro-salicylated (Pul- vis talci compositus)	12	suet	197
of talc, compound	12	Prickly ash bark (Xanthoxylum)	228
pancreatin compound	153	ash berries	228
peptonizing (Pulvis pancrea- tini compositus)	153	ash, fluidextract (Fluidextrac- tum xanthoxyli)	228
rubefacient spice	57	Protochloride of iron solution (Liquor ferri protochloridi)	92
seidlitz (Pulvis effervescens compositus)	172	of iron syrup (Syrupus ferri protochloridi)	92
soluble, antiseptic	231	of mercury (Hydrargyri chloridum mite)	109
Powdered compound extract of colocynth (Extractum colo- cynthis compositum)	70	Protolodide of mercury (Hy- drargyri iodidum flavum) ..	110
extract of aconite (Extractum aconiti)	15	Prune	174
extract of aloes (Extractum aloes)	20	Prunus virginiana	174
extract of cascara sagrada (Extractum cascarae sagra- dae)	57	Prussic acid, diluted (Acidum hydrocyanicum dilutum)	9
extract of cimicifuga (Extrac- tum cimicifugae)	63	Pulsatilla	174
extract of colchicum corm (Extractum colchici cormi) ..	69	tincture	175
extract of colocynth (Extrac- tum colocynthis)	70	Pulveres	175

	PAGE	PAGE
Pulvis		
See also under Powder.		
aloes et canellae.....	21	
antisepticus	231	
antisepticus solubilis (Pulvis antisepticus)	231	
aromaticus rubefaciens	57	
catechu compositus (Pulvis gambir compositus)	98	
cretæ aromaticus	66	
cretæ compositus	76	
cretæ et opii aromaticus	151	
effervescens compositus	172	
gambir compositus	98	
glycyrrhizæ compositus	194	
hydrargyri chloridi mitis et jalapæ	110	
myricæ compositus	135	
potassii bromidi effervescens (Sal potassii bromidi effervescens)	170	
potassii bromidi effervescens cum caffèina (Sal potassii bromidi effervescens compositus)	170	
purgans (Pulvis alapæ compositus)	120	
salis carolini factitii effervescens	207	
salis kissingensis factitii effervescens (Sal kissingense factitium effervescens)	202	
salis vichyani factitii effervescens (Sal vichyanum factitium effervescens)	200	
salis vichyani factitii effervescens cum lithio (Sal vichyanum factitium effervescens cum lithio)	200	
Pumice	175	
Pumpkin seed (Pepo)	155	
Pure extract of glycyrrhiza	102	
Purgans, pulvis (Pulvis jalapæ compositus)	120	
Purging cassia (Cassia fistula)	58	
Purified antidiphtheric serum	196	
antitetanic serum	196	
cotton	103	
infusorial earth (Terra silicea purificata)	219	
kieselguhr (Terra silicea purificata)	219	
petroleum benzin	39	
siliceous earth	219	
talc	217	
Pyrethrum	175	
tincture	175	
Pyrogallie acid (Pyrogallol)	175	
Pyrogallol	175	
Pyroxylin	175	
Quadruplex pills (Pilulæ ferri, quininae, aloes et nucis vomicæ)	94	
Quassia	176	
extract	176	
extract, powdered (Extractum quassiae)	176	
fluidextract	176	
tincture	176	
Quatuor pills (Pilulæ ferri, quininae, aloes et nucis vomicæ)	94	
Quebracho (Aspidosperma)	35	
fluidextract (Fluidextractum aspidospermatis)	35	
Queen's root (Stillingia)	209	
Quercus	176	
fluidextract	176	
Quevenne's iron (Ferrum reductionis)	95	
Quicklime (Calx)	52	
Quicksilver (Hydrargyrum)	112	
Quillaja	176	
tincture	177	
Quinidine	177	
syrup	177	
syrup, bitterless (Syrupus quinidinae)	177	
Quinine	177	
aloes, iron and nux vomica pills	94	
and urea chloride (Quininae et ureæ hydrochloridum)	178	
and urea hydrochloride	178	
bisulphate	177	
bromide (Quininae hydrobromidum)	178	
chloride (Quininae hydrochloridum)	179	
citrate and iron	88	
citrate and iron, soluble (Ferri et quininae citras)	88	
dihydrochloride	178	
ferric pyrophosphate and strychnine elixir	93	
glycerinophosphate (Quininae glycerophosphas)	178	
glycerophosphate	178	
hydrobromide	178	
hydrochloride	179	
hypophosphite	179	
iron and strychnine, syrup of the phosphates of	92	
oleate	177	
opium and digitalis pills	151	
salicylate	179	
strychnine and iron elixir	87	
strychnine and phosphates syrup	179	
strychnine, iron and arsenic pills, mild	95	

	PAGE		PAGE
Quinine—Continued		Resorcin—Continued	
strychnine, iron and arsenic		paste, Lassar's mild (Pasta	
pills, stronger	95	resorcinolis mitis)	182
sulphate	179	paste, Lassar's stronger	
tannate	181	(Pasta resorcinolis fortis)..	182
tannate troches	181	paste, strong	182
valerate	181	Rhamnus cathartica	182
valerate and strychnine elixir	181	cathartica fluidextract	183
Raspberries	186	cathartica syrup	183
Raspberry syrup	186	purshiana (Cascara sagrada).	57
Raw linseed oil (Oleum lini)...	143	Rhatany (Krameria)	122
Rectified empyroligneous oil of		Rheum	183
birch (Oleum betulæ empy-		Rhubarb	183
reumaticum rectificatum)...	139	and gentian tincture	185
oil of birch tar	139	and magnesia powder, anisated	184
oil of tar	145	and opium mixture, compound	150
oil of turpentine	147	and soda mixture (Mistura	
tar oil (Oleum picis liquidæ		rhei composita)	183
rectificatum)	145	extract	183
turpentine oil (Oleum tere-		extract, powdered (Extractum	
binthinae rectificatum)	147	rhei)	183
Red aromatic elixir	19	fluidextract	183
cinchona	65	fluidglycerate	183
clover blossoms (Trifolium)...	222	mixture, alkaline	183
elixir (Elixir aromaticum		mixture, compound	183
rubrum)	19	pills, compound	184
indigo (Persio)	157	powder, compound	184
iodide of mercury (Hydrar-		syrup	184
gyri iodium rubrum)...	110	Rhubarb syrup, aromatic	184
lead (Plumbi oxidum rubrum)	166	syrup, spiced (Syrupus rhei	
mercuric iodide	110	aromaticus)	184
mercuric oxide	111	tincture	184
mercuric oxide, ointment	111	tincture, aqueous	184
oxide of lead	166	tincture, aromatic	184
peruvian bark (Cinchona		tincture, sweet	185
rubra)	65	wine, compound	185
precipitate (Hydrargyri oxi-		Rhus glabra	185
dum rubrum)	111	glabra, fluidextract	185
rose	186	roasted coffee (Coffea tosta)...	69
Saunders	189	Rochelle salt (Potassii et sodii	
Reduced iron	95	tartrate)	171
Refined and concentrated diph-		Rock-rose (Helianthemum)	107
theria antitoxin (Serum		Root (Bryonia)	44
antidiphthericum purifica-		Rosa centifolia (Rosa)	185
tum)	196	gallica	186
and concentrated tetanus anti-		Rose	185
toxin (Serum antitetanicum		confection	185
purificatum)	196	fluidextract	186
Rennin	181	honey	132
and pepsin elixir, compound	156	honey and sodium borate	200
Replacing elixir curassao (Elixir		honey with borax (Mel rosae	
aurantii amari)	138	et sodii boratis)	200
emplastrum aromaticum (Pul-		infusion, compound	186
vis aromaticus rubefaciens)	57	syrup	186
Resin of jalap	120	water	185
of podophyllum	167	water ointment	186
of scammony	192	water, stronger	185
Resins	182	Rosemary oil	146
Resorcin (Resorcinol)	182	Rosin	181
resorcinol	182	adhesive plaster (Emplastrum	
ointment, compound	182	resinae)	182
		cerate	181

PAGE	PAGE
Rosin—Continued	Salicylic acid
cerate compound	acid glycerogelatin
plaster	acid mull
Rubber adhesive plaster (Em-	acid pencil
plastrum elasticum)	collodion, compound
plaster	Salol (Phenylis salicylas).....
Rubefacient spice powder	Salt, artificial carlsbad
Rubi fructus	artificial Carlsbad, effervescent
Idael fructus	Glauber's (Sodii sulphas) ...
Rubus	of lithium citrate, effervescent
fluidextract	of potassium bromide, effervescent
syrup	of potassium bromide, effervescent, compound
Rumex	rochelle (Potassii et sodii tartras)
fluidextract	solution, normal (Liquor sodii chloridi physiologicus)
Russian flies (Cantharis).....	solution, physiological (Liquor sodii chloridi physiologicus)
Rye, ergot	Saltpetre (Potassi nitrates)
spurred	Salts, granular effervescent
Sabal	Salve, Deshler's (Ceratum resinae compositum)
fluidextract	mother's (Unguentum fuscum)
Saccharated ferric oxide	Sambucus
ferrous carbonate	Sandal wood
iron, soluble, syrup	oil (Oleum santali)
oxide of iron, syrup (Syrupus ferri saccharati solubilis) ..	white (Santalum album).....
pepsin	Sanguinaria
Saccharin (Benzosulphinidum) ..	fluidextract
soluble (Sodii benzosulphinidum) ..	syrup
Saccharum lactis	tincture
ustum (Caramel)	Santal and saw palmetto tincture
Saffron (Crocus)	oil
tincture	Santalum album
with opium tincture	rubrum
Saigon cinnamon	Santalwood oil (Oleum santali) ..
St. Germain tea (Species laxativae)	Santonin
St. Ignatius bean (Ignatia).....	and calomel troches (Trochisci santonini compositi) ..
St. John Long's liniment (Linimentum terebinthinae acetum) ..	troches
Sal carolinum factitium	troches, compound
carolinum factitium effervescent ..	Sapo
Kissingense factitium	mollis
Kissingense factitium effervescent ..	Sarsaparilla
lithii citratis effervescent	decoction, compound
potassii bromidi effervescent	fluidextract
potassii bromidi effervescent compositus ..	root (Sarsaparilla)
Vichyanum factitium	syrup, compound
Vichyanum factitium effervescent ..	Sassafras
Vichyanum factitium effervescent cum lithio	and opium mixture
Sales effervescentes	medulla
Salicin	oil
Salicylated creosote mull	pith
mixture of iron (Liquor ferri salicylatis)	pith, mucilage
	Saw palmetto and santal tincture
	palmetto berries (Sabal)
	palmetto, fluidextract (Fluidextractum sabal)

	PAGE		PAGE
Scammony, aloes and mercury pills, compound	21	Sirup (Syrupus)	188
resin	192	Skullcap (Scutellaria)	193
root	192	Slippery elm (Ulmus)	223
Scilla	192	Smallpox vaccine (Virus vaccini)	227
Scoparius	193	Smith's solution of bromine (Liquor bromi)	43
fluidextract	193	Smyrna galls (Galla)	98
Scopolamine bromide (Scopolamine hydrobromidum)	193	Snake-root, black (Cimicifuga)	63
hydrobromide	193	Canada (Asarum)	34
Scutellaria	193	black, fluidextract (Fluidextractum cimicifuga)	63
fluidextract	193	Texas (Serpentaria)	195
Secale cornutum (Ergota)	82	Virginia (Serpentaria)	195
Sedative water (Lotio ammoniacalis camphorata)	25	Soap	190
Selditz powder (Pulvis effervescens compositus)	172	green, tincture (Linimentum saponis molles)	190
Seneca snakeroot (Senega)	194	liniment	190
Senecio	193	liniment, camphorated	190
fluidextract	194	plaster	190
Senega	194	soft	190
fluidextract	194	soft, liniment	190
snakeroot (Senega)	194	soft, liniment, compound	190
syrup	194	white castile (Sapo)	190
Senna	194	Soap-tree bark (Quillaja)	176
Alexandria, or Indian senna (Senna)	194	Soda and mint solution	199
confection	194	and rhubarb mixture (Mistura rhei composita)	183
fluidextract	194	baking (Sodii bicarbonas)	199
infusion, compound	194	caustic (Sodii hydroxidum)	203
India, or Alexandria senna (Senna)	194	chlorinata	197
syrup	195	cum calce	198
syrup, aromatic	195	mint (Liquor sodae et Menthae)	199
syrup, compound	195	solution (Liquor sodii hydroxidi)	204
Serpentaria	195	with lime	198
tincture	195	Sodio-benzoate, caffeine	45
Serum, antidiphtheric	196	Sodio-salicylate, caffeine	45
antidiphtheric, dried	196	Sodium acetate	198
antidiphtheric, purified	196	and calcium glycerophosphates	47
antitetanic	196	elixir	47
antitetanic, dried	197	and calcium hypophosphites	48
antitetanic, purified	196	syrup	198
fluidextract	195	arsenate	198
Sesame oil	147	arsenate, exsiccated	198
Seven-barks (Hydrangea)	108	arsenate solution	198
Sevum benzoïnatum	197	arsenate solution, Pearson's	198
praeparatum	197	benzoate	198
Sherry wine	227	benzosulphide	199
Short buchu	44	bicarbonate	199
Siliceous earth, purified	219	bicarbonate troches	200
Silver nitrate	32	borate	200
nitrate, fused (Argenti nitras fusus)	33	borate and rose honey	200
nitrate, molded	33	borate honey	200
oxide	33	borate solution, compound	200
Simple cerate (Ceratum)	59	boro-benzoate	201
elixir (Elixir aromaticum)	18	bromide	201
ointment (Unguentum)	15	bromide elixir	201
syrup (Syrupus)	183	cacodylate	201
<i>Sinapis alba</i>	197	carbonate, monohydrated	201
<i> nigra</i>	197	chloride	201

PAGE	PAGE
Sodium—Continued	Soluble antiseptic powder..... 231
chloride and gold 37	cocoa (Cacao praeparata).... 44
chloride, physiological solu- tion 202	ferric citrate (Ferri et am- monii citras) 87
citrate 202	ferric oxide (Ferri oxidum saccharatum) 91
citrate and manganese (Man- gani citras solubilis)..... 130	ferric phosphate (Ferric phos- phas) 91
citrate solution 202	ferric pyrophosphate (Ferri pyrophosphas) 92
citro-tartrate, effervescent so- lution 202	gun cotton (Pyroxylinum).... 175
cyanide 202	iron and quinine citrate (Ferri et quininae citras) 88
glycerinophosphate (Sodii gly- cerophosphas) 203	manganese citrate 130
glycerophosphate 203	manganese glycerophosphate.. 130
glycerophosphate solution 203	manganous glycerinophosphate (Mangani glycerophosphas solubilis) 130
hydrate (Sodii hydroxidum).. 203	oxide of iron syrup (Syrupus ferri saccharati solubilis).. 91
hydroxide 203	saccharated iron syrup 91
hydroxide solution 204	saccharin (Sodii benzosul- phinidum) 199
hypophosphite 204	Solution
hypophosphite elixir 204	See also under Liqueur
hypophosphite syrup 204	antiseptic 7
hyposulphite (Sodii thiosul- phas) 207	antiseptic, alkaline 169
indigotindisulphonate 204	Boulton's (Liquor lodi phe- nolatus) 118
iodide 204	Burow's (Liquor aluminii acetatis) 23
nitrate 204	Channing's (Liquor hydrargyri et potassii iodidi) 110
perborate 205	coal tar 165
phenolsulphonate 205	deodorant (Liquor zinci et ferri compositus) 231
phosphate 205	Dobell's (Liquor Sodii boratis compositus) 200
phosphate, effervescent 206	Donovan's (Liquor arseni et hydrargyri iodidi) 33
phosphate, exsiccated 205	Fleming's (Liquor caelis sul- phuratae) 51
phosphate solution, compound	Fowler's (Liquor potassi ar- senitis) 168
pyroborate (Sodii boras) 200	hydrastine, colorless (Liquor hydrastinae compositus) ... 113
saccharin (Sodii benzosul- phinidum) 199	hydrochloric, of arsenic 34
salicylate 206	Labarraque's (Liquor sodae chlorinatae) 197
salicylate elixir 206	Lugol's (Liquor lodi composi- tus) 118
salicylate elixir, compound.. 206	Monse's (Liquor ferri subsul- phatis) 93
sulphate 206	normal salt (Liquor sodii chloridi physiologicus) 202
sulphite, exsiccated 207	of acid phosphates (Liquor phosphatum acidus) 11
sulphocarbonate (Sodii phenol- sulphonas) 205	of albuminate of iron..... 94
tartrate and bismuth..... 41	of aluminum acetate 23
tartrate and potassium..... 171	of aluminum acético-tartrate.. 24
tetraborate (Sodii boras) 200	of aluminum subacetate..... 24
thiosulphate 207	
Soft soap 190	
soap liniment 190	
soap liniment, compound..... 190	
zinc glycerogelatin 229	
zinc paste 229	
zinc paste, Unna's (Pasta zinci mollis) 229	
Solanum 207	
fluidextract 208	
Solid extracts (Extracta)..... 85	
opodeldoo (Linimentum sapo- nato-camporatum) 190	
petrox (Petroxolinum spis- sum) 159	
Solidified copaiba (Massa co- paibae) 73	

Solution—Continued	PAGE	Solution—Continued	PAGE
of ammonia, anisated (Spiritus ammoniac anisatus)	25	of magnesium sulphate, effervescent	129
of ammonium acetate	25	of mercuric nitrate	110
of ammonium citrate	27	of mercury and potassium iodide	110
of arsenic chloride (Liquor acidi arsenosi)	34	of oxysulphuret of calcium (Liquor calcis sulphuratae)	51
of arsenic, Clemens'	168	of pancreatin	153
of arsenous acid (Liquor acidi arsenosi)	34	of pepsin	157
of arsenous and mercuric iodide	33	of pepsin, antiseptic	157
of basic ferric sulphate (Liquor subsulphatis)	93	of pepsin, aromatic	157
of bismuth	42	of peptonate of iron	94
of bromine	43	of peptonate of iron and manganese	95
of bromine, Smith's (Liquor bromi)	43	of phosphates, acid	11
of calcium hydroxide	52	of phosphates, compound	11
of carmine	56	of phosphorus	161
of chlorinated potassa	167	of pituitary body (Liquor hypophysis)	116
of chlorinated soda	197	of potassa (Liquor potassii hydroxidi)	172
of chlorine, compound	62	of potassium arsenate and bromides (Liquor arsenicallis Clemens)	168
of cresol, compound	75	of potassium citrate	171
of ferric acetate	85	of potassium hydroxide	172
of ferric chloride	86	of potassium iodo-hydrargyrate (Liquor hydrargyri et potassii iodidi)	110
of ferric citrate	87	of protochloride of iron (Liquor ferri protochloridi)	92
of ferric hypophosphite	89	of phosphorus, Thompson's (Liquor phosphori)	161
of ferric nitrate	91	of soda (Liquor sodii hydroxidi)	204
of ferric oxychloride	91	of soda and mint	199
of ferric oxysulphate	91	of sodium arsenate	198
of ferric salicylate	206	of sodium arsenate, Pearson's	198
of ferric subsulphate	93	of sodium borate, compound	200
of ferric sulphate	94	of sodium chloride, physiological	202
of ferrous chloride	92	of sodium citrate	202
of formaldehyde	96	of sodium citro-tartrate, effervescent	202
of glycerophosphates, compound (Elixir glycerophosphatum compositum)	203	of sodium glycerophosphate	203
of gold and arsenic bromide	7	of sodium hydroxide	204
of gutta percha	106	of sodium phosphate, compound	205
of hydrastine, compound	113	of strychnine acetate	212
of hydrogen dioxide	115	of strychnine, Hall's (Liquor strychninae acetatis)	212
of hydrogen peroxide (Liquor hydrogenii dioxidi)	115	of sulphurated lime	51
of hypophosphite of iron (Liquor ferri hypophosphitis)	89	of tar, alkaline	165
of hypophosphites	48	of zinc and aluminum, compound	230
of hypophosphites, compound	48	of zinc and iron, compound	231
of hypophysis	116	of zinc chloride	229
of iodine, carbolyzed (Liquor iodi phenolatus)	118	of physiological salt (Liquor sodii chloridi physiologicus)	202
of iodine, compound	118	Vlemineckx' (Liquor calcis sulphuratae)	51
of iodine, phenolated	118		
of iron and ammonium acetate	25		
of iron perchloride (Liquor ferri chloridi)	86		
of iron tersulphate (Liquor ferri tersulphatis)	94		
of lead subacetate	167		
of lead subacetate, diluted	167		
of magnesium citrate	128		

	PAGE
Solutions	125
Spanish flies (<i>Cantharis</i>)	54
Sparteine sulphate	208
Spearmint	132
oil	143
spirit	143
water	143
Species	208
ad infusum pectorale (Species pectorales)	22
emollientes	23
laxativae	195
pectorales	22
Spermaceti	60
Spice powder, rubefacient.....	57
Spiced syrup of rhubarb (Syrupus rhei aromaticus) ..	184
Spigelia	208
fluidextract	208
Spigmet (<i>Aralia</i>)	32
Spirit of ammonia, anisated....	25
of anise	138
of ants (<i>Spiritus acidi formici</i>)	8
of bitter almond	138
of camphor	53
of cardamon, compound	140
of chloroform	62
of cinnamon	140
of ether	16
of ether, compound	16
of formic acid	8
of glonoin (<i>Spiritus glycerylis nitratis</i>)	102
of glyceryl trinitrate	102
of juniper	142
of juniper, compound	142
of lavender	142
of lavender, compound (<i>Tinctura lavandulae composita</i>) ..	142
of mindererus (<i>Liquor ammonii acetatis</i>)	25
of mustard	147
of myrcia, compound	145
of nitre, sweet (<i>Spiritus aetheris nitrosi</i>)	17
of nitroglycerin (<i>Spiritus glycerylis nitratis</i>)	102
of nitrous ether	17
of orange, compound	138
of peppermint	143
of spearmint	143
of vanillin, compound	224
Spirits	208
of turpentine (<i>Oleum terebinthinae</i>)	147
of volatile oils	208
<i>Spiritus amygdalae amarae</i> ..	138
<i>aurantii compositus</i>	138
<i>glycerylis nitratis</i>	102
<i>menthae piperitae</i>	143
<i>menthae viridis</i>	143

Spiritus—Continued

odoratus	139
sinapis	147
vanillin compositus	224
Spray, eucalyptol	83
menthol	133
menthol, compound	133
Sprays	136
Spurred rye (<i>Ergota</i>).....	82
Squaw root (<i>Caulophyllum</i>)....	59
Squibb's diarrhea mixture (<i>Mistura opii et chloroformi composita</i>)	150
Squill	192
fluidextract	192
mercury and digitalis pills...	79
oxymel	192
syrup	192
syrup, compound	192
tincture	192
vinegar	192
Squirrel corn (<i>Corydalis</i>)	74
Staphisagria	209
fluidextract	209
Star anise oil	138
grass (<i>Aletris</i>)	19
Starch	29
glycerite	29
Stavesacre (<i>Staphisagria</i>)	209
fluidextract (<i>Fluidextractum staphisagriae</i>)	209
Stearic acid	13
Sterilized distilled water.....	32
Still acid salicylic diluibles... 12	
diluibles	209
Stillingia	209
fluidextract	209
fluidextract, compound	209
syrup, compound	209
Stokes' expectorant	26
liniment (<i>Linimentum terebinthinae aceticum</i>)	148
Stomach drops (<i>Tinctura amara</i>)	100
Storax	214
liquid (<i>Styrax</i>)	214
Stramonium	210
extract	210
fluidextract	210
ointment	210
tincture	210
Strong resorcinol paste.....	182
Stronger ammonia water	25
orange flower water	37
pills of iron, quinine, strychnine and arsenic	95
resorcinol paste, Lassar's (<i>Pasta resorcinolis fortis</i>)..	182
rose water	185
tincture of iodine	119
Strontium bromide	210
carbonate	210

PAGE	PAGE
Strontium—Continued	
iodide	210
salicylate	211
Strophanthin	211
Strophanthus	211
tincture	211
Strychni semen (Nux vomica)..	136
tinctura (Tinctura nucis vomicae)	137
Strychnine	212
acetate	212
acetate solution	212
and quinine valerate elixir..	181
belladonna and aloin pills..	22
belladonna and aloin pills, compound	22
callaya alkaloidal, and iron elixir (Elixir cinchonae alkaloidorum, ferri et strychninae)	213
cinchona alkaloids, and iron elixir	213
cinchona alkaloids, iron and bismuth elixir	213
ferric pyrophosphate, and quinine, elixir	93
glycerinophosphate (Strychninae glycerophosphas) . .	212
glycerophosphate	212
iron and quinine elixir..	87
iron and quinine, syrup of the phosphates of	92
iron, quinine and arsenic pills, stronger	95
iron, callaya, alkaloidal, and bismuth, elixir (Elixir cinchonae alkaloidorum, ferri, bismuthi et strychninae)...	213
nitrate	213
pepsin, and bismuth elixir..	212
quinine and phosphates syrup	179
quinine, iron and arsenic pills, mild	95
solution, Hall's (Liquor strychninae acetatis)	212
sulphate	213
valerate	213
valerate elixir	214
Styptic collodion	13
cotton	86
Stypticin (Cotarninae hydrochloridum)	74
Styrax	214
Subchloride of mercury (Hydrargyri chloridum mite)..	109
Sublimed sulphur	215
Succus citri	214
citri et pepsinum	157
limettae cum pepsino (Succus citri et pepsinum)	157
pomorum	214
Sucrose (Saccharum)	187
Suet, benzoinated	197
mutton (Sevum praeeparatum) prepared	197
Sugar	187
of lead (Plumbi acetate)	165
of milk	188
Sulphocarbonate zinc (Zinci phenolsulphonas)	230
Sulphonal (Sulphonmethanum) .	214
Sulphonethylmethane	214
Sulphonmethane	214
Sulphur and cream of tartar troches (Trochisci sulphuris et potassii bitartratis)	215
and potassium bitartrate troches	215
flowers (Sulphur sublimatum) .	215
iodide	216
liver (Potassa sulphurata) . . .	168
lotum	215
milk (Sulphur praecipitatum) .	215
ointment	215
ointment, alkaline	215
ointment, compound	216
precipitated	215
sublimed	215
washed	215
Sulphurated antimony	30
lime (Calcii sulphidum crudum)	50
lime solution	51
petrox (Petroxolinum sulphuratum)	215
petrox, compound (Petroxolinum sulphuratum compositum)	215
potassa	168
zinc paste	230
zinc paste, Unna's (Pasta zinci sulphurata)	230
Sulphuric acid	13
aromatic	13
diluted	13
Sumac berries (Rhus glabra)...	185
Sumbul	216
extract	216
fluidextract	216
tincture	216
Sun cholera mixture (Mistura opii et rhei composita)	150
Sundew (Drosera)	80
Suppositories	216
of boroglycerin	101
of glycerin	101
Suprarenal glands, desiccated (Suprarenalium siccum)	216
dried	216
Sweet almond (Amygdala dulcis) almond oil (Oleum amygdalae expressum)	138
bark (Cascarilla)	53
birch oil (Methylis salicylas) .	133

	PAGE
Sweet—Continued	
clover, yellow (<i>Melilotus</i>)....	132
orange oil (<i>Oleum aurantii</i>)..	138
orange peel	36
orange peel, tincture	37
spirit of nitre (<i>Spiritus</i> <i>aetheris nitrosi</i>)	17
tincture of rhubarb	185
Sweetwood bark (<i>Cascarilla</i>) ..	58
Sydenham's laudanum (<i>Tinctura</i> <i>opii crocata</i>)	152
Syrup	188
hive (<i>Syrupus scillae composi-</i> <i>tus</i>)	192
Jackson's pectoral (<i>Syrupus</i> <i>morphinae et acaciae</i>)	134
of acacia	5
of actaea, compound (<i>Syrupus</i> <i>cimicifugae compositus</i>) ...	64
of althaea	22
of ammonium hypophosphite..	27
of asarum, compound	34
of blackberry fruit	186
of bloodroot (<i>Syrupus san-</i> <i>guinariæ</i>)	189
of buckthorn berries (<i>Syrupus</i> <i>rhamni cathartica</i>)	183
of calcium and sodium hypo- phosphites	48
of calcium hydrochlorophos- phate	50
of calcium hypophosphite ...	48
of calcium iodide	49
of calcium lactophosphate ...	50
of calcium lactophosphate and iron	50
of cimicifuga, compound	64
of cinnamon	66
of citric acid	8
of codiene	68
of Dover's powder (<i>Syrupus</i> <i>ipeacacuanhae et opii</i>)	151
of eriodictyon, aromatic	83
of ferric hypophosphite	89
of ferrous chloride	92
of ferrous iodide	90
of figs, compound	195
of garlic	19
of ginger	232
of glycyrrhiza	103
of hydriodic acid	9
of hydrochlorophosphates, com- pound (<i>Syrupus phosphat-</i> <i>um cum quina et strych-</i> <i>nina</i>)	179
of hypophosphites	48
of hypophosphites, compound..	49
of iodotannin	118
of ipecac	119
of ipecac and opium	151
of iron and manganese iodide	90
of iron lactophosphate	90

	PAGE
Syrup—Continued	
of krameria	123
of lactucarium	123
of licorice (<i>Syrupus glycyrr-</i> <i>rhizae</i>)	103
of manna	131
of morphine and acacia	134
of orange	36
of orange flowers	37
of phosphates, compound	11
of phosphates with quinine and strychnine	179
of poppy	153
of protochloride of iron (<i>Syrupus ferri protochlor-</i> <i>idi</i>)	92
of quinine	177
of quinidine, bitterless (<i>Syr-</i> <i>upus quinidinae</i>)	177
of raspberry	186
of rhamnus cathartica	183
of rhubarb, aromatic	184
of rhubarb, spiced (<i>Syrupus</i> <i>rhei aromaticus</i>)	184
of rose	186
of rubus	187
of saccharated oxide of iron (<i>Syrupus ferri saccharati</i> <i>solubilis</i>)	91
of sanguinaria	189
of sarsaparilla, compound ...	191
of senega	194
of senna	195
of senna, compound	195
of sodium hypophosphite	204
of soluble oxide of iron (<i>Syr-</i> <i>upus ferri saccharati solu-</i> <i>bilis</i>)	91
of soluble saccharated iron..	91
of squill	192
of squill, compound	192
of stillingia, compound	209
of tar	165
of the bromides	170
of the phosphates of iron, qui- nine and strychnine	92
of tolu	37
of white pine, compound	164
of white pine with morphine, compound	135
of wild cherry	174
of yerba santa, aromatic (<i>Syrupus eriodictyi aro-</i> <i>maticus</i>)	83
simple (<i>Syrupus</i>)	188
Syrups	217
Syrups	
See also under Syrup.	
alli	19
aurantii	36
aurantii florum	37
bromidorum	170

	PAGE		PAGE
Syrupus—Continued		Tar—Continued	
corrigens (Syrupus eriodictyli aromaticus)	83	solution, alkaline	165
ferri oxydati solubilis (Syrupus ferri saccharati solubilis)	91	solution, coal	165
ferri protochloridi	92	syrup	165
ferri, quinae et strychninae phosphatum	92	wine	165
ferri saccharati solubilis	91	Taraxacum	217
papaveris	153	elixir, compound	217
pectoralis (Syrupus morphinae et acaclae)	134	extract	218
phosphatum compositum	11	fluidextract	218
phosphatum cum quina et strychnina (See under Quinae hydrochloridum)	12	Tartar, cream of (Potassii bitartras)	169
phosphatum cum quinae et strychnina	179	cream of, and sulphur troches (Trochisci sulphuris et potassii bitartras)	215
pilis liquidae	165	emetic (Antimonii et potassii tartras)	30
pini strobili compositus	164	Tartaric acid	14
pini strobili compositus (Syrupus pini strobili compositus cum morphina)	135	Tartrated antimony (Antimonii et Potassii tartras)	30
pini strobili compositus cum morphina	135	Tasteless tincture of ferri chloride (Tinctura ferri citrochloridi)	87
pruni virginianae	174	tincture of iron (Tinctura ferri citro-chloridi)	87
rhei et potassii compositus (Mistura rhei alkalina)	183	Tea, breast (Species pectorales)	22
rubi	187	Teaberry, oil (Methylis salicylas)	133
rubi fructi	186	Teel oil (Oleum sesami)	147
rubi idaei	186	Terebene	218
scillae	192	Terebinthina	218
spinae cervinae (Syrupus rhamni catharticae)	183	laricis	218
stillingiae compositus	209	Terpin hydrate	218
zingiberis	232	hydrate and codeine elixir	68
Syrupy glucose (Glucosum)	101	hydrate and diacetylmorphine elixir	78
Talc, boro-salicylated powder (Pulvis talci compositus) ..	12	hydrate elixir	218
powder, compound	12	Terra silicea purificata	219
purified	217	Tetanus antitoxic globulins (Serum antitetanicum purificatum)	196
Tamarind	217	antitoxin (Serum antitetanicum)	196
Tannic acid	13	antitoxin, concentrated (Serum antitetanicum purificatum)	196
glycerite	13	antitoxin, dried (Serum antitetanicum siccum)	197
ointment	14	antitoxin, refined and concentrated (Serum antitetanicum purificatum)	196
troches	13	Texas snakeroot (Serpentaria) ..	195
Tannin (Acidum tannicum) ..	13	Theine (Caffeina)	45
glycerite (Glyceritum acidii tannici)	13	Theobroma, oil	148
Tar	164	Theocin (Theophyllina)	219
birch, rectified oil	139	Theophylline	219
coal	165	Theobromine sodio-salicylate ..	219
glycerite	165	Thompson's solution of phosphorus (Liquor phosphori) ..	161
mixture (Mistura olei pilis) ..	145	Thomsonian number six (Tinctura capsici et myrrhae) ..	55
oil, mixture	145	Thoroughwort (Eupatorium) ..	84
oil, rectified (Oleum pilis liquidae rectificatum)	145	Three bromides elixir	25
ointment	165		
ointment, compound	146		
petrox (Petroxolinum pilis) ..	145		
pine, (Pix liquida)	164		

	PAGE
Thuja	219
fluidextract	220
Thyme	220
fluidextract	220
oil	148
Thymol	220
iodide	220
spray	220
Thyroid glands, desiccated (Thy- roideum sicum)	220
Thyroids, dried	220
Tinctura	
See also under Tincture.	
amara	100
aurantii amari	36
aurantii dulcis	37
cinchonae composita	65
coto (Tinctura paracoto)	154
croci	76
delphinii	78
ferri malatis crudi (Tinctura ferri pomata)	90
ferri pomata	90
gallae	98
humuli	108
limonis corticis	124
moschi	135
pectoralis	152
persionis	157
persionis composita	157
pimpinellae	164
rhei	184
rhei aquosa	184
rhei aromatica	184
rhei dulcis	185
rhei et gentianae	185
rhei vinosa (Vinum rhei com- positum)	185
sabal et santali	187
saponis viridis composita	190
zingiberis	232
Tincturae aetherea	221
herbarum recentium (Tinc- turae medicamentorum re- centium)	221
medicamentorum recentium	221
Tincture, antiperiodic (Tinctura antiperiodica)	177
antiperiodic, without aloes	178
aromatic	66
Bestuscheff's (Tinctura ferri chloridi aetherea)	87
bitter	100
of aconite	15
of aloes	21
of aloes and myrrh	21
of arnica	33
of asafetida	34
of belladonna leaves	38
of benzoin	39
of benzoin, compound	39
of bitter orange peel	36

	PAGE
Tincture—Continued	
of bloodroot (Tinctura san- guinariae)	189
of bryonia	44
of cactus grandiflorus	45
of calbar bean (Tinctura phys- ostigmatis)	162
of calendula	51
of calumba	51
of cannabis	54
of cantharides	55
of capsicum	55
of capsicum and myrrh	55
of caramel	55
of cardamon	56
of cardamon, compound	56
of cimicifuga	64
of cinchona	65
of cinchona, compound	65
of cinnamon	66
of cocculus indicus	67
of colchicum seed	70
of crude malate of iron (Tinc- tura ferri pomata)	90
of cubeb	76
of cudbear	157
of cudbear, compound	157
of digitalis	79
of Dover's powder (Tinctura ipacacuanhae et opii)	152
of ergot, ammoniated	82
of ferrated extract of apples	90
of ferric chloride	86
of ferric chloride, ethereal	87
of ferric chloride, tasteless (Tinctura ferri citro-chlor- idi)	87
of ferric citro-chloride	87
of ferric citro-chloride with gentian (Elixir gentianae et ferri)	87
of gambir, compound	98
of gelsemium	99
of gentian, compound	100
of ginger	232
of golden seal (Tinctura hy- drastis)	114
of green soap (Linimentum saponis mollis)	190
of gualac	105
of gualac, ammoniated	105
of gualac, compound	105
of gualac, Dewees' (Tinctura gualaci composita)	105
of henbane (Tinctura hyoscy- ami)	115
of hops	108
of hydrastis	114
of hyoscyamus	115
of ignatia	115
of iodine	119

	PAGE		PAGE
Tincture—Continued		Tincture—Continued	
of iodine, Churchill's Tinc- tura Iodi fortior)	119	of viburnum, compound	226
of iodine, decolorized	118	of zedoary, bitter	228
of iodine, stronger	119	of zedoary, compound (Tinc- tura zedoariae amara)	228
of ipecac and opium	152	pectoral	152
of iron, tasteless (Tinctura ferri citro-chloridi)	87	stomachic (Tinctura amara)	100
of jalap	120	Warburg's (Tinctura antiperi- odica)	177
of jalap, compound	120	Warburg's, without aloes (Tinctura antiperiodica sine aloe)	178
of Jamaica ginger (Tinctura zingiberis)	232	Tinctures	221
of kino	122	ethereal	221
of kino and opium, compound	122	of fresh drugs	221
of kino, compound (Tinctura kino et opii composita)	122	Tobacco, Indian (Lobelia)	126
of krameria	123	Tolu balsam (Balsamum Tolu- tanum)	37
of lactucarium	124	syrup	37
of larkspur	78	tincture	38
of lavender, compound	142	Tonquin musk (Moschus)	135
of lemon peel	124	Toxibellae	221
of lobelia	127	hydrargyri chloridi corrosivi	109
of musk	135	Tragacanth	221
of myrrh	136	glycerite	221
of nutgall	98	muellage	221
of nux vomica	137	Trichloroacetic acid	14
of opium	152	fluidextract	222
of opium, camphorated	152	Trillium	222
of opium, deodorized	152	fluidextract	222
of opium with saffron	152	Trinitrin (Glycerylls nitrates)	101
of pale catechu, compound (Tinctura gambir com- posita)	98	Trinitrophenol	222
of paracoto	154	Trional (Sulphonethylmetha- num)	214
of passion flower	155	Trioxymethylene (paraformalde- hydum)	154
of peillitory (Tinctura py- rethri)	175	Triplex pills (Pillulae aloes, hy- drargyri et podophylli)	21
of pimpinella	164	Francis' (Pillulae aloes, hy- drargyri et scammonii com- positae)	21
of physostigma	162	Triticum	222
of pulsatilla	175	fluidextract	222
of pyrethrum	175	Trituration of elaterin	81
of quassia	176	Triturations	222
of quillaja	177	Troches	223
of rhubarb	184	of ammonium chloride	27
of rhubarb and gentian	185	of charcoal	56
of rhubarb, aqueous	184	of cubeb	76
of rhubarb, aromatic	184	of elm	223
of rhubarb, sweet	185	of gambir	99
of saffron	76	of peppermint	143
of sanguinaria	189	of phenolphthalein	166
of saw palmetto and santal	187	of potassium chlorate	171
of serpentaria	195	of quinine tannate	181
of squill	192	of santonin	189
of stramonium	210	of santonin and calomel (Tro- chisci santonini compositi)	190
of strophanthus	211	of santonin, compound	190
of sumbul	216	of sodium bicarbonate	200
of sweet orange peel	37		
of tolu	38		
of valerian	224		
of valerian, ammoniated	224		
of vanilla	224		
of veratrum viride	225		

PAGE	PAGE
Troches—Continued	Unna's paste—Continued
of sulphur and cream of tar- tar (Trochisci sulphuris et potassii bitartratis)	sulphurated zinc paste (Pasta zinci sulphurata)
215	230
of sulphur and potassium bi- tartrate	Uranium nitrate
215	223
of tannic acid	Uranil nitrate (Uranil nitras). 223
13	Urea chloride and quinine (Qui- ninae et ureae hydrochlori- dum)
223	178
Trochisci	hydrochloride and quinine
carbonis ligni	178
menthae piperitae	Urethane (Aethylis carbamas).. 17
ulmi	223
223	Urisol (Hexamethylenamina)
Turkey corn (Corydalis)	107
74	Uritone (Hexamethylenamina)
Turnera (Damiana)	107
77	Urotropine (Hexamethylen- amina)
Turnerae, fluidextractum (Fluid- extractum damianae)	107
77	Uva ursi
Turner's cerate (Unguentum calaminae)	223
46	fluidextract
218	223
Turpentine	Vaccine, Jennerian (Virus vac- cinicum)
218	227
larch (Terebinthina laricis)	smallpox (Virus vaccenicum)
148	227
liniment	virus
148	227
oil (Oleum Terebinthinae)	virus, glycerinated (Virus vaccinicum)
147	227
oil, emulsion	224
148	Valerian
oil, rectified (Oleum tere- binthinae rectificatum)	224
147	fluidextract
petrox, Venice (Petroxolinum terebinthinae laricis)	224
218	tincture
218	224
petroxolin, Venice	tincture, ammoniated
147	224
spirits (Oleum terebinthinae). Venice	Vallet's mass (Massa ferri car- bonatis)
218	85
Tussilago leaves (Farfara)	224
85	Vanilla
223	bean (Vanilla)
Ulmus	224
Unguentum	tincture
See also under Ointment.	224
creosoti salicylatum extensum (Mulla creosoti salicylata)	224
12	Vanillin
fuscum	elixir, compound
53	224
gallae	spirit, compound
98	224
hydrargyri	Vegetable cathartic pills
112	71
hydrargyri ammoniati	Venice turpentine
113	218
hydrargyri chloridi corrosivi extensum (Mulla hydrargyri chloridi corrosivi)	turpentine petrox (Petroxoli- num terebinthinae laricis)
109	218
hydrargyri dilutum	turpentine petroxolin
112	218
hydrargyri nitratis	Veratria (Veratrina)
111	225
hydrargyri oxidi flavi	Veratrine
111	225
hydrargyri oxidi rubri	ointment
111	225
matris (Unguentum fuscum)	oleate
53	225
piceis compositum	Veratrum viride
146	225
piceis liquidae	viride fluidextract
165	225
salicylatum extensum (Mulla acidi salicylici)	viride tincture
12	225
zinci carbonatis crudi (Un- guentum calaminae)	Verbasel flores
46	225
zinci extensum (Mulla zinci). 229	Verbena
230	226
zinci oxidi	fluidextract
230	226
zinci stearatis	Vermuth (Absinthium)
19	5
Unicorn root (Aletris)	Vesicating collodion (Collodium cantharidatum)
209	54
Unna pencils (Stilli dilubiles)	Viburnum (Viburnum pruni- folium)
Unna's soft zinc paste (Pasta zinci mollis)	227
229	opulus
	226
	opulus elixir, compound
	226
	opulus fluidextract
	226
	prunifolium
	227
	prunifolium elixir
	227
	prunifolium extract
	227

	PAGE		PAGE
Viburnum—Continued		Water—Continued	
prunifolium extract, powdered		chlorine (Liquor chlori com-	
(Extractum viburni pruni-		positus)	62
folii)	227	chloroform	62
prunifolium fluidextract	227	cinnamon	140
tincture, compound	226	creosote	75
Vichy salt, artificial	199	distilled	32
artificial, effervescent	200	distilled, sterilized	32
with lithium, effervescent		fennel	141
artificial	200	hamamelis	106
Villate's mixture (Mistura ad-		javelle (Liquor potassae	
stringens)	77	chlorinatae)	167
Vina	227	lead (Liquor plumbi subaceta-	
Vinegar, aromatic	142	tis dilutus)	167
of opium	149	lime (Liquor calcis)	52
of squill	192	orange flower	37
Vinegars (Aceta)	5	orange flower, stronger	37
Vinum		peppermint	143
See also under Wine.		phenolated	159
aurantii compositum	36	rose	185
carnis	56	rose, ointment of	186
carnis et ferri	57	rose, stronger	185
fraxini	97	sedative (Lotio ammoniacalis	
piceis	165	camphorata)	25
pruni virginianae	174	spearmint	143
pruni virginianae ferratum	174	witch hazel (Aqua hamameli-	
xericum	227	dis)	106
Virginia snakeroot (Serpentaria)	195	Waters, aromatic	32
Viride, veratum	225	White agaric (Agaricus)	18
Virus, vaccine	227	arsenic (Arseni trioxidum)	33
vaccine, glycerinated (Virus		ash bark	97
vaccinicum)	227	ash, wine	97
Vlemineckx' lotion (Liquor calcis		castile soap (Sapo)	190
sulphuratae)	51	dextrin	78
solution (Liquor calcis sul-		lead (Plumbi carbonas)	166
phuratae)	51	mustard	197
Volatile liniment (Linimentum		oak bark (Quercus)	176
ammoniae)	25	petrolatum	158
oil of mustard	147	petroleum jelly (Petrolatum	
oils, spirits	208	album)	158
Wahoo bark (Euonymus)	84	pine bark	164
Walnut bark, white (Juglans) ..	120	pine syrup, compound	164
Warburg's pills (Pilulae anti-		pine with morphine syrup,	
periodicae)	180	compound	135
pills without aloes (Pilulae		precipitate (Hydrargyrum am-	
antiperiodicae sine aloes) ..	180	moniatum)	113
tincture (Tinctura antiperi-		precipitate ointment (Unguen-	
odica)	177	tum hydrargyri ammoniati) ..	113
tincture without aloes (Tinc-		sandal wood (Santalum al-	
tura antiperiodica sine aloes)	178	bum)	189
Washed sulphur	215	walnut bark (Juglans)	120
Washes (Lotiones)	127	wax	59
Water		Wild black cherry (Prunus vir-	
See also under Aqua.		giniana)	174
almond, bitter	137	chamomile (Matricaria)	131
ammonia	24	cherry	174
ammonia, stronger	25	cherry and cod liver oil emul-	
anise	138	sion	144
camphor	53	cherry, fluidextract	174
carbolic acid (Aqua phe-		cherry infusion	174
nolata)	159	cherry syrup	174
		cherry wine	174

INDEX

285

	PAGE		PAGE
Wild—Continued		Yellow—Continued	
cherry wine, ferrated	174	lotion	111
ginger (Asarum)	34	mercuric oxide	111
indigo root (baptisia)	38	mercuric oxide ointment	111
yam root (Dioscorea)	79	mercurous iodide	116
Wilkinson's ointment (Unguentum sulphuris compositum)	216	mustard (Sinapis alba)	197
Wine, iron and beef (Vinum carnis et ferri)	57	Peruvian bark (Cinchona)	64
of antimony	30	sweet clover (Melilotus)	132
of beef	56	wash (Lotio flava)	111
of beef and iron	57	wax	59
of citrate of iron (Vinum ferri)	88	Yerba santa (Eriodictyon)	83
of colchicum corm	69	elixir, aromatic (Elixir eriodictyi aromaticum)	83
of colchicum seed	70	syrup, aromatic (Syrupus eriodictyi aromaticus)	83
of ipecac	119	Zea	228
of iron	88	fluidextract	228
of iron, bitter	88	Zedoary	228
of orange, compound	36	tincture, bitter	228
of pepsin	157	tincture, compound (Tinctura zedoariae amara)	228
of rhubarb, compound	185	Zinc	231
of tar	165	acetate	229
of white ash	97	and aluminum solution, compound	230
of wild cherry	174	and iron solution, compound	231
of wild cherry, ferrated	174	carbonate, precipitated	229
pepsin (Vinum pepsini)	157	chloride	229
sherry	227	chloride, solution	229
Wines	227	glycerogelatin, firm	229
Wintergreen oil (Methylis salicylas)	133	glycerogelatin, soft	229
Witch hazel (Hamamelis)	106	mull	229
extract, distilled (Aqua hamamelidis)	106	ointment (Unguentum zinci oxidi)	230
leaves (Hamamelidis folia)	106	oxide	229
water (Aqua hamamelidis)	106	oxide ointment	230
Wood charcoal	55	paste	229
Wool fat (Adeps lanae)	15	paste, Lassar's (Pasta zinci)	229
fat, hydrous	16	paste, soft	229
Wormseed, American oil (Oleum chenopodii)	140	paste, sulphurated	230
Wormwood (Absinthium)	5	paste, Unna's soft (Pasta zinci mollis)	229
Xanthoxylum	228	paste, Unna's sulphurated (Pasta zinci sulphurata)	230
fluidextract	228	phenolsulphonate	230
fructus	228	stearate	230
Yeast, compressed	60	stearate ointment	230
Yellow cinchona (Cinchona)	64	sulphate	230
dock (Rumex)	187	sulphocarbolate (Zinci phenolsulphonas)	230
gentian root (Gentiana)	99	valerate	231
iodide of mercury (Hydrargyri iodidum flavum)	110	valerate elixir	231
jasmine root (Gelsemium)	99	valerianate (Zinci valerac)	231
jessamine (Gelsemium)	99	Zingiber	231

USEFUL DRUGS

A descriptive list of selected drugs, which has been prepared under the direction and supervision of the Council on Pharmacy and Chemistry of the American Medical Association.

This list was compiled for the purpose of aiding the movement toward the elimination of the mass of useless or superfluous drugs now in books on materia medica. It is particularly fitted to serve as a text for teachers of materia medica and therapeutics.

Many state medical examining boards are limiting their examinations on therapeutics to this list of drugs.

In this book the properties—pharmacologic action, therapeutic uses and dosage, and methods of administration—are briefly but practically discussed.

Cloth, flexible cover, $4\frac{1}{2}$ by $7\frac{1}{2}$ inches, 176 pages. Price, postpaid, 60 cents.

AMERICAN MEDICAL ASSOCIATION
535 North Dearborn Street
CHICAGO

NEW AND NONOFFICIAL REMEDIES

This is an official publication of the Council on Pharmacy and Chemistry of the American Medical Association. It contains descriptions of those preparations—proprietary and non-proprietary—that have been accepted by the Council after due investigation. The description of each preparation includes the chief facts physicians desire to know concerning composition, dosage, indications, cautions to be observed, etc.

The book also contains the rules of the Council which form the basis of the work.

Every physician who desires to further the cause of scientific prescribing, who is anxious to see this country purged of the blight of the nostrums, and who desires to aid in diminishing the domination of commercialism in therapeutics in this country, should have a copy of this book for ready reference. Especially should it be available when the physician is called on by "detail men" exploiting proprietary preparations.

Single copy, cloth cover, postpaid, \$1.50. Prices on quantities sent on application. A permanent subscription list is now established.

AMERICAN MEDICAL ASSOCIATION

**535 North Dearborn Street
CHICAGO**

REPORTS OF THE COUNCIL ON PHARMACY AND CHEMISTRY OF THE AMERICAN MEDICAL ASSOCIATION

These volumes contain the complete and unabridged reports of the Council's work from 1905 down to the present year. Published for the use of physicians, teachers and students who wish full information in regard to the Council's action on the hundreds of medicinal preparations submitted to it for consideration. Price, \$1.00 postpaid; cloth cover.

Vols. I (1905-1908) to XIV (1921), Inc. Numbers I, II, V, VI, VII and XI are out of print.

PUBLICATIONS OF THE PROPAGANDA DEPARTMENT OF THE AMERICAN MEDICAL ASSOCIATION

Propaganda for Reform in Proprietary Medicines Volume 2, 1922

This book is in the press. It will contain the material (similar to that in the first volume) which has appeared since 1916. It will be bound uniform in size and style with that of Volume 1. The price of the book will be announced later.

Nostrums and Quackery and Allied Matters Affecting the Public Health Volume 2, 1921

This 800-page book gives the results of analyses of "patent medicines," describes the methods of "patent medicine" exploiters, details the schemes of medical mail-order quacks and in general turns the searchlight of scientific investigation into the dark recesses of unscientific medical empiricism.

Profusely illustrated, green cloth, stamped in gold

Price, \$2.00 postpaid

AMERICAN MEDICAL ASSOCIATION
535 North Dearborn Street CHICAGO, ILLINOIS



LANE MEDICAL LIBRARY

—
This book should be returned on or before
the date last stamped below.

--	--	--

V141.2	Pharmacopoeia of the	
P56	United States.	
1921	Epitome.	97538

[illegible]

9-41-3M

